

## COHOMOLOGY GROUPS OF CIRCULAR UNITS IN $\mathbb{Z}_p$ -EXTENSIONS

JAE MOON KIM

ABSTRACT. Let  $k$  be a real abelian field such that the conductor of every nontrivial character belonging to  $k$  agrees with the conductor of  $k$ . Note that real quadratic fields satisfy this condition. For a prime  $p$ , let  $k_\infty$  be the  $\mathbb{Z}_p$ -extension of  $k$ .

The aim of this paper is to produce a set of generators of the Tate cohomology group  $\hat{H}^{-1}$  of the circular units of  $k_n$ , the  $n$ th layer of the  $\mathbb{Z}_p$ -extension of  $k$ , where  $p$  is an odd prime. This result generalizes some earlier works which treated the case when  $k$  is real quadratic field and used them to study  $\lambda$ -invariants of  $k$ .

### 1. Introduction

Let  $k$  be a real abelian field of conductor  $d$ . It is known that  $k$  admits a unique  $\mathbb{Z}_p$ -extension for each prime  $p$ , namely the basic  $\mathbb{Z}_p$ -extension  $k_\infty = \bigcup_{n \geq 0} k_n$ , where  $k_n = k\mathbb{Q}_n$ . Here  $\mathbb{Q}_n$  is the subfield of  $\mathbb{Q}(\zeta_{p^{n+1}})$  whose degree over  $\mathbb{Q}$  is  $p^n$  when  $p$  is odd. When  $p = 2$ , one can define  $\mathbb{Q}_n$  similarly with a minor modification. But we will restrict our attention only to odd primes to avoid complexity.

For each  $n \geq 0$ , let  $A_n$  be the Sylow  $p$ -subgroup of the ideal class group of  $k_n$ . Then it is well known([5],[10]) that there exist integers  $\lambda \geq 0$  and  $\mu$  such that  $\#A_n = p^{\lambda n + \mu}$  for  $n \gg 0$ . These constants  $\lambda$  and  $\mu$  are called Iwasawa invariants. One of the main topics in Iwasawa theory is to investigate these invariants. For instance, Greenberg conjectured([4]) that  $\lambda = 0$  and gave several examples supporting the conjecture. Recently, many more affirmative results on the conjecture are given in [2],[3] and [7] when  $k$  is real quadratic. In all of these works,

---

Received Aug 1, 2000.

1991 Mathematics Subject Classification: Primary 11R23; Secondary 11R18.

Key words and phrases: circular units,  $\mathbb{Z}_p$ -extension, Iwasawa invariant.

This paper was supported by research fund of Inha University,1999

analysis of cyclotomic units(circular units, in general) is essential because they are related with the class number via the index theorem of Sinnott([9]). In [7], for example, some new information on the  $\lambda$ -invariant of a real quadratic field  $k$  was obtained from a particular circular unit of  $k$  which generates a certain cohomology group.

The aim of this paper is to find generators of cohomology groups for a wider class of real abelian fields including real quadratic fields so that they could be used, hopefully, in studying  $\lambda$ -invariants. To be precise, let  $G_n = Gal(k_n/k_0)$  and  $C_n$  be the group of circular units of  $k_n$ . Recently, it is proved([8]) that the Tate cohomology group  $\hat{H}^{-1}(G_n, C_n)$  is isomorphic to  $(\mathbb{Z}/p^n\mathbb{Z})^l$ , where  $l$  is the number of prime ideals of  $k$  above  $p$ . In this paper, we will find explicit generators of  $\hat{H}^{-1}(G_n, C_n)$  when the field  $k$  satisfies the following assumption : For each divisor  $f$  of the conductor  $d$  of  $k$ ,  $k \cap \mathbb{Q}(\zeta_f)$  is either  $k$ (this happens when  $d = f$ ) or  $\mathbb{Q}$ . Note that above assumption is equivalent to saying that every nontrivial character belonging to  $k$  is of conductor  $d$ . Also note that this assumption is clearly satisfied when  $k$  is a real quadratic field or when the conductor  $d$  is a prime  $q$ . Slightly more generally, the assumption is satisfied when  $[k : \mathbb{Q}]$  is a prime. Thus the results of this paper generalize those in [6], where explicit generators of  $\hat{H}^{-1}(G_n, C_n)$  are given when  $d = q$  is a prime.

This paper is organized as follows. Next section is preliminary. We will review circular units defined by Sinnott briefly and set up notations for the subsequent section. In section 3, we introduce a theorem of Ennola on relations of cyclotomic units and prove several lemmas. And, finally, we exhibit explicit generators.

## 2. Preliminaries and notations

In this section, we briefly review the definitions of cyclotomic units and circular units and set up notations.

For each integer  $s > 0$ , we choose a primitive  $s$ th root of 1 so that  $\zeta_s^{\frac{s}{t}} = \zeta_t$  if  $t \mid s$ . Let  $n \not\equiv 2 \pmod{4}$  and  $V$  be the multiplicative subgroup of  $\mathbb{Q}(\zeta_n)^\times$  generated by  $\{\pm\zeta_n, 1 - \zeta_n^a \mid 1 \leq a \leq n-1\}$ . Define the group  $C$  of cyclotomic units of  $\mathbb{Q}(\zeta_n)$  by  $C = E \cap V$ , where  $E$  is the group of units of  $\mathbb{Q}(\zeta_n)$ . In general, for an abelian field  $M$ , Sinnott([10]) defines

the group of circular units of  $M$  as follows. For  $n > 2$ , let

$$M_n = M \cap \mathbb{Q}(\zeta_n) \text{ and } C_{M_n} = N_{\mathbb{Q}(\zeta_n)/M_n}(C_{\mathbb{Q}(\zeta_n)}),$$

where  $C_{\mathbb{Q}(\zeta_n)}$  is the group of cyclotomic units of  $\mathbb{Q}(\zeta_n)$ . Then the group  $C_M$  of circular units of  $M$  is defined to be the multiplicative subgroup of  $M^\times$  generated by  $C_{M_n}$  for all  $n > 2$  together with  $-1$ . Note that in this definition, we do not have to consider infinitely many  $n$ 's. Let  $d$  be the conductor of  $M$ . If  $(n, d) = 1$ , then  $M_n = M \cap \mathbb{Q}(\zeta_n) = \mathbb{Q}$ . Thus  $N_{\mathbb{Q}(\zeta_n)/M_n}(C_{\mathbb{Q}(\zeta_n)}) = \{1\}$ . If  $d \mid n$ , then  $M \subset \mathbb{Q}(\zeta_d) \subset \mathbb{Q}(\zeta_n)$ . So  $N_{\mathbb{Q}(\zeta_n)/M}(C_{\mathbb{Q}(\zeta_n)}) \subset N_{\mathbb{Q}(\zeta_d)/M}(C_{\mathbb{Q}(\zeta_d)})$ . Therefore in the definition of circular units it is enough to consider those  $n$ 's which divide the conductor of  $n$ .

As before,  $k$  is a real abelian field of conductor  $d$ . Let  $K = \mathbb{Q}(\zeta_d)$ ,  $F = \mathbb{Q}(\zeta_p)$  and  $K' = \mathbb{Q}(\zeta_{pd})$ . We denote their cyclotomic  $\mathbb{Z}_p$ -extension by  $K_\infty, F_\infty$  and  $K'_\infty$ , and their  $n$ th layers by  $K_n, F_n$  and  $K'_n$  respectively. Let  $\sigma$  be the topological generator of the Galois group  $\Gamma = Gal(K'_\infty/K')$  which maps  $\zeta_{p^n}$  to  $\zeta_{p^n}^{1+p}$  for all  $n \geq 1$ . Restrictions of  $\sigma$  to various subfields are also denoted by  $\sigma$ . Let  $k_{(p)}$  be the decomposition subfield of  $k$  for  $p$  and let  $\Delta = Gal(K/k)$ ,  $\bar{\Delta} = Gal(K/\mathbb{Q})$ ,  $\Delta_p = Gal(K/k_{(p)})$ ,  $\Delta_k = Gal(k/\mathbb{Q})$  and  $\Delta_{k,p} = Gal(k_{(p)}/\mathbb{Q})$ . Let  $[k : \mathbb{Q}] = m$  and  $[k_{(p)} : \mathbb{Q}] = l$ , so there are  $l$  prime ideals in  $k$  above  $p$ . Elements of  $\Delta, \bar{\Delta}$  or  $\Delta_p$  will be denoted by  $\tau$ 's, and those of  $\Delta_k$  and  $\Delta_{k,p}$  by  $\rho$ 's. The Frobenius automorphism of  $K$  for  $p$  or its restriction to  $k$  is denoted by  $\tau_p$ . Let  $R$  be the set of all roots of 1 in the ring of the  $p$ -adic integers, i.e.,  $R = \{\omega \in \mathbb{Z}_p \mid \omega^{p-1} = 1\}$ . Then  $R$  can be regarded as the Galois group  $Gal(F/\mathbb{Q})$  or any Galois group isomorphic to it such as  $Gal(F_n/\mathbb{Q}_n)$ . For  $m > n$ , let  $G_{m,n}$  be the Galois group  $Gal(K'_m/K'_n)$  and  $N_{m,n}$  the norm map  $N_{K'_m/K'_n}$  from  $K'_m$  to  $K'_n$ . We will abbreviate  $G_{m,0}$  and  $N_{m,0}$  by  $G_m$  and  $N_m$ , respectively.  $G_{m,n}$  will also mean the Galois groups  $Gal(k_m/k_n)$ ,  $Gal(F_m/F_n)$  and  $Gal(\mathbb{Q}_m/\mathbb{Q}_n)$ . Similarly,  $N_{m,n}$  will have various meanings. Finally we fix a generator  $\psi_n$  of the character group of  $Gal(\mathbb{Q}_n/\mathbb{Q})$  such that  $\psi_n(\sigma) = \zeta_{p^n}$ .

### 3. Main results

We begin this section with a theorem of Ennola on relations of cyclotomic units which plays a crucial role in our study.

**THEOREM.** (V. Ennola[1]). Suppose  $\delta = \prod_{1 \leq a < n} (1 - \zeta_n^a)^{x_a}$  is a root of 1 for some integers  $x_a$ . Then for every even character  $\chi \neq 1$  of conductor  $f$  belonging to  $\mathbb{Q}(\zeta_n)$ ,  $Y(\chi, \delta) = 0$ , where

$$Y(\chi, \delta) = \sum_{\substack{d \\ f|d|n}} \frac{1}{\phi(d)} T(\chi, d, \delta) \prod_{p|d} (1 - \bar{\chi}(p)),$$

$$\text{and } T(\chi, d, \delta) = \sum_{\substack{a=1 \\ (a,d)=1}}^{d-1} \chi(a) x_{\frac{n}{d}a}.$$

**LEMMA 1.** Let  $\chi \neq 1$  be an even character belonging to  $\mathbb{Q}(\zeta_n)$  and  $\delta_1, \delta_2, \delta$  be cyclotomic units in  $\mathbb{Q}(\zeta_n)$ . Then

- (i)  $Y(\chi, \delta_1 \delta_2) = Y(\chi, \delta_1) + Y(\chi, \delta_2)$ .
- (ii) If  $(\text{root of } 1) \times \delta_1 = (\text{root of } 1) \times \delta_2$ , then  $Y(\chi, \delta_1) = Y(\chi, \delta_2)$ .
- (iii) For any  $\sigma \in \text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ ,  $Y(\chi, \delta^\sigma) = \chi(\sigma) Y(\chi, \delta)$ .
- (iv)  $Y(\chi, \delta^{\sigma^{-1}}) = (\chi(\sigma) - 1) Y(\chi, \delta)$ .

The proof of Lemma 1 is immediate from the definition of  $Y$ .

Next, in the following lemma, we compute  $Y(\chi, \delta)$  for various cyclotomic units  $\delta$  for an even character  $\chi = \psi_n \gamma$ , where  $\gamma$  is a character of conductor  $d$ .

**LEMMA 2.** Let  $\chi = \psi_n \gamma$  be an even character of conductor  $p^{n+1}d$ . Then

- (i)  $Y(\chi, \zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k) = \frac{1}{\phi(p^{n+1}d)} \psi_n(d\sigma^i) \gamma(kp^{n+1})$
  - (ii)  $Y(\chi, \prod_{i,j,k} (\zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k)^{c_{i,j,k}(\sigma^{-1})}) = \frac{(\psi_n(\sigma) - 1)}{\phi(p^{n+1}d)} \psi_n(d) \gamma(p^{n+1}) \alpha(\gamma)$
- for some algebraic integer  $\alpha(\gamma)$  depending on  $\gamma$ , where  $c_{i,j,k}$ 's are integers.

*Proof.* Note that  $\zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k = \zeta_d^k (\zeta_{p^{n+1}}^{\sigma^i \omega^j} \zeta_d^{-k} - 1) = \zeta_d^k (\zeta_{p^{n+1}d}^{\sigma^i \omega^j d - kp^{n+1}} - 1)$ .

Also note that  $\bar{\chi}(q) = 0$  for each  $q|p^{n+1}d$  since  $\chi$  is of conductor  $p^{n+1}d$ .

Thus

$$\begin{aligned} Y(\chi, \zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k) &= \frac{1}{\phi(p^{n+1}d)} T(\chi, p^{n+1}d, 1 - \zeta_{p^{n+1}d}^{\sigma^i \omega^j d - kp^{n+1}}) \\ &= \frac{1}{\phi(p^{n+1}d)} \psi_n \gamma(\sigma^i \omega^j d - kp^{n+1}) \\ &= \frac{1}{\phi(p^{n+1}d)} \psi_n(d\sigma^i) \gamma(kp^{n+1}). \end{aligned}$$

This proves (i).

For (ii), we use Lemma 1. From Lemma 1, we have

$$\begin{aligned} Y(\chi, \prod_{i,j,k} (\zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k)^{c_{i,j,k}(\sigma-1)}) \\ &= (\psi_n(\sigma) - 1) \sum_{i,j,k} c_{i,j,k} Y(\chi, \zeta_{p^{n+1}}^{\sigma^i \omega^j} - \zeta_d^k) \\ &= (\psi_n(\sigma) - 1) \sum_{i,j,k} c_{i,j,k} \frac{1}{\phi(p^{n+1}d)} \psi_n(d\sigma^i) \gamma(kp^{n+1}) \\ &= \frac{(\psi_n(\sigma) - 1)}{\phi(p^{n+1}d)} \psi_n(d) \gamma(p^{n+1}) \sum_{i,j,k} c_{i,j,k} \psi_n(\sigma^i) \gamma(k) \\ &= \frac{(\psi_n(\sigma) - 1)}{\phi(p^{n+1}d)} \psi_n(d) \gamma(p^{n+1}) \alpha(\gamma), \end{aligned}$$

where  $\alpha(\gamma) = \sum_{i,j,k} c_{i,j,k} \psi_n(\sigma^i) \gamma(k)$ . □

LEMMA 3. Let  $\delta_n = \prod_{\omega \in R, \tau \in \Delta_p} (\zeta_{p^{n+1}}^\omega - \zeta_d^\tau)$ . Then

- (i)  $N_{n,n-1}(\delta_n) = \delta_{n-1}$
- (ii)  $N_n(\delta_n) = 1$ .

*Proof.* Note that  $1 - X^p = \prod_{0 \leq i \leq p-1} (\zeta_p^i - X)$ . By putting  $X = \zeta_d^\tau \zeta_{p^{n+1}}^{-\omega}$ , we obtain

$$\prod_{0 \leq i \leq p-1} (\zeta_p^i - \zeta_d^\tau \zeta_{p^{n+1}}^{-\omega}) = 1 - \zeta_d^{p\tau} \zeta_{p^n}^{-\omega}.$$

Hence

$$\begin{aligned}
N_{n,n-1}(\zeta_{p^{n+1}}^\omega - \zeta_d^\tau) &= \prod_{0 \leq i \leq p-1} (\zeta_{p^{n+1}}^{(1+ip^n)\omega} - \zeta_d^\tau) \\
&= \prod_{0 \leq i \leq p-1} (\zeta_{p^{n+1}}^\omega \zeta_p^{i\omega} - \zeta_d^\tau) \\
&= \zeta_{p^n}^\omega \prod_{0 \leq i \leq p-1} (\zeta_p^i - \zeta_{p^{n+1}}^{-\omega} \zeta_d^\tau) \\
&= \zeta_{p^n}^\omega (1 - \zeta_d^{p\tau} \zeta_{p^n}^{-\omega}) \\
&= \zeta_{p^n}^\omega - \zeta_d^{p\tau}.
\end{aligned}$$

Therefore

$$\begin{aligned}
N_{n,n-1}(\delta_n) &= \prod_{\omega \in R, \tau \in \Delta_p} N_{n,n-1}(\zeta_{p^{n+1}}^\omega - \zeta_d^\tau) \\
&= \prod_{\omega \in R, \tau \in \Delta_p} (\zeta_{p^n}^\omega - \zeta_d^{p\tau}) \\
&= \delta_{n-1},
\end{aligned}$$

since  $p$  permutes  $\Delta_p$ .

Similarly,

$$N_n(\delta_n) = \prod_{\omega \in R, \tau \in \Delta_p} (\zeta_p^\omega - \zeta_d^\tau).$$

Put  $X = \zeta_d^\tau$  in the identity  $1 - X^p = \prod_{0 \leq i \leq p-1} (\zeta_p^i - X)$  to obtain

$$\frac{1 - \zeta_d^{\tau p}}{1 - \zeta_d^\tau} = \prod_{\omega \in R} (\zeta_p^\omega - \zeta_d^\tau).$$

Then

$$N_n(\delta_n) = \prod_{\tau \in \Delta_p} \frac{1 - \zeta_d^{\tau p}}{1 - \zeta_d^\tau} = 1.$$

Therefore we get Lemma 3. □

Let  $\{\rho, \dots, \rho_{l-1}, \rho_l = id\}$  be a set of coset representatives of  $\bar{\Delta}/\Delta_p$ . For each  $1 \leq i \leq l-1$ , let

$$\delta_{n,i} = \prod_{\substack{\omega \in R \\ \tau \in \Delta_p}} (\zeta_{p^{n+1}}^\omega - \zeta_d^{\tau \rho_i}) \quad \text{and} \quad \pi_n = \prod_{\omega \in R} (\zeta_{p^{n+1}}^\omega - 1).$$

Then  $N_{n,n-1}(\delta_{n,i}) = \delta_{n-1,i}$  and  $N_n(\delta_{n,i}) = 1$  by Lemma 3. Also, obviously we have  $N_n(\pi_n^{\sigma^{-1}}) = 1$ . Thus we have  $l$  circular units of  $k_n$  whose norm to  $k$  equal 1. The next theorem confirms that there are indeed a set of generators of  $\hat{H}^{-1}(G_n, C_n)$ .

**THEOREM.** *Suppose that  $k \cap \mathbb{Q}(\zeta_f) = k$  or  $\mathbb{Q}$  for every  $f$ ,  $f \mid d$ . Then  $\hat{H}^{-1}(G_n, C_n)$  is generated by  $\{\delta_{n,1}, \dots, \delta_{n,l-1}, \pi_n^{\sigma^{-1}}\}$ .*

The proof of the theorem is almost identical with that of section 5 of [6]. The only difference is that the conductor of  $k$  in [6] is a prime  $q$ , while the conductor  $d$ , here, is arbitrary. Although the computations for the general case are more complicated previous lemmas and the next lemma take care of all the difficulties in generalizing the proof of [6] to arbitrary conductor. So we omit the proof.

**LEMMA 4.** *Suppose that  $k \cap \mathbb{Q}(\zeta_f) = k$  or  $\mathbb{Q}$  for every  $f$ ,  $f \mid d$ . Let  $u$  be a circular unit in  $k_n$ . Then  $u$  can be written as  $u = u_0 u_1 u_2 \cdots u_n$ , where  $u_0 \in C_0$  and for each  $1 \leq i \leq n$ ,  $u_i$  is of the form*

$$u_i = \left( \prod_{\substack{\omega \in R \\ \tau \in \Delta}} (\zeta_{p^{i+1}}^\omega - \zeta_d^\tau)^{\sum_{i, \rho_j \in \Delta_k} a_{i,j} \sigma^i \rho_j} \right) \left( \pi_i^{(\sigma^{-1}) \sum_i b_i \sigma^i} \right).$$

*Proof.* To describe circular units of  $k_n$ , it is enough to consider divisors of the conductor of  $k_n$ , which is  $p^{n+1}d$  by the remark after the definition of circular units. Thus if  $f \mid p^{n+1}d$ , then  $f = p^{i+1}d_1$ , for some  $i$  ( $-1 \leq i \leq n$ ) and  $d_1 \mid d$ . Since  $\mathbb{Q}(\zeta_f) \cap k_n = k_i$  or  $\mathbb{Q}_n$ . Hence  $u$  is of the form

$$u = (N_{\mathbb{Q}(\zeta_d)/k}(v_0))(N_{\mathbb{Q}(\zeta_{p^2d})/k_1}(v_1))(N_{\mathbb{Q}(\zeta_{p^2})/\mathbb{Q}_1}(\omega_1)) \cdots \\ \cdots (N_{\mathbb{Q}(\zeta_{p^{n+1}d})/k_n}(v_n))(N_{\mathbb{Q}(\zeta_{p^{n+1}})/\mathbb{Q}_n}(\omega_n)),$$

where  $v_0, v_1, \omega_1, \dots, v_n$  and  $\omega_n$  are cyclotomic units in  $\mathbb{Q}(\zeta_d)$ ,  $\mathbb{Q}(\zeta_{p^2d})$ ,  $\mathbb{Q}(\zeta_{p^2})$ ,  $\dots$ ,  $\mathbb{Q}(\zeta_{p^{n+1}d})$  and  $\mathbb{Q}(\zeta_{p^{n+1}})$ , respectively. Thus  $u$  is of the desired form.  $\square$

## References

1. V.Ennola, *On relations between cyclotomic units*, J. Number Theory **4** (1972), 236–247.
2. T.Fukuda and K.Komatsu, *A capitulation problem and Greenberg's conjecture of real quadratic fields*, Math. Comp. **65** (1996), 313–318.
3. T.Fukuda and H.Taya, *The Iwasawa  $\lambda$ -invariants of  $\mathbb{Z}_p$ -extensions of real quadratic fields*, Acta Arith. **69** (1995), 277–292.
4. R.Greenberg, *On the Iwasawa invariants of totally real number fields*, Amer. J. Math. **98** (1976), 263–284.
5. K.Iwasawa, *On  $\mathbb{Z}_l$ -extensions of algebraic number fields*, Ann. Math. **98** (1973), 246–326.
6. J.M.Kim, *Circular units in the  $\mathbb{Z}_p$ -extensions of real abelian fields of prime conductor*, Tohoku Math. J. (1995), 305–313.
7. J.M.Kim and S.I.Oh, *On the Iwasawa  $\lambda$ -invariants of real quadratic fields*, Acta Arith (to appear).
8. J.M.Kim and S.I.Oh, *Cohomology groups of circular units* (to appear).
9. W.Sinnott, *On the Stickelberger ideal and the circular units of an abelian field*, Invent. Math. **62** (1980), 181–234.
10. L.Washington, *Introduction to Cyclotomic Fields*, G.T.M.83, Springer-Verlag, New York, 1980.

Department of Mathematics  
 Inha University, Incheon, Korea  
*E-mail*: jmkim@math.inha.ac.kr