

## FUZZY R-CLUSTER AND FUZZY R-LIMIT POINTS

YONG CHAN KIM AND YOUNG SUN KIM

ABSTRACT. In this paper, we introduce the notions of fuzzy r-cluster and fuzzy r-limit points in smooth fuzzy topological spaces and investigate some of their properties.

### 1. Introduction and preliminaries

A.P. Sostak [11] introduced the smooth fuzzy topology as an extension of Chang's fuzzy topology [1]. It has been developed in many directions [2,5,6]. Pu and Liu [10] introduced the notions of fuzzy nets and Q-neighborhoods and established the convergence theory in fuzzy topological spaces. In 1994, Chen and Cheng [3] introduced the concepts of fuzzy cluster and fuzzy limit points in fuzzy topological spaces with respect to R-neighborhood instead of Q-neighborhood.

In this paper, we introduce the concept of fuzzy r-cluster and fuzzy r-limit points in a smooth fuzzy topological space as an extension of [10] and investigate some of their properties and give an example of those.

Throughout this paper, let  $X$  be a nonempty set,  $I = [0, 1]$  and  $I_0 = (0, 1]$ . A *fuzzy point*  $x_t$  for  $t \in I_0$  is an element of  $I^X$  such that, for  $y \in X$ ,

$$x_t(y) = \begin{cases} t & \text{if } y = x, \\ 0 & \text{if } y \neq x. \end{cases}$$

The set of all fuzzy points in  $X$  is denoted by  $Pt(X)$ . Let  $x_t \in Pt(X)$  and  $\lambda, \mu \in I^X$ .  $x_t \in \lambda$  iff  $t \leq \lambda(x)$  for  $x \in X$ .  $\lambda$  is *quasi-coincident* with  $\mu$ , denoted by  $\lambda q \mu$ , if there exists  $x \in X$  such that  $\lambda(x) + \mu(x) > 1$ . If  $\lambda$  is not quasi-coincident with  $\mu$ , we denote  $\lambda \bar{q} \mu$ . All the other notations and the other definitions are standard in fuzzy set theory.

---

Received December 7, 1999.

1991 Mathematics Subject Classification: 54A40.

Key words and phrases: fuzzy r-cluster points, fuzzy r-limit points.

DEFINITION 1.1 ([11]). A function  $\mathcal{T} : I^X \rightarrow I$  is called a *smooth fuzzy topology* on  $X$  if it satisfies the following conditions:

- (O1)  $\mathcal{T}(\tilde{0}) = \mathcal{T}(\tilde{1}) = 1$ , where  $\tilde{0}(x) = 0$  and  $\tilde{1}(x) = 1$  for all  $x \in X$ .
- (O2)  $\mathcal{T}(\mu_1 \wedge \mu_2) \geq \mathcal{T}(\mu_1) \wedge \mathcal{T}(\mu_2)$ , for any  $\mu_1, \mu_2 \in I^X$ .
- (O3)  $\mathcal{T}(\bigvee_{i \in \Gamma} \mu_i) \geq \bigwedge_{i \in \Gamma} \mathcal{T}(\mu_i)$ , for any  $\{\mu_i\}_{i \in \Gamma} \subset I^X$ .

The pair  $(X, \mathcal{T})$  is called a *smooth fuzzy topological space*.

THEOREM 1.2 ([2]). Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space. For each  $r \in I_0$  and  $\lambda \in I^X$ , we define a fuzzy closure operator  $C_{\mathcal{T}} : I^X \times I_0 \rightarrow I^X$  as follows:

$$C_{\mathcal{T}}(\lambda, r) = \bigwedge \{ \rho \in I^X \mid \lambda \leq \rho, \mathcal{T}(\tilde{1} - \rho) \geq r \}.$$

For  $\lambda, \mu \in I^X$  and  $r, s \in I_0$ , it satisfies the following properties:

- (1)  $C_{\mathcal{T}}(\tilde{0}, r) = \tilde{0}$ .
- (2)  $\lambda \leq C_{\mathcal{T}}(\lambda, r)$ .
- (3)  $C_{\mathcal{T}}(\lambda, r) \vee C_{\mathcal{T}}(\mu, r) = C_{\mathcal{T}}(\lambda \vee \mu, r)$ .
- (4)  $C_{\mathcal{T}}(\lambda, r) \leq C_{\mathcal{T}}(\lambda, s)$ , if  $r \leq s$ .
- (5)  $C_{\mathcal{T}}(C_{\mathcal{T}}(\lambda, r), r) = C_{\mathcal{T}}(\lambda, r)$ .

DEFINITION 1.3 ([7]). Let  $\lambda, \mu \in I^X$ . Define the *fuzzy quasi-difference* of  $\lambda$  and  $\mu$ , denoted by  $\lambda \setminus \mu$ , as

$$(\lambda \setminus \mu)(x) = \begin{cases} \lambda(x), & \text{if } \mu(x) = 0, \\ 0, & \text{if } \lambda(x) \geq \mu(x) > 0, \\ \lambda(x), & \text{if } \lambda(x) < \mu(x). \end{cases}$$

NOTATION 1.4. Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space,  $\mu \in I^X$ ,  $x_t \in Pt(X)$  and  $r \in I_0$ . We denote

$$\mathcal{N}(x_t, r) = \{ \mu \in I^X \mid x_t q \mu, \mathcal{T}(\mu) \geq r \}.$$

DEFINITION 1.5 ([7]). Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space,  $\lambda \in I^X$ ,  $x_t \in Pt(X)$  and  $r \in I_0$ .

- (1)  $x_t$  is called an *fuzzy  $r$ -adherent point* of  $\lambda$  if for every  $\mu \in \mathcal{N}(x_t, r)$ , we have  $\mu q \lambda$ .

- (2)  $x_t$  is called a *fuzzy r-accumulation point* of  $\lambda$  if for every  $\mu \in \mathcal{N}(x_t, r)$ , we have  $\mu q (\lambda \setminus x_t)$ .
- (3) Define the *fuzzy r-derived set* of  $\lambda$ , denote by  $D_{\mathcal{T}}(\lambda, r)$ , as

$$D_{\mathcal{T}}(\lambda, r) = \bigvee \{x_t \in Pt(X) \mid x_t \text{ is a fuzzy r-accumulation point of } \lambda\}.$$

**THEOREM 1.6** ([7]). *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space. For each  $\lambda \in I^X$  and  $r \in I_0$ , we have*

$$C_{\mathcal{T}}(\lambda, r) = \bigvee \{x_t \in Pt(X) \mid x_t \text{ is a fuzzy r-adherent point of } \lambda\}.$$

**THEOREM 1.7** ([7]). *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space. For  $\lambda, \mu \in I^X$  and  $r, s \in I_0$ , the following properties hold:*

- (1)  $D_{\mathcal{T}}(\lambda, r) \leq C_{\mathcal{T}}(\lambda, r)$ .
- (2)  $C_{\mathcal{T}}(\lambda, r) = \lambda \vee D_{\mathcal{T}}(\lambda, r)$ .
- (3)  $C_{\mathcal{T}}(\lambda, r) = \lambda$  iff  $D_{\mathcal{T}}(\lambda, r) \leq \lambda$ .
- (4) If  $r \leq s$ , then  $D_{\mathcal{T}}(\lambda, r) \leq D_{\mathcal{T}}(\lambda, s)$ .
- (5)  $D_{\mathcal{T}}(\lambda \vee \mu, r) \leq D_{\mathcal{T}}(\lambda, r) \vee D_{\mathcal{T}}(\mu, r)$ .

## 2. Fuzzy r-cluster points and r-limit points

**DEFINITION 2.1.** Let  $D$  be a directed set. A function  $S : D \rightarrow Pt(X)$  is called a *fuzzy net*. Let  $\lambda \in I^X$ . We say  $S$  is a *fuzzy net in*  $\lambda$  if  $S(n) \in \lambda$  for every  $n \in D$ .

Using Notation 1.4 , we can define the followings:

**DEFINITION 2.2.** Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space,  $\mu \in I^X$ ,  $x_t \in Pt(X)$  and  $r \in I_0$ .

- (1)  $x_t$  is called a *fuzzy r-cluster point* of  $S$ , denoted by  $S \overset{r}{\infty} x_t$ , if for every  $\mu \in \mathcal{N}(x_t, r)$ ,  $S$  is frequently quasi-coincident with  $\mu$ , i.e, for each  $n \in D$ , there exists  $n_0 \in D$  such that  $n_0 \geq n$  and  $S(n_0) q \mu$ .
- (2)  $x_t$  is called a *fuzzy r-limit point* of  $S$ , denoted by  $S \overset{r}{\rightarrow} x_t$ , if for every  $\mu \in \mathcal{N}(x_t, r)$ ,  $S$  is eventually quasi-coincident with  $\mu$ , i.e,

there exists  $n_0 \in D$  such that for each  $n \in D$  with  $n \geq n_0$ , we have  $S(n) q \mu$ . We denote

$$clu_{\mathcal{T}}(S, r) = \bigvee \{x_t \in Pt(X) \mid x_t \text{ is a fuzzy } r\text{-cluster point of } S\},$$

$$lim_{\mathcal{T}}(S, r) = \bigvee \{x_t \in Pt(X) \mid x_t \text{ is a fuzzy } r\text{-limit point of } S\}.$$

DEFINITION 2.3. Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space. Let  $S : D \rightarrow Pt(X)$  and  $W : E \rightarrow Pt(X)$  be two fuzzy nets.  $W$  is called a *subnet* of  $S$  if there exists a function  $N : E \rightarrow D$ , called by a *cofinal selection* on  $S$ , such that

- (1)  $W = S \circ N$ ;
- (2) For every  $n_0 \in D$ , there exists  $m_0 \in E$  such that  $N(m) \geq n_0$  for  $m \geq m_0$ .

THEOREM 2.4. Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space. Let  $S : D \rightarrow Pt(X)$  fuzzy net and  $W : E \rightarrow Pt(X)$  a subnet of  $S$ . For  $r, s \in I_0$ , the following properties hold:

- (1) If  $S \xrightarrow{r} x_t$ , then  $S \overset{r}{\infty} x_t$ .
- (2)  $lim_{\mathcal{T}}(S, r) \leq clu_{\mathcal{T}}(S, r)$ .
- (3) If  $S \overset{r}{\infty} x_t$  and  $x_t \geq x_s$ , then  $S \overset{r}{\infty} x_s$ .
- (4) If  $S \xrightarrow{r} x_t$  and  $x_t \geq x_s$ , then  $S \xrightarrow{r} x_s$ .
- (5)  $S \overset{r}{\infty} x_t$  iff  $x_t \in clu_{\mathcal{T}}(S, r)$ .
- (6)  $S \xrightarrow{r} x_t$  iff  $x_t \in lim_{\mathcal{T}}(S, r)$ .
- (7) If  $S \xrightarrow{r} x_t$ , then  $W \xrightarrow{r} x_t$ .
- (8)  $lim_{\mathcal{T}}(S, r) \leq lim_{\mathcal{T}}(W, r)$ .
- (9) If  $W \overset{r}{\infty} x_t$ , then  $S \overset{r}{\infty} x_t$ .
- (10)  $clu_{\mathcal{T}}(W, r) \leq clu_{\mathcal{T}}(S, r)$ .

*Proof.* (1) and (2) are clear.

(3) For every  $\mu \in \mathcal{N}(x_s, r)$ , since  $x_s \leq x_t$ , then  $\mu \in \mathcal{N}(x_t, r)$ . Since  $S \overset{r}{\infty} x_t$ , for each  $n \in D$ , there exists  $n_0 \in D$  such that  $n_0 \geq n$  and  $S(n_0) q \mu$ . Hence  $S \overset{r}{\infty} x_s$ .

(4) It is similar to (3).

(5) ( $\Rightarrow$ ) It is clear.

( $\Leftarrow$ ) Let  $x_t \in clu_{\mathcal{T}}(S, r)$  and  $\mu \in \mathcal{N}(x_t, r)$ . Since  $x_t q \mu$  and  $clu_{\mathcal{T}}(S, r)(x) \geq t$ , we have

$$\mu(x) + clu_{\mathcal{T}}(S, r)(x) \geq \mu(x) + t > 1.$$

From the definition of  $clu_{\mathcal{T}}(S, r)$ , there exists a fuzzy r-cluster point  $x_s \in Pt(X)$  of  $S$  such that

$$\mu(x) + clu_{\mathcal{T}}(S, r)(x) \geq \mu(x) + s > 1.$$

Thus  $\mu \in \mathcal{N}(x_s, r)$ . Since  $x_s$  is a fuzzy r-cluster point of  $S$ , for each  $n \in D$ , there exists  $n_0 \in D$  such that  $n_0 \geq n$  and  $S(n_0) q \mu$ . Hence  $S \overset{r}{\infty} x_t$ .

(6) It is similar to (5).

(7) For every  $\mu \in \mathcal{N}(x_t, r)$ , since  $S \overset{r}{\rightarrow} x_t$ , there exists  $n_0 \in E$  such that for all  $n \geq n_0$ ,  $S(n) q \mu$ . Let  $N : E \rightarrow D$  be a cofinal selection on  $S$ . Then for  $n_0 \in D$ , there exists  $m_0 \in E$  such that  $N(m) \geq n_0$  for all  $m \geq m_0$ . Thus  $W(m) = S(N(m)) q \mu$  for  $m \geq m_0$ . Therefore,  $W \overset{r}{\rightarrow} x_t$ .

(8) From (7), it is clear.

(9) Suppose that  $W \overset{r}{\infty} x_t$  and  $n \in D$ . If  $N : E \rightarrow D$  is a cofinal selection on  $S$ , then there exists  $m \in E$  such that  $N(k) \geq n$  for  $k \geq m$ . Since  $W \overset{r}{\infty} x_t$ , for every  $\mu \in \mathcal{N}(x_t, r)$ , there exists  $m_0 \in E$  such that  $m_0 \geq m$  and  $W(m_0) q \mu$ . We let  $n_0 = N(m_0)$ . Then  $n_0 \geq n$  and since  $S(n_0) = W(m_0)$ , we have  $S(n_0) q \mu$ .

(10) From (9), it is clear.  $\square$

**THEOREM 2.5.** *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space and  $x_t \in Pt(X)$  and  $r \in I_0$ . For every fuzzy net  $S$ ,  $S \overset{r}{\rightarrow} x_t$  iff  $W \overset{r}{\infty} x_t$ , for every fuzzy subnet  $W$  of  $S$ .*

*Proof.* ( $\Rightarrow$ ) From Theorem 2.4(7),  $S \overset{r}{\rightarrow} x_t$  implies  $W \overset{r}{\rightarrow} x_t$ . From Theorem 2.4 (9),  $W \overset{r}{\rightarrow} x_t$  implies  $W \overset{r}{\infty} x_t$ .

( $\Leftarrow$ ) Suppose  $x_t$  is not a fuzzy r-limit point  $x_t$  of  $S$ . Then there exists  $\mu \in \mathcal{N}(x_t, r)$  satisfying the followings: for each  $n \in D$ , there exists  $N(n) \in D$  such that  $N(n) \geq n$  and  $S(N(n)) \bar{q} \mu$ . We can define  $N : D \rightarrow D$ . For each  $m \geq n$ , we have  $N(m) \geq m \geq n$ . Hence  $N$  is a

cofinal selection on  $S$ . So,  $W = S \circ N$  is a fuzzy subnet of  $S$ . Since for  $\mu \in \mathcal{N}(x_t, r)$  and for each  $n \in D$ ,  $W(n) = S(N(n)) \bar{q} \mu$ ,  $x_t$  is not fuzzy  $r$ -cluster point of  $W$ .  $\square$

**THEOREM 2.6.** *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space and  $x_t \in Pt(X)$  and  $r \in I_0$ . For every fuzzy net  $S : D \rightarrow Pt(X)$ , we have  $S \overset{r}{\infty} x_t$  iff  $S$  has a fuzzy subnet  $W$  such that  $W \xrightarrow{r} x_t$ .*

*Proof.* ( $\Rightarrow$ ) Let  $E = D \times \mathcal{N}(x_t, r) = \{(m, \lambda) \mid m \in D, \lambda \in \mathcal{N}(x_t, r)\}$ . Define a relation on  $E$  by

$$\forall (m, \lambda), (n, \mu) \in E, \quad (m, \lambda) \leq (n, \mu) \Leftrightarrow m \leq n, \lambda \geq \mu.$$

For each  $(m, \lambda), (n, \mu) \in E$ , we have  $\lambda, \mu \in \mathcal{N}(x_t, r) \Rightarrow \lambda \wedge \mu \in \mathcal{N}(x_t, r)$  and there exists  $k \in D$  such that  $m \leq k$  and  $n \leq k$ . Hence there exists  $(k, \lambda \wedge \mu) \in E$  such that  $(m, \lambda) \leq (k, \lambda \wedge \mu)$  and  $(n, \mu) \leq (k, \lambda \wedge \mu)$ . So,  $E$  is a directed set. For each  $(n, \mu) \in E$ , since  $S \overset{r}{\infty} x_t$ , there exists  $N(n, \mu) \in D$  such that  $N(n, \mu) \geq n$  and  $S(N(n, \mu)) \bar{q} \mu$ . So, we can define  $N : E \rightarrow D$ . For each  $n_0 \in D$ , since  $S \overset{r}{\infty} x_t$ , for  $\mu_0 \in \mathcal{N}(x_t, r)$ , there exists  $(n_0, \mu_0) \in E$  such that  $N(n_0, \mu_0) \geq n_0$ . Hence for every  $(n, \mu) \geq (n_0, \mu_0)$ , since  $n \geq n_0$ , we have  $N(n, \mu) \geq n \geq n_0$ . Therefore  $N$  is a cofinal selection on  $S$ . So,  $W = S \circ N$  is a fuzzy subnet of  $S$ . Now we show that  $W \xrightarrow{r} x_t$ . For each  $\mu_0 \in \mathcal{N}(x_t, r)$ , since  $S \overset{r}{\infty} x_t$ , for  $n_0 \in D$ , there exists  $N(n_0, \mu_0) \in D$  such that  $S(N(n_0, \mu_0)) \bar{q} \mu_0$ . Hence for every  $(n, \mu) \geq (n_0, \mu_0)$ ,  $S(N(n, \mu)) \bar{q} \mu$  implies  $S(N(n, \mu)) \bar{q} \mu_0$  because  $\mu \leq \mu_0$ . So,  $W \xrightarrow{r} x_t$ .

( $\Leftarrow$ ) From Theorem 2.4(1),  $W \xrightarrow{r} x_t$  implies  $W \overset{r}{\infty} x_t$ . From Theorem 2.4(9),  $W \overset{r}{\infty} x_t$  implies  $S \overset{r}{\infty} x_t$ .  $\square$

**THEOREM 2.7.** *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space and  $x_t \in Pt(X)$  and  $r \in I_0$ . Then the following statements are equivalent.*

- (1)  $x_t \in C_{\mathcal{T}}(\lambda, r)$
- (2) There exists a fuzzy net  $S$  in  $\lambda$  such that  $S \overset{r}{\infty} x_t$ .
- (3) There exists a fuzzy net  $S$  in  $\lambda$  such that  $S \xrightarrow{r} x_t$ .

*Proof.* (1) $\Rightarrow$ (2) Define a relation on  $\mathcal{N}(x_t, r)$  by,

$$\nu \preceq \omega \text{ iff } \omega \leq \nu, \quad \forall \nu, \omega \in \mathcal{N}(x_t, r).$$

Then  $(\mathcal{N}(x_t, r), \preceq)$  is a directed set.

For each  $\mu \in \mathcal{N}(x_t, r)$ , since  $x_t \in C_{\mathcal{T}}(\lambda, r)$ , we have

$$C_{\mathcal{T}}(\lambda, r)(x) + \mu(x) \geq t + \mu(x) > 1.$$

From Theorem 1.6, there exists fuzzy r-adherent point  $x_s$  of  $\lambda$  such that

$$C_{\mathcal{T}}(\lambda, r)(x) + \mu(x) \geq s + \mu(x) > 1.$$

Since  $x_s$  is a fuzzy r-adherent point of  $\lambda$  and  $\mu \in \mathcal{N}(x_s, r)$ , we have  $\lambda q \mu$ . Then there exist  $y \in X$  and  $m \in I_0$  such that

$$\lambda(y) + \mu(y) \geq m + \mu(y) > 1.$$

Hence  $y_m \in \lambda$  and  $\mu \in \mathcal{N}(y_m, r)$ . Define a directed set  $(\mathcal{N}(x_t, r), \preceq)$  by

$$\nu \preceq \mu \text{ iff } \mu \leq \nu.$$

For each  $\mu \in \mathcal{N}(x_t, r)$ , we can define a fuzzy net  $S : \mathcal{N}(x_t, r) \rightarrow Pt(X)$  by  $S(\mu) = y_m$ . Then  $S(\mu) q \mu$  and  $S(\mu) \in \lambda$ .

Now we will show that  $S \overset{r}{\infty} x_t$ . Let  $\mu \in \mathcal{N}(x_t, r)$ . Then for every  $\nu \in \mathcal{N}(x_t, r)$ , we have  $\mu \wedge \nu \in \mathcal{N}(x_t, r)$  and  $S(\mu \wedge \nu) q (\mu \wedge \nu)$ . Thus  $\nu \preceq \mu \wedge \nu$  and  $S(\mu \wedge \nu) q \mu$ .

(2)  $\Rightarrow$  (1) If there exists a fuzzy net  $S$  in  $\lambda$  such that  $S \overset{r}{\infty} x_t$ , for each  $\mu \in \mathcal{N}(x_t, r)$  and for each  $n \in D$ , there exists  $n_0 \in D$  such that  $n_0 \geq n$  and  $S(n_0) q \mu$ . Since  $S(n_0) \in \lambda$ ,  $S(n_0) q \mu$  implies  $\lambda q \mu$ . Hence  $x_t$  is fuzzy r-adherent point of  $\lambda$ , that is,  $x_t \in C_{\mathcal{T}}(\lambda, r)$ .

(2) $\Rightarrow$ (3) It is easily proved from Theorem 2.6.

(3) $\Rightarrow$ (2) It is easily proved from Theorem 2.4(1).  $\square$

**THEOREM 2.8.** *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space and  $x_t \in Pt(X)$  and  $r \in I_0$ . Then the following statements are equivalent.*

(1)  $C_{\mathcal{T}}(\lambda, r) = \lambda$ .

(2) For every fuzzy net  $S$  in  $\lambda$  and  $x_t \in Pt(x)$ , if  $S \overset{r}{\infty} x_t$ , then  $x_t \in \lambda$ .

(3) For every fuzzy net  $S$  in  $\lambda$  and  $x_t \in Pt(x)$ , if  $S \overset{r}{\rightarrow} x_t$ , then  $x_t \in \lambda$ .

*Proof.* (1) $\Rightarrow$ (2) Suppose that there exists a fuzzy net  $S$  in  $\lambda$  such that  $S \overset{r}{\infty} x_t$  but  $x_t \notin \lambda$ . From Theorem 2.7,  $x_t \in C_{\mathcal{T}}(\lambda, r)$ . Hence  $C_{\mathcal{T}}(\lambda, r)(x) \geq t > \lambda(x)$ . Thus  $C_{\mathcal{T}}(\lambda, r) \neq \lambda$ .

(2)  $\Rightarrow$  (1) If  $x_t \in C_{\mathcal{T}}(\lambda, r)$ , by Theorem 2.7, there exists a fuzzy net  $S$  in  $\lambda$  such that  $S \overset{r}{\infty} x_t$ . Hence  $x_t \in \lambda$  from (2). Thus  $C_{\mathcal{T}}(\lambda, r) \leq \lambda$ . From Theorem 1.2(2), we have  $C_{\mathcal{T}}(\lambda, r) = \lambda$ .

(1) $\Rightarrow$ (3) and (3) $\Rightarrow$ (1) are similarly proved.  $\square$

Using Definition 1.5 and Theorem 2.7, we can easily prove the following corollary.

**COROLLARY 2.9.** *Let  $(X, \mathcal{T})$  be a smooth fuzzy topological space and  $x_t \in Pt(X)$  and  $r \in I_0$ . Then the following statements are equivalent.*

- (1)  $x_t \in D_{\mathcal{T}}(\lambda, r)$
- (2) There exists a fuzzy net  $S$  in  $\lambda \setminus x_t$  such that  $S \overset{r}{\infty} x_t$ .
- (3) There exists a fuzzy net  $S$  in  $\lambda \setminus x_t$  such that  $S \overset{r}{\rightarrow} x_t$ .

**EXAMPLE 2.10.** Let  $X = \{x, y\}$  be set. Define  $\mu \in I^X$  as follows:

$$\mu(x) = 0.3, \mu(y) = 0.4.$$

We define a smooth fuzzy topology  $\mathcal{T} : I^X \rightarrow I$  as follows:

$$\mathcal{T}(\lambda) = \begin{cases} 1, & \text{if } \lambda = \tilde{0} \text{ or } \tilde{1}, \\ \frac{1}{2}, & \text{if } \lambda = \mu, \\ 0, & \text{otherwise.} \end{cases}$$

Let  $N$  be a natural number set. Define a fuzzy net  $S : N \rightarrow Pt(X)$  by

$$S(n) = x_{a_n}, \quad a_n = 0.6 + (-1)^n 0.2 + \frac{0.2}{n}.$$

We can show  $clu_{\mathcal{T}}(S, \frac{1}{2}) = \tilde{1}$  from (1) to (4)

(1)  $x_t$  for  $t \leq 0.7$  is a fuzzy  $\frac{1}{2}$ -cluster point of  $S$ , for  $\tilde{1} \in \mathcal{N}(x_t, \frac{1}{2})$  and for all  $n \in N$ , we have  $S(n) q \tilde{1}$ .

(2)  $x_t$  for  $0.7 < t$  is a fuzzy  $\frac{1}{2}$ -cluster point of  $S$ , for  $\tilde{1}, \mu \in \mathcal{N}(x_t, \frac{1}{2})$  and for all  $n \in N$ , there exists  $2n \in N$  such that  $2n \geq n$ ,  $S(2n) = x_{0.8 + \frac{0.2}{2n}} q \mu$  and  $S(2n) = x_{0.8 + \frac{0.2}{2n}} q \tilde{1}$ .



(3)  $y_s$  for  $s \leq 0.6$  is a fuzzy  $\frac{1}{2}$ -cluster point of  $S$ , for  $\tilde{1} \in \mathcal{N}(y_s, \frac{1}{2})$  and for all  $n \in N$ , we have  $S(n) q \tilde{1}$ .

(4)  $y_s$  for  $0.6 < s$  is a fuzzy  $\frac{1}{2}$ -cluster point of  $S$ , for  $\tilde{1}, \mu \in \mathcal{N}(y_s, \frac{1}{2})$  and for all  $n \in N$ , there exists  $2n \in N$  such that  $2n \geq n$ ,  $S(2n) = x_{0.8+\frac{0.2}{n}} q \tilde{1}$  and  $S(2n) = x_{0.8+\frac{0.2}{n}} q \mu$ .

We can show  $\lim_{\mathcal{T}}(S, \frac{1}{2}) = \tilde{1} - \mu$  from (5) to (8).

(5)  $x_t$  for  $t \leq 0.7$  is a fuzzy  $\frac{1}{2}$ -limit point of  $S$ , for  $\tilde{1} \in \mathcal{N}(x_t, \frac{1}{2})$  and for all  $n \in N$ , we have  $S(n) q \tilde{1}$ .

(6)  $x_t$  for  $0.7 < t$  is not a fuzzy  $\frac{1}{2}$ -limit point of  $S$ , there exists  $\mu \in \mathcal{N}(x_t, \frac{1}{2})$  such that for all  $n \in N$ , there exists  $2n+1 \in N$  such that  $2n+1 \geq n$  and  $S(2n+1) = x_{0.4+\frac{0.2}{2n+1}} \bar{q} \mu$ .

(7)  $y_s$  for  $s \leq 0.6$  is a fuzzy  $\frac{1}{2}$ -limit point of  $S$ , for  $\tilde{1} \in \mathcal{N}(y_s, \frac{1}{2})$  and for all  $n \in N$ , we have  $S(n) q \tilde{1}$ .

(8)  $y_s$  for  $0.6 < s$  is not a fuzzy  $\frac{1}{2}$ -limit point of  $S$ , there exists  $\mu \in \mathcal{N}(x_t, \frac{1}{2})$  such that for all  $n \in N$ , there exists  $2n+1 \in N$  such that  $2n+1 \geq n$  and  $S(2n+1) = x_{0.4+\frac{0.2}{2n+1}} \bar{q} \mu$ . Hence  $\lim_{\mathcal{T}}(S, \frac{1}{2}) = \tilde{1} - \mu$ .

Define  $\Psi : N \rightarrow N$  by  $\Psi(n) = 2n+1$ . Then  $\Psi$  is a cofinal selection on  $S$ .  $W$  is a subnet of  $S$ . Since  $W(n) = S \circ \Psi(n) = S(2n+1) = x_{0.4+\frac{0.2}{2n+1}}$ , as the above methods, we can obtain  $\text{clu}_{\mathcal{T}}(W, \frac{1}{2}) = \tilde{1} - \mu$ .  $\square$

## References

1. C.L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182–190.
2. K.C. Chattopadhyay and S.K. Samanta, *Fuzzy topology*, Fuzzy sets and Systems **54** (1993), 207–212.
3. S.L. Chen and J.S.Cheng, *On convergence of nets of L-fuzzy sets*, J. Fuzzy Math. **2** (1994), 517-524.
4. S.L. Chen and J.S.Cheng,  *$\theta$ -Convergence of nets of L-fuzzy sets and its applications*, Fuzzy sets and Systems **86** (1997), 235-240.
5. M. Demirci, *Neighborhood structures in smooth topological spaces*, Fuzzy sets and Systems **92** (1997), 123-128.
6. R.N. Hazra, S.K. Samanta and K.C. Chattopadhyay, *Gradation of openness: Fuzzy topology*, Fuzzy sets and Systems **49(2)** (1992), 237–242.
7. Y.C. Kim and Y.S. Kim, *Fuzzy r-derived sets in fuzzy topological spaces*, J. Fuzzy Logic and Intelligent Systems **9(6)** (1999), 644–649.
8. K. Kuratowski, *Topology*, Academic Press Inc., New York, 1966.
9. Liu Ying-Ming and Luo Mao-Kang, *Fuzzy topology*, World Scientific Publishing Co., Singapore, 1997.

10. Pu Pao-Ming and Liu Ying-Ming, *Fuzzy topology I; Neighborhood structure of a fuzzy point and Moore-Smith convergence*, J.Math.Anal. Appl. **76** (1980), 571-599.
11. A.P. Sostak, *On a fuzzy topological structure*, Rend. Circ. Matem. Palermo Ser.II **11** (1985), 89-103.
12. Wang Guo-Jun, *Pointwise topology on completely distributive lattices*, Fuzzy sets and Systems **30** (1989), 53-62.

Department of Mathematics  
Kangnung National University  
Kangnung, Kangwondo 210-702, Korea

Department of Applied Mathematics  
Pai Chai University  
Taejon 302-735, Korea