

## **The Estimation of Early Health Effects for Different Combinations of Release Parameters and Meteorological Data**

**Jongtae Jeong and Wondea Jung**

Korea Atomic Energy Research Institute  
150 Dukjin-dong, Yusong-gu, Taejon 305-353, Korea  
jtjeong@kaeri.re.kr

(Received November 29, 2000)

### **Abstract**

Variations in the number of early health effects resulting from the severe accidents of the YGN 3&4 nuclear power plants were examined for different combinations of release parameters and meteorological data. The release parameters and meteorological data were selected in combination to define a limited number of basic spectra characterized by release height, heat content, release time, warning time, wind speed, rainfall rate, and atmospheric stability class. Variant seasonal spectra were also defined in order to estimate the potential significance of seasonal variations as a factor determining the incidence or number of early health effects. The results show that there are large differences in consequences from spectrum to spectrum, although an equal amount and mix of radioactive material is released to the atmosphere in each case. Also, there are large differences in the estimated number of health effects from season to season due to distinct seasonal variations in meteorological combinations in Korea. Therefore, it is necessary to consider seasonal characteristics in developing optimum emergency response strategies.

**Key Words** : early health effects, release parameters, meteorological data, seasonal characteristics

### **1. Introduction**

After the TMI and Chernobyl accidents, the importance of consequence analysis resulting from severe accidents at a nuclear power plant has increased because the influence of severe accidents on the environment and human beings is large in spite of their low occurrence probability. If a severe accident of a nuclear power plant were to

proceed to containment failure, radioactive materials would be released to the atmosphere. When such an accidental release occurs, the radioactive materials in the plume while dispersing in the atmosphere would be transported by a prevailing wind. In result, the radioactive materials would contaminate the environment, and finally the population would be exposed to radiation. The consequences resulting from such an accidental

release are health effects and economic impacts. These consequences are estimated by a sequence of mathematical and statistical models such as atmospheric dispersion, deposition of the material onto the ground, and the effects of airborne and deposited materials on humans and the environment.

The potential importance of offsite health and economic consequences of an accidental release from a nuclear power plant is a function of many factors such as the source term, weather condition, emergency response plan, and so on. The estimation of the total influence of these factors on offsite health and economic consequences is very complex. Therefore, it is useful to quantify their separate influences on offsite health and economic consequences based on their relative importance. Jeong and Ha[1] investigated the relative influences of source term release parameters such as release height, heat content of the plume, release time, release duration, and warning time on offsite health consequences. Also, Jeong and Ha[2] examined the relative influences of meteorological data such as wind speed, rainfall rate, and atmospheric stability class on offsite health consequences. They examined the variation of health effect cases such as early fatalities, early injuries, cancer fatalities, cancer injuries, early fatality risk, and cancer fatality risk resulting from the change of source term release parameters and meteorological data.

Although the unique accident sequence and source term release rate are considered in the consequence analysis, the resulting offsite health effects may be different if the source term release parameters and weather conditions are different. Therefore, we have made basic spectra based on the relative importance of source term release parameters and meteorological data on offsite health effects. We then investigated the variation of health effects resulting from the severe

accidents of the YGN 3&4 nuclear power plants from spectrum to spectrum by using MACCS code[3]. Also, we have investigated the seasonal variation of health effects based on spectra considering seasonal characteristics because we have four distinct seasons in Korea. The information obtained through this research will be very useful in developing optimum emergency response strategies for reducing offsite risks from the viewpoint of accident management.

## **2. Atmospheric Dispersion and Meteorological Data**

If a severe nuclear power plant accident were to proceed to containment failure, a fraction of radionuclides in the form of noble gases, halogens, and aerosols would be released to the atmosphere. An assessment of the impact of such releases to the environment and the general public requires the calculation of airborne and ground concentrations of each radionuclide at various distances from the reactor. When released into the atmosphere, radioactive gases and aerosols will follow prevailing winds and be diffused due to atmospheric turbulence. Predictions of dispersion are most commonly made from the Gaussian plume model[4] due to its economy of computing time, simplified input requirements, and reasonable agreement with experimental data over flat terrain. It is also very useful for repetitive calculations and sensitivity studies.

As the plume of radioactive material travel downwind from the reactor, material is removed from the plume by radioactive decay and by deposition onto the ground. Also, the basic dispersion model is usually modified to take account of a number of additional effects such as radioactive decay, the turbulent wake of the reactor building, the broadening of the time-averaged plume as a function of release duration

**Table 1. The Frequencies of Atmospheric Stability Classes for Four Seasons**

Atmospheric Stability(%) Season	A	B	C	D	E	F
Spring	0.7	1.4	3.4	33.7	59.4	1.5
Summer	3.2	7.8	20.5	34.2	33.7	0.6
Fall	4.1	11.2	38.6	31.5	13.0	1.5
Winter	2.7	5.0	12.7	32.7	44.7	2.2

to account for plume meander, mixing layer depth, surface roughness, and plume rise due to the thermal buoyancy of the plume. Plume rise, dispersion, downwind transport, and deposition onto the ground depend on the prevailing weather conditions such as wind speed, rainfall rate, and atmospheric stability. In most consequence analysis codes, the meteorological data file of the site region is used. This file is usually composed of one year of hourly wind speed, rainfall rate, and atmospheric stability recorded at the site or nearby weather service station.

The atmospheric transport models implemented in MACCS require hourly readings of wind speed, rainfall rate, and atmospheric stability as input. In addition, four values of the mixing height, one for each season of the year, must also be specified. The constant weather conditions are selected among the five ways to specify the required 120 hours of weather data that constitute a weather sequence because it is possible to estimate the variation of health effects due to the change of values of meteorological data. First of all, seasonal average wind speeds are obtained by using the monthly average wind speed that is obtained by averaging the hourly wind speed. The resulting seasonal average wind speeds are 3.0 m/sec for spring, 3.4 m/sec for summer, 3.6 m/sec for fall, 4.0 m/sec for winter. These values are assumed to be representative for each season, and are used in the formation of seasonal spectra. The maximum rainfall rate of hourly meteorological data is selected as the representative rainfall rate for each

season because the early fatalities and early fatality risk increase as the rainfall rate increases[2]. These values are 9.4 mm/hr for spring, 50.0 mm/hr for summer, 9.0 mm/hr for fall, 11.4 mm/hr for winter. The frequencies of the six atmospheric stability classes defined by Pasquill[5] are listed in Table 1. The most frequent atmospheric stability class is slightly stable(E) for spring, neutral(D) for summer, slightly unstable(C) for fall, and slightly stable(E) for winter. These atmospheric stability classes are assumed to be representative for each season and are used in the formation of seasonal spectra. A constant wind direction, west-north-west, was postulated for all spectra for the conservative results.

### 3. Modeling of Health Effects

The source term profiles which were derived from the Individual Plant Examination (IPE) of the YGN 3&4 nuclear power plants[6] were used to evaluate health consequences. According to the IPE results, 19 source term categories (STC) are defined by categorizing similar containment failure modes. The calculated source term release fractions are listed in Table 2. The core inventory data for fission products used for health effect calculations were derived from the ORIGEN2[7] calculations using the end-of-cycle inventory of fission products for the conservative evaluation because fission product buildup is greatest at the end-of-cycle conditions.

The MACCS code is used to evaluate the health

**Table 2. Fractional Source Term Release for Release Categories**

Nuclide Group	STC 1 & 2	STC 3	STC 4	STC 6 & 10	STC 7 & 11	STC 8 & 12
Noble Gases	0.0	1.0	1.0	1.0	1.0	1.0
Iodine	0.0	6.77E-02	2.22E-01	8.01E-03	8.41E-04	2.58E-02
Cesium	0.0	8.82E-02	2.23E-01	6.33E-03	1.14E-03	3.36E-02
Tellurium	0.0	1.07E-02	3.49E-02	1.71E-03	6.12E-04	3.71E-02
Barium	0.0	1.00E-03	3.29E-03	4.31E-05	1.08E-06	1.57E-03
Strontium	0.0	7.71E-04	2.52E-03	3.22E-05	8.05E-07	3.87E-03
Ruthenium	0.0	1.38E-03	4.51E-03	2.30E-05	5.75E-07	2.30E-05
Lanthanum	0.0	4.87E-04	1.59E-03	5.04E-07	1.30E-08	5.04E-07
Cerium	0.0	4.88E-04	1.60E-03	7.56E-07	1.90E-08	7.56E-07

  

Nuclide Group	STC 14	STC 15	STC 16	STC 17	STC 18	STC 19
Noble Gases	1.0	1.0	1.0	1.0	1.0	7.41E-01
Iodine	6.95E-01	0.97E-01	5.02E-03	6.02E-02	3.59E-01	1.13E-01
Cesium	5.85E-01	1.29E-01	3.29E-03	3.95E-02	2.35E-01	9.24E-02
Tellurium	1.96E-01	3.59E-02	9.12E-04	1.09E-02	6.53E-02	9.27E-02
Barium	6.45E-03	1.18E-03	3.01E-05	3.61E-04	2.15E-03	1.46E-03
Strontium	4.02E-03	7.36E-04	1.87E-05	2.24E-04	1.34E-03	1.15E-03
Ruthenium	2.04E-03	3.74E-04	9.52E-06	1.14E-04	6.79E-04	8.21E-04
Lanthanum	1.00E-04	1.83E-05	4.66E-07	5.59E-06	3.33E-05	1.80E-05
Cerium	1.50E-04	2.75E-05	6.99E-07	8.39E-06	4.50E-05	2.55E-05

effects resulting from the source terms of the YGN 3&4 nuclear power plants. In MACCS, the dispersion and deposition of radionuclides released from the reactor containment to the atmosphere were modeled with a Gaussian plume model. The site was selected as the center of a polar grid and the grid was divided into 16 equally spaced sectors with the outermost radius extending to 80 km. Each sector was divided further into 10 elements to reasonably account for the site specific population distribution. Evacuation and temporary relocation are considered as emergency response actions. These actions are to mitigate the effects of a release of radioactivity during a severe accident and are designed to reduce radiation exposures, public health effects, and economic impacts from an accident. Other parameters such as protection factors for inhalation or skin exposure, resuspension, cloud and other shielding

factors, and the specific input required for deriving chronic effects, are assumed to be the default values recommended in the MACCS User's Guide[8].

## 4. Results and Discussion

### 4.1. Formation of Spectra

The basic spectra were made by using the results of the relative influences of source term release parameters and meteorological data on health effects. The source term release parameters used in the formation of basic spectra are release height, heat content of the plume, release time, and warning time. Only the early health consequences are considered because the information obtained through the research will be used in developing the optimum emergency

**Table 3. The Basic Spectra and the Parameter Values**

Parameters Spectrum	Release Height (m)	Heat Content (MW)	Release Time (hr)	Warning Time (hr)	Wind Speed (m/sec)	Rainfall Rate (mm/hr)	Atmospheric Stability Class
Spectrum-1	0	1	2.0	3.0	2	50	F
Spectrum-2	20	10	1.0	2.0	4	30	E
Spectrum-3	40	20	0.5	1.0	6	10	D
Spectrum-4	60	30	4.0	0.5	10	0	B

**Table 4. The Seasonal Spectra and the Parameter Values**

Parameters Spectrum	Release Height (m)	Heat Content (MW)	Release Time (hr)	Warning Time (hr)	Wind Speed (m/sec)	Rainfall Rate (mm/hr)	Atmospheric Stability Class
Spring-1	0	1	2.0	3.0	3.0	9.4	E
Spring-2	20	10	1.0	2.0	3.0	9.4	E
Spring-3	40	20	0.5	1.0	3.0	9.4	E
Spring-4	60	30	4.0	0.5	3.0	9.4	E
Spring-5	0	1	2.0	3.0	3.0	0.0	E
Summer-1	0	1	2.0	3.0	3.4	50.0	D
Summer-2	20	10	1.0	2.0	3.4	50.0	D
Summer-3	40	20	0.5	1.0	3.4	50.0	D
Summer-4	60	30	4.0	0.5	3.4	50.0	D
Summer-5	0	1	2.0	3.0	3.4	0.0	D
Fall-1	0	1	2.0	3.0	3.6	9.0	C
Fall-2	20	10	1.0	2.0	3.6	9.0	C
Fall-3	40	20	0.5	1.0	3.6	9.0	C
Fall-4	60	30	4.0	0.5	3.6	9.0	C
Fall-5	0	1	2.0	3.0	3.6	0.0	C
Winter-1	0	1	2.0	3.0	4.0	11.4	E
Winter-2	20	10	1.0	2.0	4.0	11.4	E
Winter-3	40	20	0.5	1.0	4.0	11.4	E
Winter-4	60	30	4.0	0.5	4.0	11.4	E
Winter-5	0	1	2.0	3.0	4.0	0.0	E

response strategies during the emergency phase which begins immediately after the accident and could last up to seven days following the accident. The basic spectra listed in table 3 were made by considering characteristics of the relative influences of release parameters and meteorological data on early health effects[1, 2].

The parameter values of Spectrum-1 in Table 3 are composed of values that show maximum values of early health consequences. Therefore, one can use the parameter values of Spectrum-1 in order to obtain the most conservative results if the exact values of the release parameters and meteorological data are not available. As the

**Table 5. The Variation of Health Effects Based on Basic Spectra**

Health Effects Spectrum	Early Fatalities	Early Injuries	Early Fatality Risk (0.0-1.6 km)	Early Fatality Risk (1.6-3.2 km)	Pop. Weighted Early Fatality Risk
Spectrum-1	43.4	720	1.00E+00	1.58E-02	1.08E-03
Spectrum-2	38.4	806	1.00E+00	7.18E-03	9.59E-04
Spectrum-3	0.496	899	2.66E-02	0.0	1.24E-05
Spectrum-4	0.0	0	0.0	0.0	0.0

number of the Spectrum increases, the impact of the release parameters and meteorological data on early health consequences decreases. Finally, the parameter values of Spectrum-4 are those within the range of available data that show minimum values of early health consequences.

The seasonal spectra are listed in Table 4. The values of the four source term release parameters of Spectrum-1 through -4 for each season are the same values as the basic spectra. However, the meteorological data such as wind speed, rainfall rates, and atmospheric stability are the values assumed to be representative for each season as mentioned in the atmospheric dispersion and meteorological data. The Spectrum-5 of each season is the case that has the same parameter values of Spectrum-1 of each season except for the rainfall rates. In these spectra, the rainfall rate is zero. According to the analysis of the meteorological data at the YGN 3&4 nuclear power plants site, the number of hours that rainfall occurred among the 8,760 hours is only 440 hours, which is about 5% of the total data. Therefore, we added Spectrum-5 for each season in order to compare the results for the cases with and without rainfall rates.

#### 4.2 Results and Discussion

The overall health effects were calculated based on the assumptions and parameter values mentioned above. According to the results, the

values of early fatalities and total latent cancer fatalities are small relative to the total number of individuals, and the total latent cancer fatalities are larger than the early fatalities. This is due to the time span for the calculation, i.e., the calculated latent cancer fatalities occur over several decades. The individual early fatality risk and individual latent cancer fatality risk are  $7.52 \times 10^{-8}$  per year and  $2.45 \times 10^{-7}$  per year, respectively. These values are below the safety goal of the USNRC. However, these values are one or two order of magnitudes larger than the results of the five plants calculated in the NUREG-1150 studies. This can be due to the weather patterns at the site of the YGN 3&4 nuclear power plants. According to the analysis of the meteorological data at the site, the most frequent wind direction is west-north-west. The western part of the site is a marine area and the eastern part of the site is a populated region. Therefore, many individuals may be in the direct pathway of the radioactive plume.

The variations of early health consequences based on the basic spectra are listed in Table 5. As shown in Table 5, the early health consequences except early injuries considered in this study show maximum values in the case of Spectrum-1, and then decrease as the number of the Spectrum increases. The early injuries are much larger than the early fatalities. These facts are attributed to the time span for calculation and the atmospheric dispersion. Early fatalities are incurred during short

**Table 6. The Variation of Health Effects Based on Seasonal Spectra**

Health Effects Spectrum	Early Fatalities	Early Injuries	Early Fatality Risk (0.0-1.6 km)	Early FatalityRisk (1.6-3.2 km)	Pop. Weighted Early Fatality Risk
Spring-1	35.0	1464	1.00E+00	1.23E-02	8.75E-04
Spring-2	36.1	1502	1.00E+00	1.86E-02	9.00E-04
Spring-3	10.7	136	6.10E-01	0.0	2.67E-04
Spring-4	0.0	0	0.0	0.0	0.0
Spring-5	13.6	2122	7.60E-01	0.0	3.39E-04
Summer-1	48.1	617	7.93E-01	0.0	1.20E-03
Summer-2	51.2	642	1.00E+00	0.0	1.28E-03
Summer-3	45.2	572	1.00E+00	0.0	1.13E-03
Summer-4	0.0	0	0.0	0.0	0.0
Summer-5	2.51	1992	1.41E-01	0.0	6.26E-05
Fall-1	2.09	365	7.82E-02	0.0	5.21E-05
Fall-2	3.41	414	1.23E-01	0.0	8.51E-05
Fall-3	2.18	413	7.78E-02	0.0	5.45E-05
Fall-4	0.0	0	0.0	0.0	0.0
Fall-5	0.0	596	0.0	0.0	0.0
Winter-1	28.2	1298	9.98E-01	7.14E-03	7.05E-04
Winter-2	30.3	1331	9.99E-01	1.09E-02	7.57E-04
Winter-3	18.9	1226	8.94E-01	0.0	4.71E-04
Winter-4	0.0	0	0.0	0.0	0.0
Winter-5	4.41	1832	2.95E-01	0.0	1.10E-04

time periods, usually within weeks. However, early injuries due to pulmonary effects are incurred up to one year. The atmospheric dispersion increases due to the atmospheric stability and high wind speed as the number of the Spectrum increases. Also, as the release height and heat content of the plume increase, the area affected by the radioactive plume is increased due to atmospheric dispersion. Therefore, the number of people affected by the radioactive plume increases in spite of low concentrations.

In the case of Spectrum-4, all values of the health consequences are zero although equal amount of radioactive material is released to the atmosphere. From this fact, we can figure out that the source term release parameters and meteorological data have a great impact on the offsite health consequences. Therefore, relatively

exact values of release parameters and meteorological data should be provided in order to predict accident consequences accurately. One can use the parameter values of Spectrum-1 to obtain conservative results if one cannot obtain the exact values of release parameters and meteorological data.

The centerline early fatality risks between two specified intervals show similar trends for different spatial intervals. The centerline early fatality risks are on the order of  $10^{-5}$  for 3.2~4.8km,  $10^{-7}$  for 4.8~6.4km, and zero for distances beyond 6.4km. Therefore, the centerline early fatality risks between 1.6~3.2km are listed in Table 5. The results show that the early fatality risk decreases rapidly as the distance from the site increases. Therefore, actions such as evacuation should be taken to protect the workers and the general

public. Also, the warning time, which is the time after accident initiation at which the offsite alarm is rung, must be short so that evacuation can begin as soon as possible to protect the surrounding population. The population weighted risk which accounts for the population distribution shows similar trends to other health consequences.

The variation of early health consequences based on seasonal spectra considering the four distinct seasonal characteristics of Korea is shown in Table 6. The results for the spectra of each season show similar trends as those of the basic spectra. However, the results of Spectrum-2 for each season show the most conservative results. Therefore, we can use the source term release parameter values of Spectrum-2 for each season and typical meteorological data for each season when the exact values of source term release parameter are not available. In the Fall, all the values of the early health consequences are minimum. This is attributed to the weather condition of Fall. The most frequent atmospheric stability of Fall is slightly unstable(C), while the most frequent atmospheric stability classes of other seasons are neutral(D) or slightly stable(E). Therefore, the concentration of radioactive material is relatively low in the Fall, although the area affected by the radioactive plume is increased due to atmospheric dispersion. In the Summer, the early fatalities and the resulting early fatality risk show a maximum value due to the large rainfall rate. In the case of high rainfall rates, all materials washed out from the radioactive plume will be deposited onto the ground. This would produce very high ground concentrations in the spatial elements onto which deposition would occur instead of spreading over many spatial elements, and it is likely that large exposures are delivered over short time periods. Therefore, there are more early fatalities and less early injuries in Summer than other seasons.

The results of Spectrum-5 for each season show less early fatalities and more early injuries than those of Spectrum-1. The early fatality risk of Spectrum-5 for each season is also less than that of Spectrum-1 for each season. This is attributed to the plume depletion due to wet deposition because the Spectrum-5 of each season has the same parameter values without rainfall as Spectrum-1. That is, the radioactive plume is spread over much more spatial elements because of the lack of rain. Therefore, the radionuclide concentration is lower and the area affected by the radioactive plume is larger than the case of Spectrum-1.

## **5. Conclusions**

The variations of health effects resulting from the severe accidents of the YGN 3&4 nuclear power plants were examined for various combinations of release parameters and meteorological data. A small number of cases were defined based on the relative importance of the source term release parameters and meteorological data on the number and frequency of offsite early health effects. We then investigated the variation in various measures of early health effects resulting from severe accidents at the YGN 3&4 nuclear power plants from case to case by using MACCS code. Also, we investigated the variation in these measures of early health effects for meteorological conditions considered representative of the four distinct seasons in Korea.

The results show that with the same amount of radioactive material is released to the atmosphere, there are large differences in the number and frequency of early health effects from case to case. Therefore, release parameters and meteorological data have to be well characterized in order to estimate accurately the early health effects arising



from an accident. The variation of average individual early fatalities versus distance shows that the early fatality risk decreases rapidly as the distance from the site increases. Therefore, evacuation can be a very effective emergency response action in order to protect the population. There are large differences in the number of early health effects from season to season. In the fall, the early fatalities and early fatality risk show minimum values due to enhanced dispersion arising from increased atmospheric instability. The early fatalities show a maximum value in Summer, due to a large rainfall rate. It is necessary to consider seasonal characteristics in developing emergency response strategies to take account of seasonal variations in the induction of early health effects. The information obtained through this research will be very useful in developing optimum emergency response strategies for reducing offsite risks from the viewpoint of accident management.

### Acknowledgement

This project has been carried out under the Nuclear R&D Program by MOST.

### References

1. Jongtae Jeong and Jaejoo Ha., The influence of source term release parameters on health effects. *Journal of the Korean Nuclear Society* 31(3), 294-302, (1999).
2. Jongtae Jeong and Jaejoo Ha., The influence of meteorological data on health effects. KAERI/TR-1230/99, (1999).
3. H-N Jow, J. L. Sprung, J. A. Rollstin, L. T. Ritchie, and D. I. Chanin, MELCOR Accident Consequence Code System(MACCS), Model Description, Sandia National Laboratories, (1990).
4. Slade D. H. (Ed.). *Meteorology and nuclear energy*, TID-24190, U.S. Atomic Energy Commission, (1968).
5. Pasquill F., The estimation of the dispersion of windborne material, *Meteorological Magazine* 90, (1961).
6. C. K. Park et al., A study on individual plant examination of nuclear power plants (containment performance analysis) (1994).
7. A. G. Croff, A User's Manual for the ORIGEN2 Computer Code: ORNL/TM-7175, Oak Ridge National Laboratory, (1980).
8. D. I. Chanin, J. L. Sprung, L. T. Ritchie, and H-N Jow, MELCOR Accident Consequence Code System(MACCS), User's Guide, Sandia National Laboratories, (1990).
9. USNRC, Severe accident risks: an assessment of five nuclear power plants, NUREG-1150, (1990).