

1.

가 (8).

50% 가 (6,7)

가 (1)

가 , , , .

, BRAC1 BRAC2 .

가 (1).

10-15%

1 1

3-5 2

(9). 2 가

6 .

(3,4). 가

n-6 () () .

) 가 가

(4). , EPA DHEA 가

(fish oil) n-3 가

가

(5). 15-19

(6,7)

(10,11).

가 .

21 가 가 , 50 가 가 .

(21). (, ,) 가

(tyrosine kinase)

in vitro

(22) in vivo (epidermal growth factor) 가 3.

가 10-15%

(23). 가 가 가

21 mRNA 1 2 가

(24). 가 , 50 mRNA 가 가 , , II 가

50 , (EGF receptor) 가

가 가

가

21 가

(21). kg

25, 250mg

- ship to risk for breast cancer: an 8-year follow-up. *JAMA* 1992;268:2037-44.
4. Rose DP. Dietary fatty acids and prevention of hormone-responsive cancer. *Proc Soc Exp Biol Med* 1997;216:224-33.
5. Trichopoulos A, Ligiou P. Worldwide patterns of dietary lipids intake and health implications. *Am J Clin Nutr* 1997;66(suppl):961S-4S.
6. Boice JD, Monson RR. Breast cancer in woman after repeated fluoroscopic examinations of the chest cancer *Inst* 1997;59:863-7.
7. McGregor DH, Land CE, Choi K, et al. Breast cancer incidence among atomic bomb survivors, and Nagasaki. *J Natl Cancer Res* 1977;59:799-811.
8. Adami H-O, Lipworth L, Titus-Ernstoff L, et al. Organochlorine compounds and estrogen-related cancers in women. *Cancer Causes Control* 1995;6:551-66.
9. Russo J, Russo IH. DNA labeling index and structure of the rat mammary gland as determinants of its susceptibility to carcinogenesis. *J Natl Cancer Inst* 1978;61:1451-9.
10. Grubbs CJ, Farnell DR, Hill DL, McDonough KC. Chemoprevention of N-nitroso-N-methylurea-induced mammary cancers by pretreatment with 17 β -estradiol and progesterone. *J Natl Cancer Inst* 1985;74:927-31.
11. Rosso J, Wilgus G, Russo IH. Susceptibility of the mammary gland to carcinogenesis. I. Differentiation of the mammary gland an determinant of tumor incidence and type of lesion. *Am*
4. : *Am J Clin Nutr* 2000; 71S (1705-1707)
- 가
- 가
- 가
1. food, nutrition, and the prevention of cancer: a global perspective. Washington, DC: World Cancer Research Fund/American Institute for Cancer Research, 1997.
2. Colditz GA, Frazier AL. Models of cancer show that risk is set by events of early life: prevention efforts must shift focus. *Cancer Epidemiol Biomarkers Prev* 1995;4:567-71.
3. Willett WC, Hunter DJ, Stampfor MJ, et al. Dietary fat and fiber in relation-

- J Pathol 1979;96:721-36.
12. Lee HP, Gourley L, Duffy SW, Essteve J, Lee J, Day NE, Dictury effects on breast cancer risk in Singapore. *Lancet* 1991;337:1197-200.
 13. Yuan J-M, Wang Q-S, Ross RK, et al. Diet and risk of breast cancer in Shanghai and Tianjin, China. *Br J Cancer* 1995;71:1353-8.
 14. Wu AH, Ziegler RG, Horn-Ross PL, et al. Tofu and risk of breast cancer in Asian-Americans. *Cancer Epidemiol Biomarkers Prev* 1996;5:901-6.
 15. Ziegler RG, Hoover RN, Hildeshein RN, et al. Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst* 1993;85:1819-27.
 16. Lamartiniere CA, Holland MB. Neonatal diethylstilbestrol prevents spontaneously developing mammary tumors. In: Li JJ, Nandi S, Li SA, eds. *Hormonal carcinogenesis*. New York: Springer Verlag, 1992:305-8.
 17. Lamartiniere CA, Murrill WB, Manzollilo PA, et al. Genistein alters the ontogeny of mammary gland development and protects against mammary cancer in rats. *Proc Soc Exp Biol Med* 1997;217:358-64.
 18. Lamartiniere CA, Moore J, Holland MB, Barnes S. Neonatal genistein chemoprevents mammary carcinogenesis. *Proc Soc Exp Biol Med* 1995;208:120-3.
 19. Lamartiniere CA, Moore JB, Brown NM, et al. Genistein suppresses mammary cancer in rats. *Carcinogenesis* 1995;16:2833-40.
 20. Murrill WB, Brown NM, Manzollilo PA, et al. Prepubertal genistein exposure suppresses mammary cancer and enhances gland differentiation in rats. *Carcinogenesis* 1996;17:1451-7.
 21. Fritz WA, Coquard L, Wang J, Lamartiniere CA. Dietary genistein: perinatal mammary cancer prevention, bioavailability and toxicity testing in the rat. *Carcinogenesis* 1998;19:2151-8.
 22. Akiyama T, Ishida J, Nakagawa S, et al. Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J Biol Chem* 1987;262:5592-6.
 23. Dalu A, Haskell JF, Coward L, Lamartiniere CA. Genistein, a component of soy, inhibits the expression of the EGF and ErbB/Neureceptors in the rat dorsolateral prostate. *Prostate* 1998;37:36-43.
 24. Brown NM, Wang J, Cotroneo MS, et al. Prepubertal genistein treatment modulates TGF β , EGF, and EGF-receptor mRNAs and proteins in the rat mammary gland. *Mol Cell Endocrinol* 1998;144:149-45.
-
-
-