

Polymorphism of Insulin-like Growth Factor-I Gene in 13 Pig Breeds and its Relationship with Pig Growth and Carcass Traits

Wang Wenjun¹, Huang Lusheng*, Chen Kefei, Gao Jun, Ren Jun, Ai Huashui and Lin Wanhua

Jiangxi Provincial Key Laboratory for Animal Biotechnology, Jiangxi Agricultural University
Nanchang 330045, P. R. China

ABSTRACT : The polymorphism of insulin-like growth factor-I (IGF-I) in 13 pig breeds (total n=559) was detected by PCR-*Hha*I-RFLP, and allele A (151 bp and 28 bp) or allele B (116 bp, 35 bp and 28 bp) were observed. In these pig breeds, it was found that European pig breeds carried high frequencies of allele B, while Chinese native pig breeds carried high frequencies of allele A. Meanwhile the role of porcine IGF-I was investigated in 117 Nanchang White pigs and 360 Large Yorkshire pigs. Eight traits about growth and carcass were recorded for analyzing the associations between IGF-I gene polymorphism and performance quantitative traits. In the Nanchang White pigs, those with AA genotype generally had higher birth weight than those with AB genotype ($p<0.05$), but all these genotypes had no significant effect on the other traits which had been analyzed. In Large Yorkshire pigs, those with BB genotype had higher 2 months and 6 months body weight than those with AA genotype ($p<0.05$), and had a thicker hind-back-fat thickness and mid-back-thickness than those with AB and BB genotypes ($p<0.05$). And those with BB genotype were the thinnest in Large Yorkshire. Furthermore, pigs with AA genotype had a lower lean percentage than those with AB and BB genotypes ($p<0.01$), and the lean percentage of those with BB genotype was the highest. Based on these results, it is possible to make the IGF-I gene locus into the application of marker-assisted selection programmes. (*Asian-Aust. J. Anim. Sci.* 2002, Vol 15, No. 10 : 1391-1394)

Key Words : Insulin-like Growth Factor-I Gene, Performance Traits, Pig, PCR-RFLP

INTRODUCTION

Growth rate and body composition are two important characteristics in livestock production. Swine production demands high growth rate and high lean percentage, together with efficient conversation of feed to meat. Pig performance traits have been successfully improved by using traditional pig selection systems; but the contribution of individual genes involving in these processes is still known little, which hinder the livestock genetics improvement.

Important economic traits are usually controlled by many genes and modified by environment factors, such as temperature, nutrition. Insulin-like growth factor-I is the candidate gene playing a role in the growth and body composition, and had been mapped on pig chromosome 5 (Winter ϕ et al., 1994). IGF-I is the mediator of GH (Growth Hormone), and one of its functions are to control muscle growth. The effects of IGF-I gene variants on performance traits had been described for chicken (Nagaraja, et al., 2000; Kita, et al., 2000) and mouse (Collins et al., 1993; Horvat et al., 1995). In pigs, associations between variants at the IGF-I locus and performance traits require further investigation.

In this study we investigated the polymorphism of IGF-I in 13 pig breeds and the effects of one IGF-I restriction fragment length polymorphism (RFLP) described by Fang et al. (1999) on performance traits in Nanchang White pigs and Large Yorkshire pigs.

MATERIALS AND METHODS

Animals

The ear notches of 13 pig breeds were collected, including 3 European breeds, 10 Chinese native breeds. They were Landrace (n=65, Dongxiang County Pig Farm and JAU Pig Breeding), Duroc (n=52, Dongxiang County Hongxing Pig Farm and JAU Pig Breeding), Pietrain (n=11, Jiangxi Provincial Pig Farm), Jinhua two-end Black (n=60, Jinghua county Pig Farm), Jiaxin Spotted (n=58, Jiaying county Pig Farm), Shengxian Spotted (n=57), Leping Spotted (n=48, Leping county Pig Farm), Yushan Black (n=66, Yushan county Pig Farm), Erhualian (n=34, JAU experimental Pig Farm), Zixi Black (n=9, Zixi county Pig Farm), Xinzhi Black (n=12, Xinzhi county Pig Farm), Wild Pig (n=30, Jiaying Pig Farm), Shangao Two-end Black (n=57, Shangao county Pig Farm). All of the collected pigs were unrelated within 3 generations. In order to detect the effects of different IGF-I genotypes on some production performances, we also developed two pure pig breeds. Nanchang White pigs were half sibs, one male was mated fifteen females to produce 117 offsprings. For Large Yorkshire pigs, one Large Yorkshire male was mated with 10 Large Yorkshire females, and two F1 males and 36 F1

* Corresponding Author : Huang Lusheng. Tel: +86-791-3818116, Fax: +86-791-3818116, E-mail: hlsh@public.nc.jx.cn

¹ Department of Food Science, Jiangxi Agricultural University, Nanchang 330045, P. R. China.

Received March 8, 2002; Accepted May 21, 2002

females were subsequently used to produce 360 offsprings.

Large Yorkshire pigs were bred at Jiangxi Provincial Pig Farm and Nanchang White pigs were bred at Jiangxi Nanchang White Pig Breeding Farm. At ten days pigs were marked permanently by ear notching and this tissue was retained for DNA test. At ten weeks of age, the piglets were taken into single pens. The fattening test started at or from the 120 days old. The two pig breeds were fed with a cereal diet *ad libitum* containing 19%, 16%, and 13% crude protein in the nursery, grower and finisher stages, respectively, and the weights of animals were recorded weekly. A total of eight traits were recorded at an average 210 days of age. The carcass composition traits were determined by PIGLOG105 (Denmark, 1996).

IGF-I genotypes defined by PCR-RFLP

Ear notches from offspring were frozen and DNA was extracted using a phenol/chloroform extraction method followed by ethanol precipitation (Strauss, 1991). Working dilutions of extracted DNA were prepared for each individual at a concentration of 50 ng/ μ l. Primers 5' AGCC CACAGGGTACGGCTC 3' and 5' CTTCTGAGCCTTG GGCATGTC 3' were used for PCR (polymerase chain reaction) amplification of insulin-like growth factor-I gene (Fang et al., 1999). The PCR mixture containing 50 ng genomic DNA, 25 pmol of each primer, 25 μ M of each dNTP, 1 unit of Taq DNA Polymerase and 1 \times reaction buffer in a 25 μ l reaction volume. PCR were processed on PE9600 (PERKIN ELMER) according to the procedure: first 95°C for 30s then 35 cycles: 94°C for 45s, 59°C for 45s, 72°C for 60s, at last 72°C for 480s. The 179 bp PCR products were subsequently digested by *Hha* I and revealed allele A (fragments of 151 bp and 28 bp) or allele B (fragments of 116 bp, 35 bp and 28 bp). The restriction digests were separated using 3.0% agarose gel in 1 \times TAE at a constant current of 50 mA. The gels were stained with ethidium bromide and the fragments were visualized using a UV transilluminator.

Statistics

Associations between IGF-I genotypes and performance traits were analyzed using SAS system (1989). Significance between genotypes was assessed by Student's *t*-tests.

RESULTS

The genotype frequencies and allele frequencies of IGF-I in 13 pig breeds was listed in Table 1.

From Table 1, the AA genotype frequency in European pig breeds was low, especially in Pietrain, its frequency was 0%. While in Chinese native pig breeds, AA genotype

frequency was high, and Erhualian was the highest (73.5%). Meanwhile, the B frequency in European pig breeds was high, especially in Pietrain, its frequency was 100%. And in Chinese native pig breeds the A frequency was high (above 50%), especially the A frequency of Taihu pigs was 86.8%. Also in Table 1 we found that the genotype frequency and allele frequency of Wild Pig were similar to European pig breeds.

Table 2 listed the observed genotypes and their frequencies. The genotype AA in two pig breeds was rare. AA frequency was 5.98% in Nanchang White and 5.83% in Large Yorkshire.

In Nanchang White pigs, birth weight was significantly associated with IGF-I genotypes. The pigs with AA genotype had higher birth weight than the ones with AB genotype ($p < 0.05$). But no significant difference was observed among three genotypes for 2 months body weight, 4 months body weight and 6 months body weight. As far as corrected back-fat thickness and average back-fat thickness were concerned, pigs with BB genotype were the thinnest, although no significant difference was observed.

In Large Yorkshire pigs, no significant difference was observed among three genotypes for birth weight, but the birth weight of the pigs with AA genotype also was the highest, this was the same as Nanchang White pigs. For 2 months body weight, pigs with BB genotype were higher than pigs with AB genotype ($p < 0.05$). For 4 months body weight, no significant difference was observed among three genotypes. For 6 months body weight, pigs with BB genotype were the highest, and had statistic significant ($p < 0.05$) to the pigs with AA genotype. For hind-back-fat thickness and mid-back-fat thickness, furthermore, pigs with BB and AB genotypes were significantly different ($p < 0.05$) to AA genotype. The pig with the BB genotype was the thinnest. For lean percentage, pigs with BB and AB genotypes had higher lean content than AA genotype ($p < 0.01$), and the pig with BB genotype was the highest.

DISCUSSION

In Table 1, the AA genotype frequencies and A allele frequencies of European pig breeds were rare, while those of Chinese native pig breeds were much higher. As a lean-type pig breed, Wild Pig carried no AA genotype and rare A allele frequency, the reason was the long natural selection and evolution. In Table 2, the two pig breeds, AA genotypes were also rare, this was similar to the results by Fang *et al.* (1999) and Wang *et al.* (2000), in which western pigs carried AA genotype with a low frequency, while Chinese native pig breeds had higher frequencies.

In two different pure pig breeds, birth weight of pigs with AA genotype was higher than the other two genotypes, and pigs with BB genotype had a higher 2 months BW

Table 1. The genotype and allele frequencies of IGF-I in 13 pig breeds

Breeds	Number	Genotype frequencies (%)			Allele frequencies (%)	
		AA	AB	BB	A	B
Landrace	65	24.6	20.0	55.4	34.6	65.4
Duroc	52	0	26.9	73.1	13.5	86.5
Pietrain	11	0	0	100	0	100
Jinhua two-end black	60	61.7	16.7	21.6	70.0	30.0
Jiaying black	58	22.8	56.1	21.1	50.9	49.1
Shengxian spotted	57	50.0	34.5	15.5	67.2	32.8
Leping spotted	48	33.3	62.5	4.2	64.6	35.4
Yushang black	66	59.1	28.8	12.1	73.5	26.5
Erhualian	34	73.5	26.5	0	86.8	13.2
Zixi black	9	55.6	44.4	0	77.8	22.2
Xinzhi black	12	33.3	41.7	25.0	54.2	45.8
Wild pig	30	0	6.7	93.3	3.3	96.7
Shanggao two-end black	57	50.0	37.9	12.1	69.0	31.0

Table 2. Effects of different genotypes on some production performances in two pure pig breeds

Breeds	Nanchang white			Large yorkshire		
	AA	AB	BB	AA	AB	BB
Number	7	15	95	21	65	274
Birth weight (kg)	1.32±0.17 ^a	1.07±0.25 ^b	1.20±0.24 ^{ab}	1.47±0.25	1.42±0.26	1.45±0.18
2 mon. BW (kg)	20.00±2.65	19.99±3.65	20.83±3.65	23.30±2.38 ^{ab}	21.84±3.30 ^b	24.28±2.27 ^a
4 mon. BW (kg)	55.17±12.22	56.29±12.01	57.50±5.68	58.63±5.48	60.20±4.09	61.40±7.70
6 mon. BW (kg)	83.00±8.34	82.42±6.67	84.64±7.94	91.52±9.83 ^b	97.38±9.32 ^{ab}	101.68±12.01 ^a
Hind-back-fat thickness (cm)				2.16±0.18 ^a	1.93±0.21 ^b	1.90±0.16 ^b
Mid-back-fat thickness (cm)				2.15±0.31 ^a	1.86±0.20 ^b	1.77±0.21 ^b
Lean percentage (%)				56.95±2.95 ^b	59.26±2.44 ^A	61.60±2.29 ^A
Corrected back-fat depth (cm)	2.45±0.24	2.40±0.15	2.36±0.18			
Average back-fat depth (cm)	2.51±0.18	2.45±0.22	2.45±0.22			

Note: Values are M±SEM. Means with low case or capital letter within the same row in the same breed are significantly different ($p < 0.05$ or $p < 0.01$).

(body weight), 4 months body weight, 6 months body weight than those pigs with AB and BB genotypes. As far as lean percentage, mid-back-fat thickness, hind-back-fat thickness, corrected back-fat thickness and average back-fat thickness were concerned, AA genotype had negative effect. From the results of two pure pig breeds we have studied, it is concluded that, if we select pigs with AB and BB genotypes in the same population, we could get more lean percentage and less back-fat thickness. Also the more BB genotype pigs are selected, the more carcass is gotten. This is similar to the results by Casas-Carrillo et al. (1997), Horvat et al. (1995) and Seo et al. (2001).

In Nanchang White pigs, there is no significant difference among three genotypes for 2 months body weight, 4 months body weight, 6 months body weight, corrected back-fat thickness and average back-fat thickness, but the body weight at these days were highest and the fat depths are the thinnest. The affective trends were similar to Large Yorkshire and the course maybe: Nanchang white is a hybrid (Large Yorkshire×Binhu Black), and only breeding for 6 generations, the pigs contents 20% blood of Binhu Black, which can effect the function of allele B. From the primary results, it is suggested to select pigs with B allele

can improve the body weight and carcass lean percentage.

In two pig breeds, AA genotype had different significant effect on the birth weight. The plausible reasons are as follows: different management and nutritious level; different sample numbers. From the results of table 2, it is found that relationships between birth weight and 2 months body weight are contrary to the principle that the higher birth weight, the higher 2 months body weight, besides the above two reasons, there is a third reason that pigs with AA genotype have different biological characters such as weaning stress, suitability and genetics speciality.

Now many studies of IGF-I are only focused on its biochemical functions and nutritional modulation. But few studies were conducted to review whether polymorphism of IGF-I gene could be used as a genetic marker in pig and breeding system, thus the results could provide a theoretical and an experimental reference for pig genetic improvement.

ACKNOWLEDGEMENTS

The authors thank Dr. Fang (CAU, Beijing, China) for technical assistance and material gifts. This study was supported by State Major Basic Research Development

Program (Grant No. G20000161, China).

REFERENCES

- Casas-Carrillo, E., A. Prill-Adams and S. G. Price et al. 1997. Relationship of growth hormone and insulin-like growth factor-I genotypes with growth and carcass traits in swine. *Anim. Genet.* 28:88-93.
- Collins, A. C., I. C. A. Martin and B. W. Kirkpatrick. 1993. Growth and quantitative trait loci (QTL) on mouse chromosome 10 in a Quackenbush-Swiss \times 57BL/6J backcross. *Mammalian Genome.* 4:454-458.
- Fang Meiyang, Liu Honglin and Jiang Zhihua et al. 1999. Detection of insulin-like growth factor- I gene polymorphisms in 6 pig breeds (in Chinese), *Animal Husbandry & Veterinary Medicine.* 31(1):12-13.
- Horvat, S. and J. F. Medrano. 1995. Interval mapping of high growth (hg), a major locus that increase weight gain in mice. *Genetics.* 139:907-920.
- Kita, K., C. Noda and K. Miki et al. 2000. Relationship of IGF-I mRNA to tissue development in chicken embryos of different strains. *Asian-Aust. J. Anim. Sci.* 14(12):1659-1664.
- Milner, S. J., J. A. Carver, F. J. Ballard and G. L. Francis. 1999. Probing the disulfide folding pathway of insulin-like factor-I. *Biotechnology and Bioengineering,* 62(6):693-711.
- Nagaraja, S. C. and J. Aggrey SE Yao et al. 2000. Trait association of a genetic marker near IGF- I gene in egg-laying chickens. *The Journal of Heredity.* 91(2):150-156.
- PIGLOG, SFK Institute Inc. 1996. PIGLOG User's Guide: Version 3.1th, SFK-TECHNOLOGY A/S Institute Inc., Vandtaarnsvej 77, DE-2860, Soeborg, Denmark.
- Rinderknecht, E. and R. E. Humbel. 1978. The amino acid sequence of human insulin-like growth factor I and its structural homology with proinsulin. *J. Biol. Chem.* 253(8): 2679-2776.
- Salmon, W. D. Jr and W. H. Daughaday. 1958. The importance of amino acids diayzable components of rat serum which promote sulfate uptake by cartilage from hypophysectomized rats *in vitro.* *J Lab Clin Med.* 51:167-173.
- Samaras, S. E., S. F. Canning and J. A. Barber et al. 1996. Regulation of insulin-like growth factor-I biosynthesis in porcine granulosa cells. *Endocrinology.* 137(11):4657-4664.
- SAS Institute Inc. 1989. SAS/STAT User's Guide: Version 6. 4th edn. SAS Institute Inc., Cary, North Carolina.
- Seo, D. S., J. S. Yun and W. J. Kang, et al. 2001. Association of insulin-like growth factor-I (IGF-I) gene polymorphism with serum IGF- I concentration and body weight in Korean native Ogot chicken. *Asian-Aust. J. Anim. Sci.* 14(7):915-921.
- Strauss, W. M. Preparation of genomic DNA from mammalian tissue. In: *Current Protocols in molecular Biology* (ed. by F.M. Ausubel, R. Brent, R.E. Kingston et al.) Vol.1, pp. 2.2.1-2.2.3. John Wiley and Sons, New York, N. Y., USA.
- Svoboda, M. E. 1980. Purification of somatomedin-C from human plasma: Chemical and Biological properties, partial sequence analysis, and relationship to other somatomedins. *Biochemistry.* 19:790.
- Wang Wenjun, Lan Rutao and Wu Dengkuan et al. 2000. Detection of polymorphism of insulin-like growth factor- I gene locus in swine (in Chinese). *Animal Biotechnology Bulletin,* Oct: 175-178.
- Winter ϕ , A. K., M. Fredholm and L. Andersson. 1994. Assignment of the gene for insulin-like growth factor-I (IGF-I) to chromosome 5 by linkage mapping. *Anim. Genet.* 25(1):37-39.