

THE DENJOY-STIELTJES EXTENSION OF THE BOCHNER, DUNFORD AND PETTIS INTEGRALS

CHUN-KEE PARK

ABSTRACT. In this paper we introduce the concepts of Denjoy-Stieltjes-Dunford, Denjoy-Stieltjes-Pettis and Denjoy-Stieltjes-Bochner integrals of Banach-valued functions and then prove some properties of them.

1. Introduction

The Denjoy integral of a real-valued function which is an extension of the Lebesgue integral was studied by some authors ([1],[2],[3],[7]). In [5] we introduced the Denjoy-Stieltjes integral which is the generalization of the Denjoy integral and obtained some properties of the Denjoy-Stieltjes integral. R.A.Gordon[2], J.L.Gamez and J.Mendoza[1] studied the Denjoy extension of the Bochner, Pettis and Dunford integrals which is defined by the Denjoy integral. In this paper we deal with the Denjoy-Stieltjes extension of the Bochner, Pettis and Dunford integrals which is the generalization of the Denjoy extension of the Bochner, Pettis and Dunford integrals. We first define Denjoy-Stieltjes-Dunford, Denjoy-Stieltjes-Pettis and Denjoy-Stieltjes-Bochner integrals of Banach-valued functions using the Denjoy-Stieltjes integral and then prove some properties of them.

2. Preliminaries

We give some definitions and results to be used in this paper. Throughout this paper, X denotes a real Banach space and X^* its

Received July 25, 2002.

2000 Mathematics Subject Classification: 28B05, 26A39, 46G10.

Key words and phrases: Denjoy-Stieltjes integral, Denjoy-Stieltjes-Dunford integral, Denjoy-Stieltjes-Pettis integral, Denjoy-Stieltjes-Bochner integral.

This work was supported by a grant from Research Institute for Basic Science at Kangwon National University.

dual.

DEFINITION 2.1[3]. Let $F : [a, b] \rightarrow X$ and let $E \subset [a, b]$.

(a) The function F is BV on E if $V(F, E) = \sup \left\{ \sum_{i=1}^n \|F(d_i) - F(c_i)\| \right\}$

is finite where the supremum is taken over all finite collections $\{[c_i, d_i] : 1 \leq i \leq n\}$ of nonoverlapping intervals that have endpoints in E .

(b) The function F is AC on E if for each $\epsilon > 0$ there exists $\delta > 0$ such that $\sum_{i=1}^n \|F(d_i) - F(c_i)\| < \epsilon$ whenever $\{[c_i, d_i] : 1 \leq i \leq n\}$ is a finite collection of nonoverlapping intervals that have endpoints in E and satisfy $\sum_{i=1}^n (d_i - c_i) < \delta$.

(c) The function F is BVG on E if E can be expressed as a countable union of sets on each of which F is BV.

(d) The function F is ACG on E if F is continuous on E and if E can be expressed as a countable union of sets on each of which F is AC.

DEFINITION 2.2[2]. Let $F : [a, b] \rightarrow X$ and let $t \in (a, b)$. A vector z in X is the approximate derivative of F at t if there exists a measurable set $E \subset [a, b]$ that has t as a point of density such that $\lim_{\substack{s \rightarrow t \\ s \in E}} \frac{F(s) - F(t)}{s - t} = z$. We will write $F'_{ap}(t) = z$.

A function $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy integrable on $[a, b]$ if there exists an ACG function $F : [a, b] \rightarrow \mathbb{R}$ such that $F'_{ap} = f$ almost everywhere on $[a, b]$. The function f is Denjoy integrable on the set $E \subset [a, b]$ if $f\chi_E$ is Denjoy integrable on $[a, b]$.

DEFINITION 2.3[2]. (a) A function $f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on $[a, b]$ if for each $x^* \in X^*$ the function x^*f is Denjoy integrable on $[a, b]$ and if for every interval I in $[a, b]$ there exists a vector x_I^{**} in X^{**} such that $x_I^{**}(x^*) = \int_I x^*f$ for all $x^* \in X^*$.

(b) A function $f : [a, b] \rightarrow X$ is Denjoy-Pettis integrable on $[a, b]$ if f is Denjoy-Dunford integrable on $[a, b]$ and if $x_I^{**} \in X$ for every interval I in $[a, b]$.

(c) A function $f : [a, b] \rightarrow X$ is Denjoy-Bochner integrable on $[a, b]$ if there exists an ACG function $F : [a, b] \rightarrow X$ such that F is approximately differentiable almost everywhere on $[a, b]$ and $F'_{ab} = f$ almost everywhere on $[a, b]$.

DEFINITION 2.4[5]. Let $F : [a, b] \rightarrow X$ and let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function and let $E \subset [a, b]$.

(a) The function F is BV with respect to α on E if $V(F, \alpha, E) = \sup \left\{ \sum_{i=1}^n \|F(d_i) - F(c_i)\| \frac{\alpha(d_i) - \alpha(c_i)}{d_i - c_i} \right\}$ is finite where the supremum is taken over all finite collections $\{[c_i, d_i] : 1 \leq i \leq n\}$ of nonoverlapping intervals that have endpoints in E .

(b) The function F is AC with respect to α on E if for each $\epsilon > 0$ there exists $\delta > 0$ such that $\sum_{i=1}^n \|F(d_i) - F(c_i)\| < \epsilon$ whenever $\{[c_i, d_i] : 1 \leq i \leq n\}$ is a finite collection of nonoverlapping intervals that have endpoints in E and satisfy $\sum_{i=1}^n [\alpha(d_i) - \alpha(c_i)] < \delta$.

(c) The function F is BVG with respect to α on E if E can be expressed as a countable union of sets on each of which F is BV with respect to α .

(d) The function F is ACG with respect to α on E if F is continuous on E and if E can be expressed as a countable union of sets on each of which F is AC with respect to α .

THEOREM 2.5[5]. Let $F : [a, b] \rightarrow X$ and let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subset [a, b]$. Then F is BV on E if and only if F is BV with respect to α on E .

THEOREM 2.6[5]. Let $F : [a, b] \rightarrow X$ and let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subset [a, b]$. Then F is AC on E if and only if F is AC with respect to α on E .

3. Denjoy-Stieltjes integral

In [5] we introduced the Denjoy-Stieltjes integral and obtained some results for the integral. In this section we give another result.

DEFINITION 3.1[5]. Let $F : [a, b] \rightarrow X$ and let $t \in (a, b)$ and let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$. A vector $z \in X$ is the approximate derivative of F with respect to α at t if there exists a measurable set $E \subset [a, b]$ that has t as a point of density such that $\lim_{\substack{s \rightarrow t \\ s \in E}} \frac{F(s) - F(t)}{\alpha(s) - \alpha(t)} = z$. We will write $F'_{\alpha, ap}(t) = z$.

We note that $F'_{ap}(t) = F'_{\alpha, ap}(t) \cdot \alpha'(t)$ for each $t \in (a, b)$.

DEFINITION 3.2[5]. Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$. A function $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy-Stieltjes integrable with respect to α on $[a, b]$ if there exists an ACG function $F : [a, b] \rightarrow \mathbb{R}$ with respect to α such that $F'_{\alpha, ap} = f$ almost everywhere on $[a, b]$. The function f is Denjoy-Stieltjes integrable with respect to α on a set $E \subset [a, b]$ if $f\chi_E$ is Denjoy-Stieltjes integrable with respect to α on $[a, b]$.

THEOREM 3.3[5]. Let $f : [a, b] \rightarrow \mathbb{R}$ and let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subset [a, b]$. Then f is Denjoy-Stieltjes integrable with respect to α on E if and only if $\alpha'f$ is Denjoy integrable on E .

THEOREM 3.4. Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$. If $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy-Stieltjes integrable with respect to α on each interval $[c, d] \subseteq (a, b)$ and $\int_c^d f d\alpha$ converges to a finite limit as $c \rightarrow a^+$ and $d \rightarrow b^-$, then f is Denjoy-Stieltjes integrable with respect to α on $[a, b]$ and $\int_a^b f d\alpha = \lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d f d\alpha$.

Proof. Since $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy-Stieltjes integrable with respect to α on each interval $[c, d] \subseteq (a, b)$, by Theorem 3.3 $\alpha'f : [a, b] \rightarrow \mathbb{R}$ is Denjoy integrable on each interval $[c, d] \subseteq (a, b)$ and $\int_c^d f d\alpha = \int_c^d \alpha'f$ for each interval $[c, d] \subseteq (a, b)$. Hence $\lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d \alpha'f = \lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d f d\alpha$ exists by hypothesis. By [3, Theorem 15.12], $\alpha'f$ is Denjoy integrable on

$[a, b]$ and $\int_a^b \alpha' f = \lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d \alpha' f$. By Theorem 3.3, f is Denjoy-Stieltjes integrable with respect to α on $[a, b]$ and $\int_a^b f d\alpha = \lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d f d\alpha$. \square

4. Denjoy-Stieltjes extension of the Bochner, Pettis and Dunford integrals

We introduce Denjoy-Stieltjes-Bochner, Denjoy-Stieltjes-Pettis and Denjoy-Stieltjes-Dunford integrals and investigate some properties of those integrals.

DEFINITION 4.1. Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$.

(a) $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on $[a, b]$ if for each $x^* \in X^*$ $x^* f$ is Denjoy-Stieltjes integrable with respect to α on $[a, b]$ and if for every interval I in $[a, b]$ there exists a vector $x_I^{**} \in X^{**}$ such that $x_I^{**}(x^*) = \int_I x^* f d\alpha$ for all $x^* \in X^*$.

(b) $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Pettis integrable with respect to α on $[a, b]$ if f is Denjoy-Stieltjes -Dunford integrable with respect to α on $[a, b]$ and if $x_I^{**} \in X$ for every interval I in $[a, b]$.

(c) $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Bochner integrable with respect to α on $[a, b]$ if there exists an ACG function $F : [a, b] \rightarrow X$ with respect to α such that F is approximately differentiable with respect to α almost everywhere on $[a, b]$ and $F'_{\alpha, ap} = f$ almost everywhere on $[a, b]$.

$f : [a, b] \rightarrow X$ is integrable in one of the above senses on the set $E \subseteq [a, b]$ if $f \chi_E$ is integrable in that sense on $[a, b]$.

THEOREM 4.2. Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subset [a, b]$. Then $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Bochner integrable with respect to α on E if and only if $\alpha' f : [a, b] \rightarrow X$ is Denjoy-Bochner integrable on E .

Proof. If $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Bochner integrable with respect to α on E , then there exists an ACG function $F : [a, b] \rightarrow X$ with respect to α such that F is approximately differentiable with

respect to α almost everywhere on $[a, b]$ and $F'_{\alpha, ap} = f\chi_E$ almost everywhere on $[a, b]$. By Theorem 2.6, F is ACG. F is also approximately differentiable almost everywhere on $[a, b]$ and $F'_{ap} = F'_{\alpha, ap}\alpha' = \alpha'f\chi_E$ almost everywhere on $[a, b]$. Hence $\alpha'f$ is Denjoy-Bochner integrable on E .

Conversely, if $\alpha'f : [a, b] \rightarrow X$ is Denjoy-Bochner integrable on E , then there exists an ACG function $F : [a, b] \rightarrow X$ such that F is approximately differentiable almost everywhere on $[a, b]$ and $F'_{ap} = \alpha'f\chi_E$ almost everywhere on $[a, b]$. By Theorem 2.6, F is ACG with respect to α on $[a, b]$. F is also approximately differentiable with respect to α almost everywhere on $[a, b]$ and $F'_{\alpha, ap} = \frac{1}{\alpha'}F'_{ap} = \frac{1}{\alpha'}\alpha'f\chi_E = f\chi_E$ almost everywhere on $[a, b]$. Hence f is Denjoy-Stieltjes-Bochner integrable with respect to α on E . \square

The following corollary is obtained from Theorem 4.2 and [2, Theorem 28].

COROLLARY 4.3. *Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$. If $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Bochner integrable with respect to α on $[a, b]$, then each perfect set in $[a, b]$ contains a portion on which $\alpha'f$ is Bochner integrable.*

THEOREM 4.4. *Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subseteq [a, b]$. Then $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on E if and only if $\alpha'f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on E .*

Proof. If $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on E , then for each $x^* \in X^*$ x^*f is Denjoy-Stieltjes integrable with respect to α on E and for every interval I in $[a, b]$ there exists a vector $x_I^{**} \in X^{**}$ such that $x_I^{**}(x^*) = \int_I x^*f\chi_E d\alpha$ for all $x^* \in X^*$. By Theorem 3.3, for each $x^* \in X^*$ $\alpha'(x^*f) = x^*(\alpha'f)$ is Denjoy integrable on E and $x_I^{**}(x^*) = \int_I x^*f\chi_E d\alpha = \int_I x^*(\alpha'f\chi_E)$ for all $x^* \in X^*$. Hence $\alpha'f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on E .

Conversely, if $\alpha'f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on E , then for each $x^* \in X^*$ $x^*(\alpha'f) = \alpha'(x^*f)$ is Denjoy integrable on E and for every interval I in $[a, b]$ there exists a vector $x_I^{**} \in X^{**}$ such

that $x_I^{**}(x^*) = \int_I x^*(\alpha' f \chi_E)$ for all $x^* \in X^*$. By Theorem 3.3, for each $x^* \in X^*$ $x^* f$ is Denjoy-Stieltjes integrable with respect to α on E and $x_I^{**}(x^*) = \int_I x^*(\alpha' f \chi_E) = \int_I \alpha'(x^* f \chi_E) = \int_I x^* f \chi_E d\alpha$ for all $x^* \in X^*$. Hence $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on E . \square

COROLLARY 4.5. *Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subseteq [a, b]$. Then $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Pettis integrable with respect to α on E if and only if $\alpha' f : [a, b] \rightarrow X$ is Denjoy-Pettis integrable on E .*

Proof. The proof is similar to Theorem 4.4. \square

The following Corollary is obtained from Theorem 3.3, Theorem 4.4 and [1, Theorem 3].

COROLLARY 4.6. *Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$ and let $E \subseteq [a, b]$. Then $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on E if and only if $x^* f$ is Denjoy-Stieltjes integrable with respect to α on E for all $x^* \in X^*$.*

THEOREM 4.7. *Let $\alpha : [a, b] \rightarrow \mathbb{R}$ be a strictly increasing function such that $\alpha \in C^1([a, b])$. If $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy-Stieltjes-Dunford integrable with respect to α on $[a, t]$ for all $t \in [a, b]$ and for each $x^* \in X^*$ $\lim_{t \rightarrow b} \int_a^t x^* f d\alpha$ exists, then f is Denjoy-Stieltjes-Dunford integrable with respect to α on $[a, b]$ and $\langle x^*, (DSD) \int_a^b f d\alpha \rangle = \lim_{t \rightarrow b} \langle x^*, (DSD) \int_a^t f d\alpha \rangle$ for each $x^* \in X^*$.*

Proof. If $f : [a, b] \rightarrow X$ is Denjoy-Stieltjes-Dunford integrable with respect to α on $[a, t]$ for all $t \in [a, b)$ and for each $x^* \in X^*$ $\lim_{t \rightarrow b} \int_a^t x^* f d\alpha$ exists, then by Theorem 3.4 $x^* f$ is Denjoy-Stieltjes integrable with respect to α on $[a, b]$ and $\int_a^b x^* f d\alpha = \lim_{t \rightarrow b} \int_a^t x^* f d\alpha$ for all $x^* \in X^*$. Take $c \in [a, b)$ and any sequence (t_n) in $[a, b)$ convergent to

b. Define $L_c(x^*) = \lim_{n \rightarrow \infty} \int_c^{t_n} x^* f d\alpha = \lim_{n \rightarrow \infty} \langle x^*, (DSD) \int_c^{t_n} f d\alpha \rangle$ for each $x^* \in X^*$. By the uniform bounded principle, the linear functional L_c is continuous on X^* , that is, $L_c \in X^{**}$. Hence it is immediate that f is Denjoy-Stieltjes-Dunford integrable with respect to α on $[a, b]$. Taking $c = a$, we get

$$\begin{aligned} \langle x^*, (DSD) \int_a^b f d\alpha \rangle &= \int_a^b x^* f d\alpha \\ &= \lim_{t \rightarrow b} \int_a^t x^* f d\alpha \\ &= \lim_{t \rightarrow b} \langle x^*, (DSD) \int_a^t f d\alpha \rangle \end{aligned}$$

for each $x^* \in X^*$. □

References

1. J.L. Gamez and J. Mendoza, *On Denjoy-Dunford and Denjoy-Pettis integrals*, Studia Math. **130** (1998), 115-133.
2. R.A. Gordon, *The Denjoy extension of the Bochner, Pettis and Dunford integrals*, Studia Math. **92** (1989), 73-91.
3. R.A. Gordon, *The integrals of Lebesgue, Denjoy, Perron and Henstock*, Grad. Stud. Math. 4, Amer. Math. Soc., Providence, R.I., 1994.
4. S. Hu and V. Lakshmikantham, *Some Remarks on Generalized Riemann Integral*, J. Math. Anal. Appl. **137** (1989), 515-527.
5. C.K. Park, *On Denjoy-Stieltjes integral*, Kangweon-Kyungki Math. Jour. **9(2)** (2001), 105-114.
6. S. Saks, *Theory of the integral*, Dover, New York, 1964.
7. D.W. Solomon, *Denjoy integration in abstract spaces*, Mem. Amer. Math. Soc. **85**, 1969.

Department of Mathematics
Kangwon National University
Chuncheon 200-701, Korea