

Synthesis of 2-Benzylphenols: Transformation of the Baylis-Hillman Adducts Derived from 2-Cyclohexen-1-one

Ka Young Lee, Jeong Eun Na, and Jae Nyoung Kim*

Department of Chemistry and Institute of Basic Science, Chonnam National University, Gwangju 500-757, Korea
Received December 5, 2002

Key Words : Baylis-Hillman adducts, 2-Cyclohexen-1-one, 2-Benzylphenols

Recently, we have reported on the novel synthesis of quinoline derivatives from the Baylis-Hillman adducts.^{1,2} As an extension of the reaction, we intended to prepare acridine skeleton from the Baylis-Hillman adducts of 2-cyclohexen-1-one as shown in Scheme 1. However, we could not prepare the desired compounds *via* the reaction scheme (*vide infra*). Instead, we could obtain 2-benzylphenol derivatives in good yields as shown in Scheme 2 and wish to report herein the results.

The Friedel-Crafts alkylation reaction is one of the most powerful methods to form the carbon-carbon bond in organic reactions. The Friedel-Crafts benzylation reaction is of great synthetic significance in industrial processes.³ However, synthesis of 2-benzylphenols regioselectively from phenols or benzyl phenyl ethers is difficult due to the formation of *ortho*-/*para*- mixtures.⁴ Synthesis of these compounds by Fries rearrangement of phenyl phenylacrylates also suffers from the formation of mixtures.⁵ *Ortho*-specific alkylation of phenols *via* 1,3,2-benzodioxaborins was known.⁶ 1,3,2-Benzodioxaborins can be reduced to *ortho*-alkyl phenols with *tert*-butylamine borane in the presence of aluminum chloride.


The Baylis-Hillman reaction of 2-fluorobenzaldehyde (**1a**) and 2-cyclohexen-1-one (**2**) was carried out in aqueous THF with the aid of DMAP at room temperature to give the corresponding adduct **3a** in 58% yield as reported previously.⁷ Acetylation of **3a** with Ac₂O/DMAP gave **4a** in 91% yield. Initially, we examined the reaction of **4a** and *p*-toluenesulfonamide in the presence of K₂CO₃ in DMF. However, 2-(2-fluorophenyl)methylphenol (**5a**) was isolated in 74% yield, unexpectedly. The formation of **5a** occurred

well without tosylamide. Actually, the yield of **5a** was improved up to 94% without tosylamide as shown in Table 1 (entry 1).


Thus, we prepared some Baylis-Hillman acetates of 2-cyclohexen-1-one and examined their conversion to 2-arylmethylphenols and the results are summarized in Table 1. As mentioned previously, the Baylis-Hillman reaction of **1** and **2** was carried out in aqueous THF in the presence of 0.1 equiv. of DMAP.⁷ The corresponding adducts **3a-h** were obtained in reasonable yields (41-64%) at room temperature. Following conversion to their acetate **4a-h** was excellent in all cases (CH₂Cl₂, Ac₂O/DMAP, rt, 90-98%). The reaction of **4a-f** in DMF in the presence of K₂CO₃ (1.0 equiv) gave **5a-f** in good yields (89-96%) in short time (1-2 h) at 60-70 °C.⁸ The formation of **5g**, the quinoline derivative, was carried out at room temperature. The reaction is believed to occur as depicted in Scheme 2: potassium carbonate assisted elimination of acetic acid and the following keto-enol tautomerization and 1,5-hydrogen transfer.⁹

We could not obtain the corresponding phenol derivative from the analogous reaction with **4h**, derived from hexanal. Intractable mixtures were observed on tlc at 60-70 °C. Instead, we could isolate cyclohexenone derivative **6** in 40% yield at room temperature.¹⁰ The structure of **6** was determined by ¹H-, ¹³C-, and ¹H-¹H COSY.⁸

In conclusion we disclosed unusual transformation of the Baylis-Hillman acetates of 2-cyclohexen-1-one into 2-arylmethylphenols. Further chemical transformation of the products to xanthene derivatives *via* the nucleophilic aromatic substitution strategy and the synthesis of acridines are underway.



Scheme 1



Scheme 2

Table 1. Synthesis of 2-benzylphenols **5**

Entry	Aldehyde 1	3a-h	4a-h	Conditions	Product
1		3a (58%) ⁷	4a (91%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 1 h	 5a (94%)
2		3b (55%) ⁷	4b (90%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 2 h	 5b (89%)
3		3c (63%) ⁷	4c (96%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 2 h	 5c (92%)
4		3d (41%)	4d (92%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 1 h	 5d (96%)
5		3e (53%) ⁷	4e (96%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 1 h	 5e (95%)
6		3f (56%) ⁷	4f (98%)	K ₂ CO ₃ (1.0 equiv) DMF, 60-70 °C 2 h	 5f (93%)
7		3g (53%) ⁷	4g (97%)	K ₂ CO ₃ (1.0 equiv) DMF, rt 6 h	 5g (61%)
8	CH ₃ (CH ₂) ₄ CHO	3h (64%) ⁷	4h (98%)	K ₂ CO ₃ (1.0 equiv) DMF, rt 20 h	 6 (40%)

Acknowledgments. This work was supported by the grant (R05-2000-000-00074-0) from the Basic Research Program of the Korea Science & Engineering Foundation.

References

- (a) Kim, J. N.; Chung, Y. M.; Im, Y. J. *Tetrahedron Lett.* **2002**, *43*, 6209. (b) Kim, J. N.; Lee, H. J.; Lee, K. Y.; Kim, H. S. *Tetrahedron Lett.* **2001**, *42*, 3737.
- Chung, Y. M.; Lee, H. J.; Hwang, S. S.; Kim, J. N. *Bull. Korean Chem. Soc.* **2001**, *22*, 799.
- (a) Keum, G.; Lim, H. J.; Kang, S. B.; Kim, Y.; Chung, B. Y. *Bull. Korean Chem. Soc.* **2000**, *21*, 809. (b) Yamato, T.; Fukumoto, M.; Sakae, N.; Furusawa, T.; Tashiro, M. *Synthesis* **1991**, 699. (c) Teubner, H.; Kramer, A.; Weuffen, W.; Schritter, E.; Grbel, G. EP 69,374 A2 (1983) (*Chem. Abstr.* **1983**, *98*, 215314q). (d) Citerio, A.; Fancelli, D. EP 309,009 A1 (1989) (*Chem. Abstr.* **1989**, *111*, 133777r). (e) Langer, R.; Buysch, H. J. EP 538,704 (1993) (*Chem. Abstr.* **1995**, *122*, 31111d). (f) Yamaguchi, K.; Tanabe, Y.; Hasegawa, K.; Yamaguchi, A. EP 0,523,697 B1 (1993) (*Chem. Abstr.* **1993**, *119*, 149558). (g) Olah, G. A.; Olah, J. A.; Ohyama, T. *J. Am. Chem. Soc.* **1984**, *106*, 5284. (h) Goswami, A. *J. Chem. Research (S)* **2000**, 554.
- Regioselective synthesis of 2-benzylphenols from benzyl phenyl ether was achieved by using K10-montmorillonite, see Venkatachalapathy, C.; Pitchumani, K.; Sivasultramanian, S. *Indian J. Chem.* **1998**, *37B*, 301. For the synthesis of 2-benzylphenol from the benzylation of the dianion of *o*-cresol, see Bates, R.; Siahaan, T. J. *J. Org. Chem.* **1986**, *51*, 1432.
- Gu, W.; Weiss, R. G. *J. Org. Chem.* **2001**, *66*, 1775.
- Lau, C. K.; Williams, H. W. R.; Tardif, S.; Dufresne, C.; Scheiget, J.; Belanger, P. C. *Can. J. Chem.* **1989**, *67*, 1384.
- Lee, K. Y.; Gong, J. H.; Kim, J. N. *Bull. Korean Chem. Soc.* **2002**, *23*, 659.
- Representative spectroscopic data of **5a** is as follows: oil; ¹H NMR (CDCl₃) δ 3.99 (s, 2H), 4.88 (br s, 1H), 6.75-7.15 (m, 8H); ¹³C NMR (CDCl₃) δ 28.82 (d, *J* = 2.9 Hz), 115.21 (d, *J* = 22.0 Hz), 115.58, 121.02, 124.11 (d, *J* = 3.3 Hz), 125.76, 126.96 (d, *J* = 15.8 Hz), 127.86, 127.99, 130.78, 130.84, 153.51, 160.99 (d, *J* = 244.7 Hz); Mass (70 eV) *m/z* (rel. intensity) 78 (14), 106 (30), 152 (15), 181 (41), 183 (36), 202 (M⁺, 100). Representative spectroscopic data of **6** is as follows: oil; ¹H NMR (CDCl₃) δ 0.89 (t, *J* = 7.5 Hz, 3H), 1.27-1.34 (m, 4H), 1.43-1.50 (m, 2H), 2.22 (q, *J* = 7.5 Hz, 2H), 2.49-2.61 (m, 4H), 5.99-6.03 (m, 1H), 6.54-6.58 (m, 2H); ¹³C NMR (CDCl₃) δ 13.93, 22.44, 24.40, 27.25, 28.38, 31.52, 38.20, 123.99, 128.34, 131.79, 136.64, 199.49.
- We examined the synthesis of **5a** from **3a** by direct dehydration. The reaction of **3a** in benzene in the presence of *p*-TsOH showed the formation of complex mixtures. For the synthesis of similar 2-arylmethylphenols by direct dehydration in acidic medium in low yields, see (a) Patra, A.; Batra, S.; Joshi, B. S.; Roy, R.; Kundu, B.; Bhaduri, A. P. *J. Org. Chem.* **2002**, *67*, 5783. (b) Iwamura, T.; Fujita, M.; Kawakita, T.; Kinoshita, S.; Watanabe, S.-i.; Kataoka, T. *Tetrahedron* **2001**, *57*, 8455.
- (a) Negishi, E.-i.; Owczarczyk, Z. R.; Swanson, D. R. *Tetrahedron Lett.* **1991**, *32*, 4453. (b) Urdaneta, N.; Ruiz, J.; Zapata, A. J. *J. Organometal. Chem.* **1994**, *464*, C33.