

Communications

Bismuth(III) Nitrate Pentahydrate $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$: An Inexpensive and Mild Reagent for the Efficient and Clean Oxidation of Thiols to Disulfides


M. M. Khodaei,^{*} I. Mohammadpoor-Baltork,^{†*} and K. Nikoofar

Department of Chemistry, Razi University, Kermanshah 67149, Iran
†Department of Chemistry, Isfahan University, Isfahan 81746-73441, Iran
Received January 8, 2003

Key Words : Bismuth(III) nitrate pentahydrate, Thiols, Disulfides

Oxidation of thiols to their disulfides is an important transformation from both biological and synthetic points of views.^{1,2} Disulfides are important reagents in organic synthesis and can be used for the preparation of sulfinyl and sulfonyl compounds.³ A variety of reagents have already been used for the conversion of thiols to disulfides.⁴ However, some of these reagents suffer from disadvantages such as long reaction times, availability, toxicity, difficult work-up, preparation and instability. Consequently, there is still a need for introducing readily available, safe, stable and cheap reagents for the oxidation of thiols to disulfides.

The applications of bismuth compounds to organic transformations have been extensively investigated.⁵ Recently we introduced Bi(III) salts as efficient reagents for different organic transformations including the conversion of epoxides to thiiranes,^{6a,b} deprotection of 1,1-diacetates,^{6c} alcoholysis, hydrolysis and acetolysis of epoxides,^{6d} deprotection of silyl and tetrahydropyranyl ethers,^{6e,f} conversion of epoxides to 1,3-dioxolanes,^{6g} acetylation, benzoylation and formylation of alcohols, phenols, trimethylsilyl and tetrahydropyranyl ethers,^{6h,j} and conversion of thiocarbonyls to their carbonyl compounds.^{6k} In continuation of our ongoing work on Bi(III) catalysis, we now report an efficient and environmentally benign procedure for the oxidative coupling of thiols to their corresponding disulfides using bismuth(III) nitrate pentahydrate (Scheme 1).



Scheme 1

Bismuth(III) nitrate pentahydrate is an inexpensive, crystalline solid and commercially available reagent and requires no special handling. Oxidation of thiols to their disulfides was performed efficiently in the presence of 30 mol% of this reagent. The results of this study are summarized in Table 1. Aromatic thiols such as benzenethiol (**1a**), 4-methylbenzenethiol (**1b**), 4-bromobenzenethiol (**1c**), 4-chlorobenzenethiol (**1d**) and 2-naphthalenethiol (**1e**) exhibited a similar reactivity, leading to the corresponding disulfides **2a-2e** in excellent yields. Heteroaromatic thiol such as 2-pyridinethiol (**1f**) underwent oxidative coupling with $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$, to afford 2,2'-dipyridine disulfide (**2f**) in 93% yield. Under the same reaction conditions, benzylthiol (**1g**) and aliphatic thiols **1h-1i** were oxidized to their disulfides **2g-2i** in excellent yields. The selectivity of the present method is evident by the oxidation of 2-mercaptoethanol (**1h**), where only mercaptan functionality is converted to the disulfide **2h**.

In conclusion, the paper describes a facile synthesis of disulfides using catalytic amounts of $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$. The method offers several advantages including high yields of the products, short reaction times, ease of isolation of the products, and stable, inexpensive and relatively non-toxic reagent which make the reaction process convenient and environmentally benign.

^{*}Co-Corresponding authors. I. Mohammadpoor-Baltork (E-mail: imbaltork@sci.ui.ac.ir; Fax: +98-311-6689732), M. M. Khodaei (E-mail: mmkhoda@razi.ac.ir)

Table 1. Catalytic oxidation of thiols to disulfides with $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$

Thiol	R	Disulfide ^a	Time (min)	Isolated Yield %	mp (°C)	Lit. mp (°C)
1a	C_6H_5	2a	20	93	59-60	58-59 ^{4f}
1b	4-MeC ₆ H ₄	2b	5	96	44-45	45-46 ^{4e}
1c	4-BrC ₆ H ₄	2c	5	97	91-93	91-92 ^{4f}
1d	4-ClC ₆ H ₄	2d	10	98	70-71	72-74 ^{4g}
1e	2-Naphthyl	2e	45	97	142-144	144-145 ^{4h}
1f	2-Pyridyl	2f	40	93	55-57	55-57 ^{4f}
1g	$\text{C}_6\text{H}_5\text{CH}_2$	2g	40	93	70-71	69-70 ^{4f}
1h	HOCH_2CH_2	2h	40	98	oil ⁴ⁱ	
1i	<i>n</i> -C ₄ H ₉	2i	25	93	oil ^{4j}	
1j	<i>n</i> -C ₈ H ₁₇	2j	35	93	oil ^{4l}	
1k	$(\text{CH}_3)_2\text{CHCH}_2$	2k	20	96	oil ^{4k}	
1l	<i>c</i> -C ₆ H ₁₁	2l	40	94	oil ^{4k}	

^aAll products were identified by comparison of their physical and spectral data with those of authentic samples.

Experimental Section

General procedure: To a solution of thiol (1 mmol) in acetonitrile (10 mL) was added $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ (0.3 mmol) and the mixture was refluxed for the indicated time according to Table 1. The reaction mixture was filtered and the solid material was washed with acetonitrile (10 mL). Evaporation of the solvent followed by recrystallization or chromatography on silica gel afforded the pure disulfide in 93-98% Yields (Table 1).

Acknowledgement. We are thankful to the Razi and Isfahan University Research Councils for partial support of this work.

References

- Jocelyn, D. C. *Biochemistry of the Thiol Group*, Academic Press: New York, 1992; p 1.
- Capozzi, G.; Modena, G. *The Chemistry of Thiol Group*, Part 2; Patai, S., Ed.; Wiley: New York, 1974; p 785.
- (a) Kuhle, E. *The Chemistry of the Sulfenic Acids*, Georg Thieme Verlag: Stuttgart, 1973. (b) Douglass, I. B.; Norton, R. V. *J. Org. Chem.* **1968**, *33*, 2104. (c) Douglass, I. B. *J. Org. Chem.* **1974**, *39*, 563.
- (a) Christensen, L. W.; Heacock, D. J. *Synthesis* **1978**, 50. (b) Drabowicz, J.; Mikolajczyk, M. *Synthesis* **1980**, 32. (c) Firouzabadi, H.; Mohammadpoor-Baltork, I. *Bull. Chem. Soc. Jpn.* **1992**, *65*, 1485. (d) Iranpoor, N.; Firouzabadi, H.; Zolfigol, M. A. *Synth. Commun.* **1998**, *28*, 367. (e) Iranpoor, N.; Zeynizadeh, B. *Synthesis* **1999**, 49. (f) McKillop, A.; Koyuncu, D. *Tetrahedron Lett.* **1990**, *31*, 5007. (g) Wu, X.; Rieke, R. D.; Zhu, L. *Synth. Commun.* **1996**, *26*, 191. (h) Meshram, H. M.; Kache, R. *Synth. Commun.* **1997**, *27*, 2403. (i) Hirano, M.; Yakabe, S.; Ando, K.-I.; Morimoto, T. *J. Chem. Res. (S)* **1998**, 816. (j) Hirano, M.; Yakabe, S.; Monobe, H.; Morimoto, T. *J. Chem. Res. (S)* **1998**, 472. (k) Sridhar, M.; Vadivel, S. K.; Bhalerao, U. T. *Synth. Commun.* **1998**, *28*, 1499. (l) Rao, T. V.; Sain, B.; Murthy, P. S.; Rao, T. S. R. P.; Jain, A. K.; Joshi, G. C. *J. Chem. Res. (S)* **1997**, 300. (m) Noureldine, M. C.; Hendry, J.; Lee, D. G. *Synthesis* **1998**, 1587. (n) Kesavan, V.; Bonnet-Delpon, D.; Begue, J.-P. *Synthesis* **2000**, 223. (o) Movassagh, B.; Lakouraj, M. M.; Ghodrati, K. *Synth. Commun.* **1999**, *29*, 3597. (p) Hajipour, A. R.; Mallakpour, S. E. *J. Chem. Res. (S)* **2000**, 32. (q) Raghavan, S.; Rajender, A.; Joseph, S. C.; Rasheed, M. A. *Synth. Commun.* **2001**, *31*, 1477. (r) Salehi, P.; Farrokhi, A.; Gholizadeh, M. *Synth. Commun.* **2001**, *31*, 2777. (s) Wang, J.-X.; Gao, L.; Huang, D. *Synth. Commun.* **2002**, *32*, 963. (t) Sun, Y.-H.; Ko, K.-Y. *Bull. Korean Chem. Soc.* **2000**, *21*, 669. (u) Chen, F.-E.; Lu, Y.-W.; He, Y.-P.; Luo, Y.-F.; Yan, M.-G. *Synth. Commun.* **2002**, *32*, 3487.
- (a) Suzuki, H.; Ikegami, T.; Matano, Y. *Synthesis* **1997**, 249. (b) Leonard, N. M.; Wieland, L. C.; Mohan, R. S. *Tetrahedron* **2002**, *58*, 8373.
- (a) Mohammadpoor-Baltork, I.; Aliyan, H. *Synth. Commun.* **1998**, *28*, 3943. (b) Mohammadpoor-Baltork, I.; Khosropour, A. R. *Molecules* **2001**, 996. (c) Mohammadpoor-Baltork, I.; Aliyan, H. *Synth. Commun.* **1999**, *29*, 2741. (d) Mohammadpoor-Baltork, I.; Tangestaninejad, S.; Aliyan, H.; Mirkhani, V. *Synth. Commun.* **2000**, *30*, 2365. (e) Firouzabadi, H.; Mohammadpoor-Baltork, I.; Kolagar, S. *Synth. Commun.* **2001**, *31*, 905. (f) Mohammadpoor-Baltork, I.; Kharameshi, B.; Kolagar, S. *Synth. Commun.* **2002**, *32*, 1633. (g) Mohammadpoor-Baltork, I.; Khosropour, A. R.; Aliyan, H. *Synth. Commun.* **2001**, *31*, 3411. (h) Mohammadpoor-Baltork, I.; Khosropour, A. R.; Aliyan, H. *J. Chem. Res. (S)* **2001**, 280. (i) Mohammadpoor-Baltork, I.; Aliyan, H.; Khosropour, A. R. *Tetrahedron* **2001**, *57*, 5851. (j) Mohammadpoor-Baltork, I.; Khosropour, A. R. *Monatsh. Chem.* **2002**, *133*, 189. (k) Mohammadpoor-Baltork, I.; Khodaei, M. M.; Nikoofar, K. *Tetrahedron Lett.* **2003**, *44*, 591.