

Peptic Hydrolysate of Porcine Crude Myosin Has Many Active Fractions Inhibiting Angiotensin I-converting Enzyme

Kazunori Katayama¹, Hidetaka Fuchu¹, Masaaki Sugiyama¹, Satoshi Kawahara, Kiyoshi Yamauchi
Yukio Kawamura² and Michio Muguruma*

Department of Biochemistry and Applied Biosciences, Faculty of Agriculture, Miyazaki University
Miyazaki 889-2192, Japan

ABSTRACT : In order to clarify one of the biological functions of pork, we investigated whether a peptic hydrolysate of denatured porcine crude myosin showed inhibitory activity against angiotensin I-converting enzyme (ACE), which contributed to hypertension. Our results indicated that this hydrolysate showed relatively strong activity, and we therefore attempted to separate the involved peptides, which were considered to be active substances. To isolate these active peptides, the hydrolysate was separated using a solid-phase separation, gel filtration high-performance liquid chromatography (HPLC), and two kinds of reverse phase HPLC. In each stage of separation, many fractions were detected, almost all of which showed ACE inhibitory activity. Thus, we suggested that the activity of the hydrolysate as a whole was a result of the activities of the many individual peptides. Six peaks were distinguished, with yields from 34 to 596 ppm of original crude myosin. In addition to the six peaks, many other active fractions were found throughout the separation steps, strongly suggesting that whole porcine crude myosin itself had ACE inhibitory activity. Moreover, pork as food was considered to function as an ACE inhibitory material *in vivo*, because pork proteins consist primarily of crude myosin, which included almost all the myofibrillar structural proteins. (*Asian-Aust. J. Anim. Sci.* 2003, Vol 16, No. 9 : 1384-1389)

Key Words : Angiotensin I-converting Enzyme, Peptide, Porcine Crude Myosin, Protease Digestion, Pepsin

INTRODUCTION

The tertiary (physiologically active) function of food, which contributes to the modulation of our physiological systems, has been widely investigated, and certain functional foods have already been developed in Japan (Arai, 1996). Some of these studies have proposed to use food peptides as functional substances generated from foods by digestion with proteases. Among these peptides, peptide inhibitors for angiotensin I-converting enzyme (ACE) have attracted considerable attention. ACE plays an important role in the regulation of blood pressure: it contributes to hypertension by converting angiotensin I to angiotensin II (Skeggs et al., 1956) and by inactivating bradykinin. Although there are other factors involved in high blood pressure, many studies have focused on ACE inhibition and hypotension, because the inhibition of ACE brings about a significant decrease in blood pressure (Ondetti et al., 1977). ACE inhibitory peptides from animal-originated foods have been previously reported by Kawamura et al. (1992), Yokoyama et al. (1992), Seki et al. (1995), Ohta et al. (1999) and Fujita et al. (2000), among others. Since pork contains significant amounts of protein, it is fair to speculate that it contains many physiologically functional

peptides. ACE inhibitory peptides from porcine myosin have been reported by Arihara et al. (2001), and the research has been conducted on their hypotensive activity *in vivo* upon oral administration to spontaneously hypertensive rats (Nakashima et al., 2002). Nevertheless, almost all of the researches described above have primarily examined peptides derived from hydrolysates with non-digestive proteases, such as thermolysin and alkalase; a detailed investigation of peptides derived from pork proteins through gastrointestinal protease digestion is important in any discussion of the functions of these peptides in a biologically active food.

In a previous study, we have reported that the enzymatic hydrolysates of a variety of porcine proteins, such as myosin, actin, tropomyosin, troponin, and water-soluble proteins, showed ACE inhibitory activities (Katayama et al., 2003). In the present study, the purpose of this research was to clarify the ACE inhibitory activities of peptides from pork. For this purpose, the peptic hydrolysate of porcine crude myosin, which included almost all myofibrillar structural proteins and was denatured by heating, was separated to isolate ACE inhibitory peptides. We also include a discussion of the activity of the fractions derived through the various stages of separation.

MATERIALS AND METHODS

Preparation of porcine crude myosin

Pork loin (*Longissimus dorsi*) was obtained from Marudai Shimane Farm Co., Ltd. (Shimane, Japan). Crude

* Corresponding Author: M. Muguruma, Tel: +81-985-58-7203, Fax: +81-985-58-7203, E-mail: muguruma@cc.miyazaki-u.ac.jp

¹ Marudai Food Co., Ltd., Takatsuki, Osaka 569-8577, Japan.

² Graduate School of Applied Life Science, Kinki University, Nara 631-8505, Japan.

Received August 16, 2002; Accepted May 28, 2003




Figure 1. SDS-PAGE profiles of porcine crude myosin, which was denatured by heating at 95°C for 10 min, and was digested by pepsin. 0, 0.5, 2, 6, and 24: incubation time (h) for digestion. S: supernatant of peptic hydrolysate incubated for 6 h. MK: molecular weight marker for (b). For (a), myosin heavy chain (200 k Da) and actin (42 k Da) were indicated as molecular weight markers.

myosin was extracted from minced meat using Guba-Straub solution (0.3 M KCl, 0.1 M KH_2PO_4 , 0.05 M K_2HPO_4 , 2 mM adenosine triphosphate) following the method described by Margossian and Lowey (1982). The crude myosin extract included myosin, actin, tropomyosin, troponin and other myofibrillar structural proteins. It was dialyzed against phosphate-buffered saline (PBS; 0.8% NaCl, 10 mM Na-phosphate, pH 7.5) before hydrolysis.

Hydrolysis of crude myosin

Pepsin was obtained from Sigma Chemical Co. (St. Louis, MO, USA). Crude myosin (5 mg/ml) was suspended in PBS and denatured by heating for 10 min at 95°C. Its pH was adjusted to 2 with 1 M HCl, and pepsin was added at a 1:100 ratio of enzyme to substrate. After 6 h digestion at 37°C, the buffer pH was adjusted to 7.5 with 1 M NaOH, and enzymatic activity was terminated by heating for 10 min at 95°C. The reaction mixture was centrifuged for 10 min at 14,000 rpm, and the resulting supernatant was used for experiments.

ACE inhibitory assay

ACE inhibitory activity was measured following the method described by Cushman and Cheung (1971), with slight modifications (Katayama et al., 2003). Briefly,

samples were incubated with hippuryl-L-histidyl-L-leucine (HHL, Nacalai Tesque Inc., Kyoto, Japan) as the substrate and rabbit lung ACE (Wako Pure Chemical Industries Ltd., Osaka, Japan) in a buffer containing 0.13 M borate (pH 8.5) and 0.4 M NaCl. The enzyme reaction was terminated by adding 1 M HCl. Hippuric acid liberated from HHL was extracted by ethyl acetate, dried, and dissolved with 1 M NaCl. The absorbance of hippuric acid at 228 nm was measured and the ACE inhibitory activity was calculated. Inhibitory activity was identified as a 50% inhibitory protein concentration (IC_{50}) of the sample in assay. One unit of the ACE inhibitory activity was defined as the amount of inhibitor that inhibited 50% of ACE activity in 1 ml of reaction mixture (Seki et al., 1995). The specific activity (activity per weight of protein) of each fraction was calculated (Nakamura et al., 1995).

Isolation of ACE inhibitory peptides from peptic hydrolysate of porcine crude myosin

The peptic hydrolysate of porcine crude myosin was separated using a SEP-PAK Plus C_{18} cartridge (Waters Co., Milford, MA, USA) with 10, 20 and 30% CH_3CN . The active fractions were then separated by reverse phase (RP) high-performance liquid chromatography (HPLC) using a Cosmosil 5C₁₈ AR-II (4.5×150 mm) column (Nacalai Tesque), and eluted with a gradient of 10 to 80% CH_3CN in 0.1% trifluoroacetic acid (TFA) at a flow rate of 0.5 ml/min. The active fractions were then applied to the same column and eluted with an isocratic condition of CH_3CN in 0.1% TFA at a flow rate of 0.5 ml/min. The active fractions obtained from RP-HPLC were subjected to gel filtration HPLC using a TSK-gel G2000SW_{XL} (7.8×300 mm) (Tosoh Co., Tokyo, Japan), and eluted with 45% CH_3CN in 0.1% TFA at a flow rate of 0.5 ml/min. Finally, the active fraction was separated through RP-HPLC using a Cosmosil 5PE-MS (4.6×250 mm) column (Nacalai Tesque), and eluted with an isocratic condition of CH_3CN in 0.1% TFA at a flow rate of 1 ml/min. All elutions were monitored using absorbance at 215 nm.

Electrophoresis

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed following the method described by Laemmli (1970), and using a gradient (7.5 to 17.5% acrylamide) slab gel for separating proteins with a stacking gel (4.5%). Protein bands were stained with Coomassie Brilliant Blue R-250.

Determination of protein concentrations

The protein concentrations of the crude myosin and its hydrolysate were measured using the biuret method (Gornall et al., 1949) with bovine serum albumin as a standard. The peptide concentrations of fractions from the

Table 1. Angiotensin I-converting enzyme inhibitory activity of digestive protease hydrolysates from various foods

Origin	Protease	IC ₅₀ (μg/ml)	Reference
Porcine crude myosin	Pepsin	112	This paper
	Chymotrypsin	121	Katayama et al. (2003)
	Trypsin	166	
Sardine	Pepsin	620	Matsui and Kawasaki (2000)
	Chymotrypsin	1,280	
	Trypsin	2,030	
Dried bonito	Pepsin	47	Yokoyama et al. (1992)
	Chymotrypsin	117	
	Trypsin	161	
Ovalbumin	Pepsin	45	Fujita et al. (2000)
	Chymotrypsin	>1,000	
	Trypsin	>1,000	




Figure 2. First RP-HPLC chromatogram and activity of fraction eluted through SEP-PAK cartridge C₁₈ with 20% CH₃CN. Eluted fractions were collected every 5 min, and measured their ACE inhibitory activities. W, X, Y and Z were highly active fractions and subjected to the next separation step.

various separation steps were measured by the UV method (Murphy and Kies, 1960) using wavelengths of 215 and 225 nm.

RESULTS AND DISCUSSION

ACE inhibitory activity of peptic hydrolysate of porcine crude myosin

The SDS-PAGE pattern of the crude myosin hydrolysate with pepsin showed that many proteins with high molecular weight (MW) decreased as hydrolysis proceeded (Figure 1a); in fact, the supernatant from the peptic hydrolysate after 6 h digestion showed no relatively high MW proteins (Figure 1b). The protein concentration of this supernatant was 3.7 mg/ml, and about 74% of the original crude myosin (5 mg/ml) was recovered as water-soluble proteins, peptides

or free amino acids.

The peptic hydrolysate of crude myosin showed ACE inhibitory activity with 112 μg/ml of IC₅₀, while the original crude myosin showed no such activity. We suggested that peptides generated from crude myosin by peptic digestion were responsible for this elevated level of inhibition. The activity of this peptic hydrolysate was found to be stronger than that of hydrolysate from sardine and a little weaker than those of hydrolysates from bonito or ovalbumin (Table 1). These data indicated that porcine crude myosin included some peptides with high activity. Crude myosin hydrolyzed with chymotrypsin or trypsin showed weaker activity than the peptic hydrolysate (Katayama et al., 2003), and the activity levels of other hydrolysates (Table 1) showed the same tendencies found with crude myosin. Cheung et al. (1980) stated that dipeptides with hydrophobic amino acid residues showed ACE inhibitory activity. Furthermore, pepsin cleaves protein at hydrophobic amino acid residues such as Phe, Tyr, Leu, and Met, then peptic hydrolysate with hydrophobic amino acid residues could be speculated to include many ACE inhibitory peptides. We therefore carried out the present study to separate and isolate the active peptides from this hydrolysate.

Separation and activity of ACE inhibitory fractions

The peptic hydrolysate of crude myosin (371 mg) was first separated with a SEP-PAK Plus C₁₈ cartridge. Fractions eluted by 10, 20 and 30% CH₃CN showed ACE inhibitory activities, and their IC₅₀s were 36, 57 and 46 μg/ml, respectively. The fraction eluted by 20% CH₃CN, which had the most protein, was applied to a C₁₈ HPLC column and separated with a CH₃CN gradient. Almost all peptides were eluted at CH₃CN concentrations between 20 and 30%, that is, at the concentrations at which relatively hydrophobic peptides were expected to elute (Figure 2). Fractions were collected every 5 min and their ACE inhibitory activities were measured. Four fractions (W, X, Y, and Z), which were eluted between 35 and 55 min, had a relatively high protein concentration and showed strong activity with 43, 49, 44 and 14 μg/ml of IC₅₀. These




Figure 3. Second RP-HPLC chromatograms and activities of fractions (W, X, Y and Z) from the first RP-HPLC. Every peak was collected and its ACE inhibitory activity was measured. \square : ACE inhibitory activity of each fraction. X-10, X-11, Y-9 and Y-12 were subjected to the next separation step.




Figure 4. Gel filtration HPLC chromatograms of fractions (X-10, X-11, Y-9 and Y-12) from second RP-HPLC. Six fractions (A to F) were subjected to the next separation step. Aprotinin and oxytocin were used as molecular weight standards.

fractions were applied to the same column and separated with an isocratic elution: the fractions W, X, Y, and Z were eluted with 15, 18, 20 and 25% CH_3CN , respectively (Figure 3). Every peak was collected, dried and dissolved in 100 μl of H_2O , and its ACE inhibitory activities were measured. Almost all peaks showed inhibitory activity, suggesting that the activity of whole crude myosin hydrolysate resulted from the activities of many individual peptides. The peaks with relatively strong activity (over 50% inhibition, see Figure 3) were separated during the next step using gel filtration HPLC. Because many fractions in Figure 3 showed very weak activity after further purification using gel filtration HPLC, only 4 chromatograms of gel filtration HPLC (X-10, X-11, Y-9 and Y-12) are shown in Figure 4. Six fractions (A to F) with strong activity were obtained: their activity levels were 22, 26, 131, 70, 37 and 135 $\mu\text{g}/\text{ml}$ of IC_{50} , respectively. The molecular weights of these fractions were estimated at less than 6.5 kDa by comparing their retention times to those of molecular weight markers (aprotinin and oxytocin). Finally, these six active fractions were applied to a PE-MS column.




Figure 5. RP-HPLC chromatograms using PE-MS column of fractions (A, D, E and F) from gel filtration HPLC. Six fractions (A5, D2, E5, E7, F2 and F7) showed ACE inhibitory activity.

Because B and C showed very weak activity after further purification using a PE-MS column, only 4 chromatograms (A, D, E and F) are shown in Figure 5. Fractions A, D, E and F were eluted with 15, 13, 13 and 14% CH_3CN in 0.1% TFA, respectively. The peaks in Figure 5 (A5, D2, E5, E7, F2 and F7) showed ACE inhibitory activity; their corresponding IC_{50} s were 8, 76, 5, 10, 21 and 65 $\mu\text{g}/\text{ml}$. These activities were stronger than those of peptides described by Arihara et al. (2001) and as strong as those of peptides described by Fujita et al. (2000), Matsui and Kawasaki (2000) and Yokoyama et al. (1992).

Whole activity of porcine crude myosin

The yield of each fraction (A5, D2, E5, E7, F2 and F7) was between 3.4 and 59.6 mg/100 g of original crude myosin, and totalled about 120 mg/100 g (Table 2). The unit recovery of these fractions to whole hydrolysate was approximately 5,500 ppm and the activity of these fractions was concentrated over 4.5 times. With the exception of the last six fractions, many active fractions were found throughout all separation steps, suggesting that whole porcine crude myosin has ACE inhibitory activity in itself. Moreover, pork as food functions as an ACE inhibitory material *in vivo*, because pork proteins consist primarily of crude myosin, which includes almost all the myofibrillar structural proteins (Hattori, 1996). Although Arihara et al. (2001) have reported previously on ACE inhibitory peptides identified from porcine myosin, the active fractions in the present study were considered to differ from their peptides, because they used another enzyme (thermolysin) to digest myosin, which was not denatured. Furthermore, the ACE inhibitory peptides from chicken muscle reported by Fujita et al. (2000) were also considered to differ from the fractions used here, because they used thermolysin, too.

The role of digestive enzymes in producing biologically active peptides from food is considered to be highly significant in the modern study of foods (Katayama et al., 2003; Li et al., 1999). In the further research, we plan to

Table 2. Angiotensin I-converting enzyme inhibitory activity and recovery of fractions from porcine crude myosin

Separation step	Fraction	Protein		IC ₅₀	Unit ¹	Unit recovery	Specific activity ²
		µg	Yield mg/100 g	µg/ml		ppm	U/mg
	Intact crude myosin	500,000	100,000	>2,000			
	Peptic hydrolysate	371,840	74,368	112	3,320	1,000,000	8.9
1	SEP-PAK C ₁₈						
	20% CH ₃ CN	126,605	25,321	57	2,221	669,017	17.5
2	RP-HPLC						
	W	17,242	3,448	43	400	120,493	23.2
	X	20,157	4,031	49	415	124,924	20.6
	Y	18,075	3,615	44	412	124,012	22.8
	Z	6,080	1,216	14	444	133,681	73.0
	Subtotal	61,553	12,311		1,670	503,111	
4	Gel-HPLC						
	A	270	54	22	12.6	3,781	46.7
	B	427	85	26	16.2	4,878	37.9
	C	987	197	131	7.5	2,262	7.6
	D	434	87	70	6.2	1,858	14.3
	E	467	93	37	12.7	3,838	27.2
	F	844	169	135	6.3	1,887	7.5
	Subtotal	3,429	686		61.4	18,504	
5	RP-HPLC						
	(PE-MS)						
	A5	17	3.4	8	2.1	625	123.5
	D2	185	37.0	76	2.4	736	13.0
	E5	17	3.4	5	3.4	1,024	200.0
	E7	34	6.8	10	3.5	1,045	102.9
	F2	48	9.7	21	2.4	709	50.0
	F7	298	59.6	65	4.6	1,376	15.4
	Subtotal	599	119.9		18.3	5,515	

¹ One unit=protein (µg)/IC₅₀ (µg/ml). ² Units of ACE inhibitory activity per milligram of peptide.

identify these active peptides and to examine the *in vivo* activity of meat after oral administration.

REFERENCES

- Arai, S. 1996. Studies on functional foods in Japan-States of the art. *Biosci. Biotechnol. Biochem.* 60:9-15.
- Arihara, K., Y. Nakashima, T. Mukai, T. Ishikawa and M. Itoh. 2001. Peptide inhibitors for angiotensin I-converting enzyme from enzymatic hydrolysates of porcine skeletal muscle proteins. *Meat Sci.* 57:319-324.
- Cheung, H. S., F. L. Wang, M. A. Ondetti, E. F. Sabo and D. W. Cushman. 1980. Binding of peptide substrate and inhibitors of angiotensin-converting enzyme. *J. Biol. Chem.* 255:401-407.
- Cushman, D. W. and H. S. Cheung. 1971. Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung. *Biochem. Pharmacol.* 20:1637-1648.
- Fujita, H., K. Yokoyama and M. Yoshikawa. 2000. Classification and antihypertensive activity of angiotensin I-converting enzyme inhibitory peptides derived from food proteins. *J. Food Sci.* 65:564-569.
- Gomall, A. G., C. J. Bardawill and M. M. David. 1949. Determination of serum protein by means of the biuret reaction. *J. Biol. Chem.* 177:751-766.
- Hattori, A. 1996. *Shokuniku no kouzou to seibun*. In: *Niku no Kagaku* (Ed. A. Okitani) Asakura shoten, Tokyo, Japan. pp. 48-49. (in Japanese).
- Katayama, K., H. Fuchu, A. Sakata, S. Kawahara, K. Yamauchi, Y. Kawamura and M. Muguruma. 2003. Angiotensin I-converting enzyme inhibitory activities of porcine skeletal muscle proteins following enzyme digestion. *Asian-Aust. J. Anim. Sci.* 16:417-424.
- Kawamura, Y., T. Takane, M. Satake and T. Sugimoto. 1992. Physiologically active peptide motif in proteins: Peptide inhibitor of ACE from the hydrolysates of antarctic krill muscle protein. *J. Agric. Res. Quart.* 26:210-213.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227:680-685.
- Li, D. F., X. H. Zhao, T. B. Yang, E. W. Johnson and P. A. Thacker. 1999. A comparison of the intestinal absorption of amino acids in piglets when provided in free form or as a dipeptide. *Asian-Aust. J. Anim. Sci.* 12:939-943.
- Margossian, S. S. and S. Lowey. 1982. *Methods in Enzymology*, 85:55-59. Academic Press, New York.
- Matsui, T. and K. Kawasaki. 2000. Antihypertensive effects of bioactive peptides derived from food proteins. *J. Jpn. Soc. Nutr. Food Sci.* 53:77-85. (in Japanese)
- Murphy, J. B. and M. W. Kies. 1960. Note on spectrophotometric determination of proteins in dilute solutions. *Biochim. Biophys. Acta*, 45:382-384.
- Nakamura, Y., N. Yamamoto, K. Sakai, A. Okubo, S. Yamazaki and T. Takano. 1995. Purification and characterization of angiotensin I-converting enzyme inhibitors from sour milk. *J. Dairy Sci.* 78:777-783.
- Nakashima, Y., K. Arihara, A. Sasaki, H. Mio, S. Ishikawa and M. Itoh. 2002. Antihypertensive activities of peptides derived from porcine skeletal muscle myosin in spontaneously hypertensive rats. *J. Food Sci.* 67:434-437.
- Ohta, T., A. Iwashita, S. Sasaki and Y. Kawamura. 1999. Antihypertensive action of the orally administered protease hydrolysates of chum salmon head and their angiotensin I-

- converting enzyme inhibitory peptides. *Food Sci. Technol. Int.*, Tokyo, 3:339-343.
- Ondetti, M. A., B. Rubin and D. W. Cushman. 1977. Design of specific inhibitors of angiotensin-converting enzyme: New class of orally active anti hypertensive agents. *Sci.* 196:441-444.
- Seki, E., K. Osajima, H. Matsufuji, T. Matsui and Y. Osajima. 1995. Val-Tyr, an angiotensin I converting enzyme inhibitor from sardines that have resistance to gastrointestinal proteases. *Nippon Nogeikagaku Kaishi*, 69:1013-1020. (in Japanese)
- Skeggs, L. T., J. R. Kahn and N. P. Shumway. 1956. The preparation and function of the angiotensin-converting enzyme. *J. Exp. Med.* 103:295-299.
- Yokoyama, K., H. Chiba and M. Yoshikawa. 1992. Peptide inhibitors for angiotensin I-converting enzyme from thermolysin digest of dried bonito. *Biosci. Biotechnol. Biochem.* 56:1541-1545.