

:

1, 2
1 . 2 . 2 . 2 . 2
1 . 1 . 1 . 1

Multiple Attempts at Embryo Transfer do not Adversely Affect In-vitro Fertilization Pregnancy Rates: Related Mucus Contamination

**Byeong Jun Jung¹, Jong Sik Kim², Cheo Jin Kwon², Mi Jin Ryu², Myung Sin Kim²,
Eun Hee Kang¹, Jong Ok Sim¹, Hyun Jin Song¹, Ik Hwan Oh¹**

*¹Department of Obstetrics and Gynecology, ²Infertility Center,
Seoul Women's Hospital, Incheon*

Objective: We investigate the effects of multiple attempts of embryo transfer because of retained embryos in the catheter and of contaminated mucus on the transferred catheter.

Materials and Methods: We respectively analysed data between November 1998 and August 2002 from 305 patients of 369 cycles who underwent IVF-ET. Of these patients, 47 patients of 50 cycles (Group 2) were required multiple trial of embryo transfer. They were compared with an age-matched control groups (Group 1) with female factor infertility.

Pearson's χ^2 and Fisher's tests were used to compare proportions between discrete variables. Non-categorical data were compared using t-test. Statistical significance was set at $p < 0.05$.

Results: Embryos were significantly more likely to be retained when catheter was contaminated with mucus (Group 1: 22.4%; Group 2: 44.0%). The clinical pregnancy rates, however, for the contaminated mucus or not, were 46.8%, 43.5% respectively. There was no significant difference clinical pregnancy rate between those who had all their embryos transferred at the first attempt (45.4%) and those who required more than one attempt (48.0%).

Conclusions: Contaminated mucus in the catheter is associated with failed embryo transferred at the first attempt. Embryo transfers, however, that are repeated attempts do not adversely affect pregnancy rates following IVF-ET.

Key Words: Retained embryos, Mucus contamination, IVF-ET

가

가

2.

1)

gonadotropin releasing hormone agonist (GnRHa, Lucrine subQ, AB-BOT) follicle stimulating hormone (FSH, Metrodin, Serono) human menopausal gonadotropin (hMG, Pergonal, Serono) GnRHa (GnRHa long protocol)

21 GnRHa

3

FSH

FSH

hMG

estradiol(E₂)

18 mm

가 3

10,000 IU

human

chorionic gonadotropin (hCG, Profasi, Serono)

hCG

36

Tur-Kaspa

(Single Lumen Ovum Pick-Up Needle Set, COOK)

37 5% CO₂

(SZH10, Olympus)

(oocytes cumulus complex,

OCC)

OCC가

(germinal vesicle,

GV)

10% synthetic serum substrate (SSS

Irvine) 가 P 1 (Preimplantation 1)

(metaphase II, MII)

(metaphase I, MI)

(immature, GV) 20

3~4

8~10

~48

CO₂

3~5

가
가
Nabi (1997)⁵
가
(1998)⁶
가
가
가
1.
1998 11 2002 6
258 319
(Group 1)
가
47 50 (Group 2)

가

2)

F-10 Nutrient Mixture Medium

(Ham's F-10, Gibco) 0.5% antibiotics (Penicillin-G, Streptomycin sulfate) 가 . 5)

P 1 37

(2.04 mM CaCl₂ · 2H₂O, 101.6 mM NaCl, 4.69 mM KCl, 0.2 mM MgSO₄, 25 mM NaHCO₃, 0.33 mM Na-pyruvate, 21.4 mM Na-lactate, 0.05 mM Taurine, 0.15 mg/l Na-citrate, 10 ? g/l Gentamicin, 0.005 g/l Phenol red) 5% CO₂ 10% SSS가

가 P 1 72

가 . 가

가 Grade 1

Sigma (G1), 가

95% 가 G2, 가

, 37 5% CO₂ (CO₂ incubator, Heraeus) 10 20% G3,

50%

G4, 가

3) (human follicular fluid, hFF) 가 가

가 G5 . 8

G1, G2 GQER

(good quality embryo rates)

30 35 (heat inactivation) 0.22? m (Sterivex GV, Millipore) -20 . bivalve speculum

4) (masturbation) G1~G3 4~5

37 20~30 (WHO)⁸ Tomcat

10% (Tomcat Catheter, 8890-793021, Sherwood medical, USA) . Tomcat

가 Ham's F-10 2 1 ml TDT

pellet Ham's F-10 5% CO₂ (T. D. T Catheter set, 2583-R106, Laboratoire CCD, France) 3

30 (swim up) (insemination) 5 ?1 , 10~20 ?1

18~22 (fertilization) , 5 ?1

(male pronuclear) (female pronuclear) 0.5~1 cm

(polar body)가

2

1 가 4~5 가

가

Table 1. Comparison of clinical characteristics and outcomes in patients without (Group 1) and with (Group 2) retained embryos at transfer

	Group 1 (n=319)	Group 2 (n=50)
Age	31.7 ±3.1	31.3 ±2.9
Duration of Infertility (yrs)	4.8 ±2.9	5.2 ±2.9
IVF indication		
Uterine factors	8	1
Tubal factors	148	22
Endometriosis	47	12
Ovulatory factors	22	0
Peritoneal factors	2	0
Unexplained	91	15
No. of oocyte retrieval	11.0 ±6.3	13.9 ±7.2*
Fertilization rate (%)	80.9 ±16.4	77.6 ±13.4
GQER (%)	39.4 ±26.0	37.7 ±21.3
No. of embryo transfer	4.8 ±1.1	5.0 ±0.8
Mucus/cycles	24.4 (78/319)	44.0 (22/50)†
Pregnancy rate (%)	45.5 (145/319)	48.0 (24/50)
Implantation rate (%)	34.5 (248/718)	35.3 (42/119)

*† p<0.05

6) 가 Table 1
 (Progesterone in oil, Progest, Samil Pharma, Korea) 50 mg, GQER 가
 13 ?-hCG 1 24.4%
 , ?-hCG 가 (78/319), Group 2 44.0% (22/50) Group 2
 3. (p=0.0038, χ^2 -test). 45.4%, 48.0%
 SAS Pearson's
 χ^2 test Fisher's test (Table 1).
 ,p 0.05 2.

Table 2. Comparison of pregnancy outcome with and without mucus

	Mucus (+) (n=100)	Mucus (-) (n=269)
Age	31.4 ±2.7	31.8 ±3.2
Duration of infertility (yrs)	4.8 ±2.7	4.9 ±2.9
No. of oocytes retrieval	10.9 ±5.4	11.6 ±6.8
Fertilization rates (%)	79.7 ±15.2	80.8 ±16.4
GQER (%)	43.1 ±25.3	37.7 ±25.4
No. of embryo transfer	4.9 ±1.0	4.8 ±1.1
Multiple attempts rates at ET (%)	22.0 (22/100)	8.2 (22/269)*
Pregnancy rates (%)	43.0 (43/100)	46.8 (126/269)
Implantation rates (%)	35.6 (73/205)	34.3 (217/632)

*p<0.05

가 , , GQ- ER, Tomas (2002)¹⁰ multiple logistic regression 가 43.0%, 46.8% 1.7 가 (p=0.5, Table 2). 가 100 22 가 가 16 2 11,12 6,13 8 ,6 3 3 269 28 가 , 가 22 2 10 ,6 가 , 가 . Kovacs (1999)¹⁶ , , , 가 . Goudas (1998)¹⁷

가

Visser (1993)¹ Nabi (1997)⁵
가
, Mansour (1994)²
가 가

(1993)¹ 가 Visser 가
, Nabi (1997)⁵ 가

가 가

44.0% 24.4%

(1997)⁵ Nabi
18 71% 46%, 24%
가 49% (1985)²³ Nazari (1993)²⁴ 가 Yovich
29.6%가 0.5 cm
57% Fanchin 가 가
(1998)¹⁹ 24%, 37% 12, 25
McNamee (1998)²⁰ mock transfer 3
Glass (2000)²¹

가 가

가 가

(Visser et al., 36 가

1993)¹, 가

.Nabi (1997)⁵ Tur-Kasp a (1998)⁶

- more: Williams and Wilkins 1991; 121-49.
1. Visser DS, Fourie FLER, Kruger HF. Multiple attempts at the embryo transfer: effect on pregnancy outcome in an in vitro fertilisation and embryo transfer program. *J Assit Reprod Genet* 1993; 10: 37-43.
 2. Mansour RT, Aboulghar MA, Serour GI, Amin YM. Dummy embryo transfer using methylene blue dye. *Hum Reprod* 1994; 9: 1257-9.
 3. Schoolcraft WB, Surrey ES, Gardner DK. Embryo transfer: techniques and variables affecting success. *Fertil Steril* 2001; 76: 863-70.
 4. Wisanto A, Janssen R, Deschacht J, Camus M, Devroey P, Van Steirteghem AC. Performance of different embryo transfer catheters in a human in vitro fertilization program. *Fertil Steril* 1989; 52: 72-84.
 5. Nabi A, Awonuga A, Birch H, Barlow S, Stewart B. Multiple attempts at the embryo transfer: does this affect in-vitro fertilization treatment outcome? *Hum Reprod* 1997; 12: 1188-90.
 6. Tur-kaspa I, Yuval Y, Bider D, Levron J, Shulman A, Dor J. Difficult or repeated sequential embryo transfers do not adversely affect in-vitro fertilization pregnancy rates or outcome. *Hum Reprod* 1998; 13: 2452-5.
 7. Jung BJ, Song HJ, Oh IH. Can endometriosis affect the clinical outcomes in patients undergoing IVF-ET? *Korea J Fertil Steril* 2002; 29: 223-7.
 8. World Health Organization. Laboratory manual for the examination of human semen and sperm cervical mucus interaction. 3rd ed., Cambridge, Cambridge University Press; 1992.
 9. Veeck LL. Preembryo grading: In atlas of the human oocyte and early embryo conceptus. Vol 2, Baltimore: Williams and Wilkins 1991; 121-49.
 10. Tomas C, Tikkinen K, Tuomivaara L, Tapanainen JS, Martikainen H. The degree of difficulty of embryo transfer is an independent factor for predicting pregnancy. *Hum Reprod* 2002; 17: 2632-5.
 11. Lass A, Abusheikha N, Brinsden P, Kovacs GT. The effect of a difficult embryo transfer on the outcome of IVF. *Hum Reprod* 1999; 14: 2417.
 12. Wood EG, Batzer FR, Go KJ, Gutmann JN, Corson SL. Ultrasound-guided soft catheter embryo transfers will improve pregnancy rates in in-vitro fertilization. *Hum Reprod* 2000; 15: 107-12.
 13. Burke LM, Davenport AT, Russell GB, Deaton JL. Predictors of success after embryo transfer: experience from a single provider. *Am J Obstet Gynecol* 2000; 182: 1001-4.
 14. Woolcott R, Stanger J. Potentially important variables identified by transvaginal ultrasound-guided embryo transfer. *Hum Reprod* 1997; 12: 963-6.
 15. Lesny P, Killick SR, Tetlow RL, Robinson J, Maguiness SD. Embryo transfer--can we learn anything new from the observation of junctional zone contractions? *Hum Reprod* 1998; 13: 1536-40.
 16. Kovacs GT. What factors are important for successful embryo transfer after in-vitro fertilization? *Hum Reprod* 1999; 14: 590-2.
 17. Goudas VT, Hammitt DG, Damario MA, Session DR, Singh AP, et al. Blood on the embryo transfer catheter is associated with decreased rates of embryo implantation and clinical pregnancy with the use of in vitro fertilization-embryo transfer. *Fertil Steril* 1998; 70: 878-82.
 18. Egbase PE, al-Sharhan M, al-Othman S, al-Mutawa M, Udo EE, Grudzinskas JG. Incidence of microbial growth from the tip of the embryo transfer catheter after embryo transfer in relation to clinical pregnancy rate following in-vitro fertilization and embryo transfer. *Hum Reprod* 1996; 11: 1687-9.
 19. Fanchin R, Harmas A, Benaoudia F, Lundkvist U, Olivennes F, Frydman R. Microbial flora of the cervix assessed at the time of embryo transfer adversely

affects in vitro fertilization outcome. *Fertil Steril* 1998; 70: 866-70.

20. McNamee P, Huang T, Carwile A. Significant increase in pregnancy rates achieved by vigorous irrigation of endocervical mucus prior to embryo transfer with a Wallace catheter in an IVF-ET program [abstract]. *Fertil Steril* 1998; 70 Suppl 1: S228.
 21. Glass KB, Green CA, Fluker MR, Schoolcraft WB, McNamee PI, Meldrum D. Multicenter randomized trial of cervical irrigation at the time of embryo transfer [Abstract]. *Fertil Steril* 2000; 74 Suppl 1: S31.
 22. Waterstone J, Curson R, Parsons J. Embryo transfer to low uterine cavity. *Lancet* 1991; 337: 1413.
 23. Yovich JL, Turner SR, Murphy AJ. Embryo transfer technique as a cause of ectopic pregnancies in in vitro fertilization. *Fertil Steril* 1985; 44: 318-21.
 24. Nazari A, Askari HA, Check JH, O'Shaughnessy A. Embryo transfer technique as a cause of ectopic pregnancy in in vitro fertilization. *Fertil Steril* 1993; 60: 919-21.
 25. Coroleu B, Carreras O, Veiga A, Martell A, Martinez F, Belil I, et al. Embryo transfer under ultrasound guidance improves pregnancy rates after in vitro fertilization. *Hum Reprod* 2001; 15: 616-20.
-