

Cytotoxic Coumarins from the Root of *Angelica dahurica*

Pham Ngoc Thanh, WenYi Jin, GyuYong Song, KiHwan Bae, and Sam Sik Kang¹

College of Pharmacy, Chungnam National University, Daejeon 305-764, Korea and ¹Natural Products Research Institute, Seoul National University, Seoul 110-460, Korea

(Received October 11, 2004)

Ten coumarins were isolated from the root of *Angelica dahurica* by repeated silica gel column chromatography. Their chemical structures were elucidated on the basis of physicochemical and spectroscopic data. Among them, oxypeucedanin hydrate acetonide (**7**) was isolated for the first time from this plant. Cytotoxicity of coumarins isolated were determined *in vitro* against L1210, HL-60, K562, and B16F10 tumor cell lines by MTT method. Pangelin (**5**) and oxypeucedanin hydrate acetonide (**7**) showed a potent cytotoxic activity with the IC₅₀ values of 8.6 to 14.6 µg/mL against four kinds of tumor cell lines. Other compounds showed the moderate cytotoxic activity or no activity against the tumor cell lines.

Key words: *Angelica dahurica*, Coumarins, Pangelin, Oxypeucedanin hydrate acetonide, Tumor cells, Cytotoxicity

INTRODUCTION

Radix Angelicae Dahuricae, the dried root of *Angelica dahurica* Benth et Hook (Umbelliferae) has long been used in Chinese medicine. It has been frequently prescribed as a sedative and an analgesic (Soka, 1985). As a part of our continuing studies to identify novel antitumor agents from natural sources, the CH₂Cl₂ fraction from MeOH extract of *A. dahurica* was found to have the potent cytotoxic activity in L1210 tumor cell line by MTT method (Mosmann, 1983). Several coumarins, a polyacetylene, and essential oil (Fujiwara *et al.*, 1980; Zhang *et al.*, 1980; Kozawa *et al.*, 1981; Zhou *et al.*, 1987; Kim *et al.*, 1992; Qiao *et al.*, 1996; Lechner *et al.*, 2004; Qiao *et al.*, 1997), and many bioactivities such as anti-staphylococcal activity, anti-inflammatory activity, hepatoprotective activity, anti-oxidative activity, antimicrobial activity, antimutagenic activity (Lechner *et al.*, 2004; Lin *et al.*, 2002; Oh *et al.*, 2002; Tsai *et al.*, 1997; Kwon *et al.*, 1997) have been reported on the root of *A. dahurica*. But no detailed research has been reported about the cytotoxic activity. In this paper, we report the isolation, structure elucidation of ten coumarins, which include one first isolated compound and cytotoxic activity against L1210, HL-60, K562, and B16F10 tumor cell lines.

MATERIALS AND METHODS

General experimental procedures

Melting points were measured on an electrothermal melting point apparatus, and are uncorrected. The IR spectra were recorded on a Jasco Report-100 IR spectrometer. ¹H- and ¹³C-NMR spectra were recorded with BRUKER AVANCE 300 NMR spectrometer in CDCl₃ using TMS as the internal standard. MS spectra were measured with JEOL JMSAX505WA mass spectrometer. Column chromatography was carried out using silica gel (Merck, 70-230 mesh). The TLC was performed on a precoated Merck Kieselgel 60 F₂₅₄ plate (0.25 mm) and a Merck RP-18 F₂₅₄ plate (5×10 cm).

Plant material

The root of *Angelica dahurica* were purchased from herb market in Taejeon and the voucher specimen (CNU 1513) were deposited at the herbarium in the College of Pharmacy, Chungnam National University.

Extraction and isolation

The dried root (5 kg) was ground and extracted 3 times with MeOH and obtained the MeOH extract (413 g). The extract was diluted with H₂O to give an aqueous solution. The aqueous solution was subsequently partitioned into CH₂Cl₂ and aqueous fractions. The CH₂Cl₂ extract (187 g) was chromatographed on a silica gel using stepwise gradient elution with the solvents hexane-EtOAc to give 7

Correspondence to: KiHwan Bae, College of Pharmacy, Chungnam National University, Daejeon 305-764, Korea
Tel: 82-42-821-5925 Fax: 82-42-823-6566
E-mail: baekh@cnu.ac.kr

fractions (AD-1~AD-7). Silica gel column chromatography of fraction AD-2 (18 g) eluting with hexane-EtOAc (5:1) gave two subfractions (AD-2-1~AD-2-2). Subfraction AD-2-1 was subjected to silica gel column eluting with hexane-EtOAc (6:1) afford compound **1** (174 mg). Subfraction AD-2-2 was subjected to silica gel column eluting with hexane-EtOAc (4:1) to afford compounds **2** (84 mg) and **3** (126 mg). Fraction AD-3 (32 g) was subjected to silica gel column eluting with hexane-EtOAc (5:1) to afford compounds **4** (21 mg), **5** (18 mg), and **6** (34 mg). Silica gel column chromatography of fraction AD-4 (24 g) eluting with hexane-EtOAc (3:1) afforded three subfractions (AD-4-1~AD-4-3). Subfraction AD-4-1 was subjected to silica gel column eluting with hexane-acetone (5:1) to afford compound **7** (9 mg). Subfraction AD-4-2 was subjected to silica gel column eluting with hexane-EtOAc (3:1) to afford compound **8** (63 mg). Silica gel column chromatography of fraction AD-5 (16 g) eluting with hexane-EtOAc (2:1) afforded three subfractions (AD-5-1~AD-5-3). Subfraction AD-5-2 was subjected to silica gel column eluting with hexane-acetone (3:1) to afford compound **9** (56 mg). Subfraction AD-5-3 was subjected to silica gel column eluting with hexane-EtOAc (1:1) to afford compound **10** (18 mg).

Isoimperatorin (1)

White crystal, mp. 105-107°C, IR ν_{\max} cm^{-1} : 1729 (C=O), 1628, 1604 (aromatic ring, C=C), 1129 (benzofuran), 821, $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 1.82 (3H, s, H-6"), 1.72 (3H, s, H-5"), 4.94 (2H, d, $J = 6.9$ Hz, H-2"), 5.56 (1H, t-like, $J = 6.9$ Hz, H-3"), 6.29 (1H, d, $J = 9.6$ Hz, H-3), 6.97 (1H, dd, $J = 0.9, 2.1$ Hz, H-3'), 7.17 (1H, br.s, H-8), 7.61 (1H, d, $J = 2.1$ Hz, H-2'), 8.17 (1H, d, $J = 9.6$ Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 18.6 (C-6"), 26.2 (C-5"), 70.2 (C-2"), 94.7 (C-8), 105.4 (C-3'), 108.0 (C-10), 113.0 (C-3), 114.7 (C-6), 119.5 (C-2"), 139.9 (C-4), 140.2 (C-4"), 145.3 (C-2'), 149.4 (C-5), 153.1 (C-9), 158.5 (C-7), 161.7 (C-2).

Osthohol (2)

White crystal, mp. 82-83°C, IR ν_{\max} cm^{-1} : 1720 (C=O), 1600, 1435 (aromatic ring, C=C), 1110 (benzofuran), $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : 1.68 (3H, s, H-5'), 1.85 (3H, s, H-4'), 3.54 (2H, d, $J = 7.2$ Hz, H-1'), 3.93 (3H, s, 7-OMe), 5.24 (1H, t, $J = 7.2$ Hz, H-2'), 6.23 (1H, d, $J = 9.6$ Hz, H-3), 6.84 (1H, d, $J = 8.4$ Hz, H-6), 7.29 (1H, d, $J = 8.4$ Hz, H-5), 7.61 (1H, d, $J = 9.6$ Hz, H-4), $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ : 161.7 (C-2), 160.6 (C-7), 153.2 (C-9), 144.1 (C-4), 133.0 (C-3'), 126.6 (C-5), 121.6 (C-2'), 118.4 (C-8), 113.4 (C-10), 113.4 (C-3), 107.8 (C-6), 56.4 (7-OMe), 26.2 (C-5'), 22.3 (C-1'), 18.3 (C-4').

Imperatorin (3)

White crystal, mp. 102, IR ν_{\max} cm^{-1} : 1722 (C=O), 1587

(aromatic ring, C=C), 1150 (benzofuran), 838, $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : 1.74 (3H, s, H-6"), 1.76 (3H, s, H-5"), 5.02 (2H, d, $J = 6.9$ Hz, H-2"), 5.60 (1H, t-like, $J = 6.9$ Hz, H-3"), 6.37 (1H, d, $J = 9.6$ Hz, H-3), 6.83 (1H, dd, $J = 0.9, 2.1$ Hz, H-3'), 7.36 (1H, s, H-5), 7.70 (1H, d, $J = 2.1$ Hz, H-2'), 7.76 (1H, d, $J = 9.6$ Hz, H-4). $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ : 18.5 (C-5"), 26.2 (C-6"), 70.5 (C-2"), 107.1 (C-3'), 113.1 (C-5), 115.1 (C-3), 116.9 (C-10), 120.2 (C-3'), 126.2 (C-6), 132.1 (C-8), 140.1 (C-4"), 144.3 (C-9), 144.7 (C-4), 147.0 (C-2'), 149.0 (C-7), 160.9 (C-2).

Psoralen (4)

White crystal, mp. 164-165°C, IR ν_{\max} cm^{-1} : 1725 (C=O), 1635, 1577 (aromatic ring, C=C), 1136 (benzofuran), 824, $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 6.39 (1H, d, $J = 9.6$ Hz, H-3), 6.85 (1H, dd, $J = 0.9, 2.4$ Hz, H-3'), 7.49 (1H, br.s, H-8), 7.70 (1H, s, H-5), 7.71 (1H, d, $J = 2.4$ Hz, H-2'), 7.81 (1H, d, $J = 9.6$ Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 100.3 (C-8), 106.8 (C-3'), 115.1 (C-3), 115.8 (C-10), 120.2 (C-5), 125.3 (C-6), 144.4 (C-4), 147.3 (C-2'), 152.5 (C-9), 156.8 (C-7), 161.4 (C-2).

Pangelin (5)

White crystal, mp. 135-136, IR ν_{\max} cm^{-1} : 3649 (OH), 1717 (C=O), 1624 (aromatic ring, C=C), 1134 (benzofuran), $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : 8.19 (1H, dd, $J = 9.6$ Hz, H-4), 7.61 (1H, d, $J = 2.4$ Hz, H-2'), 7.18 (1H, s, H-8), 6.98 (1H, dd, $J = 0.9, 2.4$ Hz, H-3'), 6.30 (1H, d, $J = 9.6$ Hz, H-3), 5.22 (1H, d, $J = 0.9$ Hz, H-5"), 4.55 (1H, m, H-3"), 4.48 (1H, dd, $J = 9.6, 3.6$ Hz, H-2"), 4.41 (1H, dd, $J = 9.6, 7.2$ Hz, H-2"), 2.33 (1H, s, 3"-OH), 1.86 (3H, s, H-6"), $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ : 161.4 (C-2), 158.5 (C-7), 153.0 (C-9), 148.9 (C-5), 145.6 (C-2'), 143.7 (C-4"), 139.5 (C-4), 114.6 (C-6), 113.8 (C-5"), 113.4 (C-3), 107.8 (C-10), 105.1 (C-3'), 95.1 (C-8), 76.1 (C-2"), 74.6 (C-3"), 19.1 (C-6").

Bergapten (6)

White crystal, mp. 187-188°C, IR ν_{\max} cm^{-1} : 1731 (C=O), 1625, 1580 (aromatic ring, C=C), 1124 (benzofuran), 834, $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 4.28 (3H, s, 5-OCH₃), 6.28 (1H, d, $J = 9.9$ Hz, H-3), 7.03 (1H, d, $J = 2.4$ Hz, H-3'), 7.15 (1H, s, H-8), 7.61 (1H, d, $J = 2.4$ Hz, H-2'), 8.16 (1H, d, $J = 9.9$ Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 60.5 (5-OCH₃), 94.3 (C-8), 105.4 (C-3'), 106.8 (C-10), 113.0 (C-3), 113.1 (C-6), 139.6 (C-4), 145.2 (C-2'), 150.0 (C-5), 153.1 (C-9), 158.8 (C-7), 161.6 (C-2).

Oxypeucedanin hydrate acetonide (7)

White crystal, mp. 157-158°C, FAB-MS m/z : 345 $[\text{M}+\text{H}]^+$, IR ν_{\max} cm^{-1} : 1736 (C=O), 1620 (aromatic ring, C=C), 1370, 1340 (C-O), 1120 (benzofuran), 820, $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : 8.19 (1H, d, $J = 9.6$ Hz, H-4), 7.63 (1H, d, $J = 2.4$ Hz, H-2'), 7.20 (1H, s, H-8), 7.00 (1H, dd, $J = 0.9,$

2.4 Hz, H-3'), 6.30(1H, d, $J = 9.6$ Hz, H-3), 4.55 (1H, dd, $J = 7.2, 9.9$ Hz, H-2''), 4.43 (1H, dd, $J = 4.5, 9.9$ Hz, H-2''), 4.24 (1H, dd, $J = 4.5, 7.2$ Hz, H-3''), 1.51 (3H, s, H-8''), 1.45 (3H, s, H-9''), 1.43 (3H, s, H-10''), 1.23 (3H, s, H-11''), $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ : 161.3 (C-2), 158.5 (C-7), 153.0 (C-9), 148.8 (C-5), 145.6 (C-2'), 139.4 (C-4), 114.5 (C-6), 113.5 (C-3), 108.7 (C-5''), 107.7 (C-10), 105.1 (C-3'), 95.2 (C-8), 81.7 (C-3''), 80.0 (C-7''), 72.4 (C-2''), 28.9 (C-8''), 27.4 (C-9''), 27.2 (C-10''), 23.4 (C-11'').

Oxypeucedanin (8)

White crystal, mp. 142-143°C, IR ν_{max} cm^{-1} : 1720 (C=O), 1602 (aromatic ring, C=C), 1386, 1362 (C-O), $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 1.35 (3H, s, H-6''), 1.43 (3H, s, H-5''), 3.25 (1H, dd, $J = 4.5, 6.6$ Hz, H-3''), 4.45 (1H, dd, $J = 6.6, 10.8$ Hz, H-2a''), 4.61 (1H, dd, $J = 4.5, 10.8$ Hz, H-2b''), 6.33 (1H, d, $J = 9.6$ Hz, H-3), 6.96 (1H, dd, $J = 0.9, 2.4$ Hz, H-3'), 7.22 (1H, br.s, H-8), 7.63 (1H, d, $J = 2.4$ Hz, H-2'), 8.22 (1H, d, $J = 9.6$ Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 19.4 (C-6''), 25.0 (C-5''), 58.7 (C-4''), 61.5 (C-3''), 72.8 (C-2''), 95.4 (C-8), 104.9 (C-3'), 107.9 (C-10), 113.6 (C-3), 114.7 (C-6), 139.3 (C-4), 145.7 (C-2'), 148.8 (C-5), 153.0 (C-9), 158.6 (C-7), 161.4 (C-2).

Xanthotoxin (9)

White crystal, mp. 146-147°C, IR ν_{max} cm^{-1} : 1710 (C=O), 1620, 1586 (aromatic ring, C=C), 1155 (benzofuran), 821, $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 4.29 (3H, s, 8-OMe), 6.38 (1H, d, 9.6 Hz, H-3), 6.83 (1H, d, 2.1 Hz, H-3'), 7.36 (1H, s, H-5), 7.70 (1H, d, 2.1 Hz, H-2'), 7.77 (1H, d, 9.6 Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 61.7 (8-OMe), 107.1 (C-3'), 113.3 (C-5), 115.2 (C-3), 116.9 (C-10), 126.5 (C-6), 133.2 (C-8), 143.4 (C-9), 144.7 (C-4), 147.0 (C-2'), 148.1 (C-7), 160.8 (C-2).

Oxypeucedanin hydrate (10)

Pale yellowish crystal, mp. 134-135, IR ν_{max} cm^{-1} : 3400 (OH), 1716 (C=O), 1604 (aromatic ring, C=C), 1389, 1370 (C-O), $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 1.33 (3H, s, H-6''), 1.38 (3H, s, H-5''), 2.18 (1H, s, 4''-OH), 3.04 (1H, s, 3''-OH), 3.93 (1H, dd, $J = 3.0, 7.8$ Hz, H-3''), 4.45 (1H, dd, $J = 7.8, 9.6$ Hz, H-2a''), 4.57 (1H, dd, $J = 3.0, 9.6$ Hz, H-2b''), 6.26 (1H, d, $J = 9.9$ Hz, H-3), 7.00 (1H, dd, $J = 0.9, 2.4$ Hz, H-3'), 7.13 (1H, br.s, H-8), 7.61 (1H, d, $J = 2.4$ Hz, H-2'), 8.17 (1H, d, $J = 9.9$ Hz, H-4), $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ : 25.6 (C-6''), 27.0 (C-5''), 72.1 (C-4''), 74.9 (C-2''), 77.0 (C-3''), 95.1 (C-8), 105.2 (C-3'), 107.6 (C-10), 113.3 (C-3), 114.6 (C-6), 139.5 (C-4), 145.6 (C-2'), 148.9 (C-5), 152.9 (C-9), 158.5 (C-7), 161.6 (C-2).

Tumor cells

The murine leukemia L1210 cell, human leukemia HL-60 cell, human leukemia K562 cell, and murine melanoma

B16F10 cell were supplied by Dr. HyeongKyu Lee at Immunomodulator Research Laboratory in KRIBB. Cells were maintained in RPMI 1640 including L-glutamine (JBI) with 10% FBS (JBI) and 2% penicillin-streptomycin (GIBCO). Trypsin-EDTA was used in separating cell from culture flask. All the cell lines were cultured at 37 in an atmosphere of 5% CO_2 incubator. The confluent cells were used for the MTT assay.

Cytotoxicity assay

Cytotoxicity was measured by a modification of the Microculture Tetrazolium (MTT) assay (Mosmann, 1983). Viable cells were seeded in the growth medium (180 μL) into 96 well microtiter plate (1×10^4 cells per each well) and allowed to attach in 37°C, 5% CO_2 incubator. The test sample was dissolved in DMSO and adjusted to the final sample concentrations ranging from 30 $\mu\text{g}/\text{mL}$ to 1.875 $\mu\text{g}/\text{mL}$ by diluting with the growth medium. Each sample was prepared in triplicate. The final DMSO concentration was adjusted to <0.1%. After allowing 2 h, 20 μL test samples were added to each wells the same concentration DMSO were added in the control group. 48 h after test sample addition, 20 μL MTT (final concentration, 5 mg/mL) was added to the each wells. Two hours later, the plate was centrifuged for 5 minutes in 1500 rpm, the medium was removed and formed formazan crystals were dissolved with 150 μL DMSO. The optical density (O.D.) was measured at 570 nm using a Titertek microplate reader (Multiskan MCC/340, Flow). The IC_{50} value was defined as the concentration of sample needed to reduce a 50% of absorbance relative to the vehicle-treated control.

RESULTS AND DISCUSSION

Repeated silica gel column chromatography of the CH_2Cl_2 fraction of the MeOH extract from *A. dahurica* (root) led to the isolation of ten coumarins (1-10). Compounds of 1-6 and 8-10 were identified as isoimperatorin (1), osthol (2), imperatorin (3), psoralen (4), pangelin (5), bergapten (6), oxypeucedanin (8), xanthotoxin (9), and oxypeucedanin hydrate (10) by comparing their spectral and physical data with those previously reported (Saiki *et al.*, 1971; Razdan *et al.*, 1987; Fujioka *et al.*, 1999; Stavri *et al.*, 2003).

Compound 7 was obtained as a white crystal. An $[\text{M}+\text{H}]^+$ peak at 345 in FAB-MS along with the analysis of $^{13}\text{C-NMR}$, DEPT spectra showed its molecular formula to be $\text{C}_{19}\text{H}_{20}\text{O}_6$. In the $^1\text{H-NMR}$ spectrum of compound 7, two proton doublet signals (1H, $J = 9.6$ Hz) at δ 6.30 and 8.19 were assignable to be protons of pyrone ring. A proton doublet signal (1H, $J = 2.4$ Hz) at δ 7.63 and another proton double of doublet signal (1H, $J = 0.9, 2.4$




Fig. 1. Structure of compounds 1-10 from the root of *Angelica dahurica*

Hz) at δ 7.00 were attributable to the protons of furan ring. The signal for H-4 was observed rather lower field (δ 8.19) to indicate the absence of proton at C-5 (Harkar *et al.*, 1984). And the signal for H-3' was observed as a doublet ($J = 0.9, 2.4$ Hz) to mean the occurrence of long-range coupling with H-8 because long-range coupling between H-3' and H-5 in the linear furanocoumarin skeleton was not usually observed (Sasaki *et al.*, 1982). Therefore, a side chain was suggested to be substituted at C-5 position. The presence of (2,2,5,5-tetramethyl-1,3-

Table I. Cytotoxicity of compounds 1-10 from *Angelica dahurica*

Compounds	IC ₅₀ (μ g/mL) ^a			
	L1210	HL-60	K562	B16F10
1	24.5	22.3	20.1	23.5
2	26.4	15.5	25.2	28.2
3	>30	20.2	>30	>30
4	>30	28.9	19.3	>30
5	13.3	14.6	10.1	12.5
6	>30	>30	>30	>30
7	9.4	9.5	8.6	9.8
8	>30	27.5	>30	>30
9	28.9	16.7	24.5	25.6
10	>30	>30	>30	>30
AM ^b	0.8	2.8	1.4	0.9

^a IC₅₀ value was the 50% inhibition concentration and calculated from regression lines using five different concentrations in triplicate experiments.

^b Adriamycin used in positive control.

dioxolan-4-yl)methoxy unit was confirmed by the proton signals observed at δ 4.55 and δ 4.43 (H-2''), δ 4.24 (H-3''), δ 1.51, δ 1.45, δ 1.43, and δ 1.23 (H-8'', 9'', 10'', and 11''). Based on these results and on values previously reported in the literature (Abyshev *et al.*, 1979), compound **7** was identified as oxypeucedanin hydrate acetone. It has been isolated from this plant for the first time.

The cytotoxicity assay was carried out against four cancer cell lines, L1210 (murine leukemia cancer), HL-60 (human leukemia cancer), K562 (human leukemia cancer), and B16F10 (murine melanoma cancer) according to the MTT assay described previously (Mosmann, 1983). The results were summarized in Table I. There is no relationship between the cytotoxicity and the structure of isolated coumarins. Among the tested coumarins isolated from the CH₂Cl₂ fraction of *A. dahurica*, compounds **5** and **7** exhibited the most potent cytotoxic activity against all tumor cell lines with IC₅₀ values ranging between 8.6 to 14.6 μ g/mL while compounds **6** and **10** did not exhibit any cytotoxic activity (IC₅₀, > 30 μ g/mL). Up to now there is no report on cytotoxicity of compounds **5** and **7** against tumor cell lines. Compounds **1**, **2**, and **9** exhibited moderate to weak cytotoxic activity against all tumor cell lines with IC₅₀ values ranging between 15.5 to 28.9 μ g/mL. According to the previous data (Yang *et al.*, 2003), compounds **1**, **2**, and **9** showed moderate to weak cytotoxic effect against HL-60, P-388, CoLo 205 and Hela tumor cell lines with IC₅₀ values ranging between 14.9 to 35.3 μ g/mL, and in accordance with our data. Interestingly, compound **3** and **8** showed weak cytotoxic activity (IC₅₀, 20.2 and 27.5 μ g/mL) against only HL-60 cell line, respectively. Also, com-

pound 4 showed weak cytotoxic activity against only HL-60 and K562 cell lines with IC₅₀ values of 28.9 and 19.3 µg/ml, respectively.

These results suggest that the main cytotoxic of the fraction of *A. dahurica* might be attributed to the pangelin (5) and oxypeucedanin hydrate acetone (7).

ACKNOWLEDGEMENTS

This research was supported by a grant of the Korea Health 21 R & D Project, Ministry of Health & Welfare, Republic of Korea (01-PJ2-PG6-01NA01-0002). We are grateful to Korea Basic Science Institute (KBSI) for supplying the NMR spectra, and also grateful to Dr. Hyeongkyu Lee at Immunomodulator Research Laboratory in KRIBB for supplying tumor cell lines.

REFERENCES

- Abyshev, A. Z., Azhdarov, B., and Gashimov, N. F., Acetone of oxypeucedanin hydrate as a new component from *Peucedanum turcomanicum*. *Khimiya Prirodnykh Soedinenii*, 6, 847-848 (1979).
- Fujioka, T., Furumi, K., Fujii, H., Okabe, H., Mihashi, K., Nakano, Y., Matsunaga, H., Katano, M., and Mori M., Antiproliferative constituents from Umbelliferae plants. V. A new furanocoumarin and falcarindiol furanocoumarin ethers from the root of *Angelica japonica*. *Chem. Pharm. Bull.*, 47, 96-100 (1999).
- Fujiwara, H., Yokoi, T., Tani, S., Saiki, Y., and Kato, A., Studies on constituents of *Angelicae dahuricae* Radix. I. On a new furocoumarin derivative. *Yakugaku Zasshi*, 100, 1258-1261 (1980).
- Harkar, S., Razdan, T. K., and Waight, E. S., Steroids, chromone, and coumarins from *Angelica officinalis*. *Phytochemistry*, 23, 419-426 (1984).
- Kim, S. H., Kang, S. S., and Kim, C. M., Coumarin glycosides from the roots of *Angelica dahurica*. *Arch. Pharm. Res.*, 15, 73-77 (1992).
- Kozawa, M., Baba, K., Okuda, K., Fukumoto, T., and Hata, K., Studies on chemical components of "Bai Zhi". On coumarins from "Japanese Bai Zhi". *Shoyakugaku Zasshi*, 35, 90-95 (1981).
- Kwon, Y. S., Kobayashi, A., Kajiyama, S. I., Kawazu, K., Kanzaki, H., and Kim, C. M., Antimicrobial constituents of *Angelica dahurica* roots. *Phytochemistry*, 44, 887-889 (1997).
- Lechner, D., Stavri, M., Oluwatuyi, M., Pereda-Miranda, R., and Gibbons, S., The anti-staphylococcal activity of *Angelica dahurica* (Bai Zhi). *Phytochemistry* 65, 331-335 (2004).
- Lin, C. H., Chang, C. W., Wang, C. C., Chang, M. S., and Yang, L. L., Byakangelicol, isolated from *Angelica dahurica*, inhibits both the activity and introduction of cyclooxygenase-2 in human pulmonary epithelial cells. *J. Pharm. Pharmacol.*, 54, 1271-1278 (2002).
- Mosmann T., Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods*, 65, 55-63 (1983).
- Oh, H., Lee, H. S., Kim, T., Chai, K. Y., Chung, H. T., Kwon, T. O., Jun, J. Y., Jeong, O. S., Kim, Y. C., and Yun, Y. G., Furocoumarin from *Angelica dahurica* with hepatoprotective activity on tacrine-induced cytotoxicity in Hep G2 cell. *Planta Med.*, 68, 463-464 (2002).
- Qiao, S., Yao, X., Liu, C., and Li, Y., Chemical constituents of essential oil from root of *Angelica dahurica*. *Zhongguo Yaowu Huaxue Zazhi*, 7, 200-201, 222, (1997).
- Qiao, S. Y., Yao, X. S., and Wang, Z. Y., Coumarins of the roots of *Angelica dahurica*. *Planta Med.*, 62, 584 (1996).
- Razdan, T. K., Qadri, B., Harkar, S., and Waight, E. S., Chromones and coumarins from *Skimmia laureola*. *Phytochemistry*, 26, 2063-2069 (1987).
- Saiki, Y., Morinaga, K., Okegawa, O., Sakai, S., Amaya, Y., Ueno, A., and Fukushima, S., Coumarins of the roots of *Angelica dahurica*. *Yakugaku Zasshi*, 91, 1313-1316 (1971).
- Sasaki, H., Taguchi, H., Endo, T., and Yosioka, I., The constituents of *Ledebouriella seseloides* Wolff. I. Structures of three new chromones. *Chem. Pharm. Bull.*, 30, 3555-3562 (1982).
- Soka, T., *Dictionary of Chinese Drug*. Shanghai Science Technology Shogakukan (Eds.) Shogakukan Press, Tokyo, pp. 2243-2246 (1985).
- Stavri, M., Mathew, K. T., Bucar, F., and Gibbons, S., Pangelin, an antimycobacterial coumarin from *Ducrosia anethifolia*. *Planta Med.*, 69, 956-959 (2003).
- Tsai, G. C. and Yang, L. L., Antioxidative principles of *Angelica dahurica* var. *pai chi*. *Taiwan Kexue*, 50, 139-153 (1997).
- Yang, L. L., Wang, M. C., Chen, L. G., and Wang, C. C., Cytotoxic activity of coumarins from the fruits of *Cnidium monnieri* on leukemia cell lines. *Planta Med.*, 69, 1091-1095 (2003).
- Zhang, H. Q., Yuan, C. Q., Chen, G. Y., Ding, Y. M., Chen, S. Q., and Deng, Y. Q., Study on the chemical constituents of the root of *Angelica dahurica* var. *formosana*. *Yaouxue Tongbao*, 15, 2-4 (1980).
- Zhou, J., Yu, C., and Hang, Y., Studies on Baizhi (*Angelica dahurica*) V. Studies on the chemical constituents. *Zhongcaoyao*, 18, 242-246 (1987).