

STUDY ON BROWDER'S SPECTRUMS AND WEYL'S SPECTRUMS

DONG HARK LEE

ABSTRACT. In this paper we give several necessary and sufficient conditions for an operator on the Hilbert space H to obey Browder's theorem. And it is shown that if S has totally finite ascent and $T \prec S$ then $f(T)$ obeys Browder's theorem for every $f \in H(\sigma(T))$, where $H(\sigma(T))$ denotes the set of all analytic functions on an open neighborhood of $\sigma(T)$.

Furthermore, it is shown that if $T \in B(H)$ is a compact operator or a Riesz Operator then T obeys Browder's theorem and Weyl's theorem holds if and only if Browder's holds.

1. Introduction

Throughout this note let $B(H)$ and $K(H)$ denote respectively the algebra of bounded linear operators and the ideal of compact operators acting on an infinite dimensional Hilbert space H .

If $T \in B(H)$ write $N(T)$ and $R(T)$ for the null space and the range of T ; $\alpha(T) = \dim N(T)$; $\beta(T) = \text{codim} R(T)$; $\sigma(T)$ for the spectrum of T ; $\pi_0(T)$ for the set of eigenvalues of T ; $\pi_{0f}(T)$ for the eigenvalues of finite multiplicity; $\pi_0^{left}(T)$ for the isolated points of $\sigma(T)$; which are eigenvalues of finite multiplicity; $\pi_{00}(T) = \sigma(T) \setminus \sigma_b(T)$ for the Riesz Points of T .

An operator $T \in B(H)$ is called Fredholm if it has closed range with finite dimensional null space and its range of finite co-dimension.

The index of a Fredholm Operator $T \in B(H)$ is given by $i(T) = \alpha(T) - \beta(T)$.

An operator $T \in B(H)$ is called Weyl if it is Fredholm of index zero. An operator $T \in B(H)$ is called Browder if it is Fredholm "of finite

Received May 27, 2004. Revised August 22, 2004.

2000 Mathematics Subject Classification: 47A53 .

Key words and phrases: Browder's spectrums, Weyl's spectrums.

This study was supported by 2002 Kangwon National University Research Fund.

ascent and descent": equivalently ([9, Theorem 7.9.3]) if T is Fredholm and $T - \lambda I$ is invertible for sufficiently small $\lambda \neq 0$ in \mathbb{C} .

The essential spectrum $\sigma_e(T)$, the Weyl's spectrum $w(T)$, the Browder's spectrum $\sigma_b(T)$, the regular spectrum $\sigma_r(T)$ of $T \in B(H)$ are defined by ([8, 9]):

$$\begin{aligned}\sigma_e(T) &= \{ \lambda \in \mathbb{C}; T - \lambda I \text{ is not Fredholm } \}; \\ w(T) &= \{ \lambda \in \mathbb{C}; T - \lambda I \text{ is not Weyl } \}; \\ \sigma_b(T) &= \{ \lambda \in \mathbb{C}; T - \lambda I \text{ is not Browder } \}; \\ \sigma_r(T) &= \{ \lambda \in \mathbb{C}; T - \lambda I \text{ is not regular } \};\end{aligned}$$

By [9, Theorem 6.4.2] $\sigma_r(T) \subset \sigma_e(T)$.

Evidently $\sigma_r(T) \subseteq \sigma_e(T) \subseteq w(T) \subseteq (\sigma_b(T) \cap \sigma_e(T)) \cup acc\sigma(T)$ where we write $accK$ for the accumulation points of $K \subseteq \mathbb{C}$.

We say Weyl's theorem holds for $T \in B(H)$ if there is equality (1.1) $\sigma(T) \setminus w(T) = \pi_0^{left}(T)$ and that Browder's theorem holds for $T \in B(H)$ if there is equality (1.2) $\sigma(T) \setminus \sigma_b(T) = \pi_{00}(T)$.

An operator $T \in B(H)$ is a **Gm**-operator ($m \geq 1$) if there exists a constant M such that $\|(T - \lambda I)\| \leq \frac{M}{(d(\lambda, \sigma(T)))^m}$ for every $\lambda \notin \sigma(T)$.

The condition $N\lambda$ is said to be satisfied at a particular λ if $N(T - \lambda I) \cap N[((T - \lambda I)^*)^n]$ is nontrivial for some positive integer n , which may depend on λ . An operator $T \in B(H)$ is said to be dominant if for every $\lambda \in \mathbb{C}$ there exists a constant M such that $(T - \lambda I)(T - \lambda I)^* \leq M\lambda(T - \lambda I)^*(T - \lambda I)$ and an operator $T \in B(H)$ is said to be paranormal if $\|Tx\|^2 \leq \|T^2x\|\|x\|$ for all $x \in H$.

In Particular, T is called totally paranormal if $T - \lambda I$ is paranormal for every $\lambda \in \mathbb{C}$.

$T \in B(H)$ is called a quasiaffinity if it has trivial kernel and dense range.

$S \in B(H)$ is said to be a quasiaffine transform of $T \in B(H)$ (notation; $S \prec T$) if there is a quasiaffinity $K \in B(H)$ such that $KS = TK$.

If both $S \prec T$ and $T \prec S$; then we say that S and T are quasisimilar.

An operator $T \in B(H)$ has totally finite ascent if $T - \lambda I$ has finite ascent for each $\lambda \in \mathbb{C}$.

It is known ([10]) that if $T \in B(H)$ then we have; Weyl's theorem \Rightarrow Browder's theorem.

If $T \in B(H)$ is a compact operator then it is proved that Weyl's theorem holds if and only if Browder's theorem holds where $T \in B(H)$

is a compact operator on H if for any bounded sequence $\langle x_n \rangle_{n=1}^\infty$ in H , the sequence $\langle Tx_n \rangle_{n=1}^\infty$ has a convergent subsequence $\langle Tx_{n_k} \rangle_{k=1}^\infty$ such that this limit $\lim_{k \rightarrow \infty} Tx_{n_k} = T_{x_0} \in H$.

Example: Let $l_2 = \{ \langle x_n \rangle_{n=1}^\infty : \text{sequences in } \mathbb{C}, | \sum_{n=1}^\infty |x_n|^2 < +\infty \}$ be a Hilbert space.

Consider $T = B(l_2)$ defined by $T(\langle x_n \rangle_{n=1}^\infty) = \langle \frac{x_n}{n} \rangle_{n=1}^\infty$. Then by [18, Theorem 2], T is a compact operator on l_2 . Since $Tx_n = \frac{x_n}{n}$, we have $\sigma(T) = \{ \frac{1}{n} | n \in \mathbb{N} \} \cup \{0\}$.

2. Main result

THEOREM 2.1. *Let $T \in B(H)$. Then the following statements are equivalent:*

- (1) T obeys Browder's theorem;
- (2) $\sigma(T) \setminus w(T) \subseteq \text{iso}\sigma(T)$;
- (3) $\gamma T(\lambda)$ is discontinuous for each $\lambda \in \sigma(T) \setminus w(T)$, where $\gamma T(\cdot)$ denotes the reduced minimum modulus;
- (4) Every $\lambda \in \sigma(T) \setminus w(T)$ satisfies the condition $N\lambda$;
- (5) $T - \lambda I$ has finite ascent for each $\lambda \in \sigma(T) \setminus w(T)$.

Proof. (1) \Leftrightarrow (2) :If T obeys Browder's theorem then $\sigma(T) \setminus w(T) = \pi_{00}(T) \subseteq \text{iso}\sigma(T)$. Conversely, suppose that $\lambda \in \sigma(T) \setminus w(T)$. Then $T - \lambda I$ is Weyl. But $\lambda \in \text{iso}\sigma(T)$; hence by the punctured neighborhood theorem $\lambda \in \sigma(T) \setminus \sigma_b(T)$. Therefore T obeys Browder's theorem.

(1) \Leftrightarrow (3): If T obeys Browder's theorem then it follows from [6, Lemma 5.52] that $\gamma T(\lambda)$ is discontinuous for each $\lambda \in \sigma(T) \setminus w(T)$. Conversely, suppose that $\gamma T(\lambda)$ is discontinuous for each $\lambda \in \sigma(T) \setminus w(T)$. Let $\lambda_0 \in \sigma(T) \setminus w(T)$. Then $T - \lambda_0 I$ is Weyl and $\alpha(T - \lambda_0 I) > 0$. Therefore $\gamma T(\lambda) > 0$ for all λ near λ_0 , and so by [6, Corollary 5.74] $\alpha(T - \lambda I) < \alpha(T - \lambda_0 I)$; for otherwise $\gamma T(\lambda)$ would be continuous at λ_0 . Since all nearby values λ are also in $\sigma(T) \setminus w(T)$, the discontinuity of $\gamma T(\lambda)$ requires that $\alpha(T - \lambda I) = 0$ in $\sigma(T) \setminus w(T)$. Therefore λ_0 is an isolated point of $\sigma(T)$.

(1) \Leftrightarrow (4): The forward implication follows from [7, Theorem 1]. Conversely, suppose that $\lambda_0 \in \sigma(T) \setminus w(T)$. Then $T - \lambda_0 I$ is Weyl and $\alpha(T - \lambda_0 I) > 0$. Since every $\lambda \in \sigma(T) \setminus w(T)$ satisfies the condition

N_λ , by the punctured neighborhood theorem there exists a neighborhood $N(\lambda_0; P)$ for some $P > 0$ such that $\alpha(T - \lambda I)$ is constant (say n_0) on $N(\lambda_0; P) \setminus \{\lambda_0\}$ and $0 \leq \alpha(T - \lambda I) < \alpha(T - \lambda_0)$. We now claim that $n_0 = 0$. Assume to the contrary that $n_0 \neq 0$. Also by the punctured neighborhood theorem there exists a neighborhood $N(\lambda_0; q)$ for some $q > 0$ such that $\lambda_1 \in N(\lambda_0; q) \setminus \{\lambda_0\}$ implies $\alpha(T - \lambda_1 I) > 0$ and $T - \lambda_1 I$ is Weyl. Thus we have $\lambda_1 \in \sigma(T) \setminus w(T)$. Now by the same reason as for λ_0 ; there exists a neighborhood $N(\lambda_1; r)$ for $r > 0$ such that $\alpha(T - \mu I)$ is constant (say n_1) and $0 \leq \alpha(T - \mu I) < \alpha(T - \lambda_1 I)$. Thus $\lambda \in [N(\lambda_0; q) \cap N(\lambda_1; r)] \setminus \{\lambda_0, \lambda_1\} \Leftrightarrow \alpha(T - \lambda I) = n_1 < n_0$, a construction. Therefore $n_0 = 0$ and hence λ_0 is an isolated point of $\sigma(T)$. Hence it follows from (2) that Browder's theorem holds for T .

(1) \Leftrightarrow (5): If T obeys Browder's theorem then $\sigma(T) \setminus w(T) = \pi_{00}(T)$. Therefore $T - \lambda I$ has finite ascent for each $\lambda \in \sigma(T) \setminus w(T)$. Conversely, Suppose that $T - \lambda I$ has finite ascent for each $\lambda \in \sigma(T) \setminus w(T)$. Then by the Index Product Theorem, $\alpha((T - \lambda I)^n) - \beta(T - \lambda I)^n = i((T - \lambda I)^n) - n \cdot i(T - \lambda I) = 0$. Thus if $\alpha((T - \lambda I)^n)$ is a constant then so is $\beta((T - \lambda I)^n)$. Therefore $T - \lambda I$ is Browder. Thus T obeys Browder's theorem. \square

We can't expect that Weyl's theorem holds for operators having totally finite ascent. Consider the following example:

Let $T \in B(l_2)$ be defined by $T(x_1, x_2, x_3, \dots) = (0, x_1, \frac{1}{2}x_2, \frac{1}{3}x_3, \dots)$. Then T has totally finite ascent. But $\sigma(T) = w(T) = \{0\}$ and $\pi_0^{left}(T) = \phi$; hence Weyl's theorem does not hold for T . However, Browder's theorem performs better:

COROLLARY 2.2. *Suppose that $S \in B(H)$ has totally finite ascent and $T \in B(H)$ satisfies $T \prec S$. Then $f(T)$ obeys Browder's theorem for every $f \in H(\sigma(T))$. In particular, if S is a dominant operator and $T \prec S$ then Browder's theorem holds for $f(T)$ for every $f \in H(\sigma(T))$, where $H(\sigma(T))$ denotes the set of all analytic functions on an open neighborhood of $\sigma(T)$.*

Proof. Since $T \prec S$, then exists a quasiaffinity $K \in B(H)$ such that $KT = SK$. But S has totally finite ascent; hence for each λ there exists a natural number n_λ such that $N((S - \lambda I)^{n_\lambda}) = N((S - \lambda I)^{n_\lambda + 1})$. We claim that $N((T - \lambda I)^{n_\lambda}) = N((T - \lambda I)^{n_\lambda + 1})$. Let $x \in N((TS -$

$\lambda I)^{n\lambda}$, Then $(T - \lambda I)^{n\lambda+1}x = 0$, and so $(S - \lambda I)^{n\lambda+1}Kx = K(T - \lambda I)^{n\lambda+1}x = 0$. Therefore $Kx \in N((S - \lambda I)^{n\lambda+1}) = N((S - \lambda I)^{n\lambda})$, and so $(S - \lambda I)^{n\lambda}Kx = 0$. Since $K(T - \lambda I)^{n\lambda}x = 0$ and K is a quasiaffinity, $x \in N(T - \lambda I)^{n\lambda}$. Since T has totally finite ascent, it follows from Theorem 2.1 that $w(T) = \sigma_b(T)$. Let $f \in H(\sigma(T))$. We shall show that $w(f(T)) \subseteq f(w(T))$ for every $f \in H(\sigma(T))$ with no other restriction on T ([5, Theorem 2]), it suffices to show that $f(w(T)) \subset w(f(T))$.

Suppose that $\lambda \notin w(f(T))$. Then $f(T) - \lambda I$ is Weyl and (2.2.1):

$$f(T) - \lambda I = c(T - \alpha_1 I)(T - \alpha_2 I) \cdots (T - \alpha_n I)g(T),$$

where $c, \alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{C}$ and $g(T)$ is invertible. Since the operators right side of (2.2.1) commute, $(T - \alpha_i I)$ is Fredholm. Now we show that $i(T - \alpha_i I) \leq 0$. Observe that if $A \in B(H)$ is Fredholm of finite ascent then $i(A) \leq 0$: Indeed, either if A has finite descent then A is Browder and hence $i(A) = 0$, or if A does not have finite descent then $n \cdot i(A) - \alpha(A^n) - \beta(A^n) \rightarrow -\infty$ as $n \rightarrow \infty$, which implies that $i(A) < 0$. Therefore $\lambda \notin f(w(T))$, and hence $f(w(T)) = w(f(T))$. Hence $\sigma_b(f(T)) = f(\sigma_b(T)) = f(w(T)) = w(f(T))$, and so Browder's theorem holds for $f(T)$. If S is a dominant operator, then $N(S - \lambda I) \subseteq N(S^* - \overline{\lambda I})$ for all $\lambda \in \mathbb{C}$. Therefore S has totally finite ascent, and hence the conclusion is evident from the precious assertion. \square

COROLLARY 2.3. *Let $T \in B(H)$ be a **Gm**-operator. If T has totally finite ascent then $f(T)$ obeys Weyl's theorem for every $f \in H(\sigma(T))$.*

Proof. Since T has totally finite ascent, it follows from Theorem 2.1 that T obeys Browder's theorem. But T is a **Gm**-operator, it follows from [10, Theorem 14] that T obeys Weyl's theorem. Let $f \in H(\sigma(T))$. Then by Corollary 2.2 $f(w(T)) = w(f(T))$. Remembering ([13, Lemma]) that if T is isoloid then $f(\sigma(T) \setminus \pi_0^{left}(T)) = \sigma(f(T) \setminus \pi_0^{left}(T))$ for every $f \in H(\sigma(T))$. Hence $\sigma(f(T)) \setminus \pi_0^{left}(f(T)) = f(\sigma(T) \setminus \pi_0^{left}(T)) - f(w(T)) = w(f(T))$, which implies that Weyl's theorem holds for $f(T)$. \square

Recall that if $T \in B(H)$ and F is a closed subset of \mathbb{C} then we define a spectral subspace $\mathcal{HT}(F)$ as follows; $\mathcal{HT}(F) = \{x \in H : (T - \lambda I)f(\lambda) = x \text{ has an analytic solution } f : \mathbb{C} \setminus F \rightarrow H\}$.

THEOREM 2.4. *Let $T \in B(H)$. If $\mathcal{HT}(\{\lambda\}) = N(T - \lambda I)$ for every $\lambda \in \pi_{0f}(T)$, then T obeys Weyl's theorem.*

Proof. Let $\lambda \in \sigma(T) \setminus w(T)$. Then $\lambda \in \pi_{0f}(T)$, and so $\mathcal{H}T(\lambda) = N(T - \lambda I)$. Since $\mathcal{H}T(\{\lambda\})$ is invariant under T , T can be represented as the following 2×2 operator matrix with respect to the decomposition $\mathcal{H}T(\{\lambda\}) \oplus \mathcal{H}T(\{\lambda\})^\perp$:

$$T = \begin{pmatrix} \lambda & T_1 \\ 0 & T_2 \end{pmatrix} \text{ and } \mathcal{H}T(\{\lambda\}) \text{ is finite dimension, } T_2 - \lambda I \text{ is invertible.}$$

Therefore $\lambda \in \text{iso}\sigma(T)$, and hence $\lambda \in \pi_0^{\text{left}}(T)$. Conversely, let $\lambda \in \pi_0^{\text{left}}(T)$. Then using the spectral projection $P = \frac{1}{2\pi i} \int_{\partial D} (\lambda I - T)^{-1} d\lambda$ where D is an open disk of center λ which contains no other points of $\sigma(T)$, we can represent T as the direct sum $T = \begin{pmatrix} T_1 & 0 \\ 0 & T_2 \end{pmatrix}$, where $\sigma(T_1) = \{\lambda\}$ and $\sigma(T_2) \setminus \{\lambda\}$. Since $P(H) = \{x \in H : \lim_{n \rightarrow \infty} \|(T - \lambda I)_x^n\|^{\frac{1}{n}} = 0\} = \mathcal{H}T(\{\lambda\})$ and $\mathcal{H}T(\{\lambda\})$ is finite dimensional, $w(T) = w(T_2)$. But $(T - \lambda I)$ is invertible; hence $T - \lambda I$ Weyl. Therefore $\lambda \in \sigma(T) \setminus (T)$. \square

COROLLARY 2.5. *If $T \in B(H)$ is totally paranormal operator then Weyl's theorem holds for $f(T)$ for every $f \in H(\sigma(T))$.*

Proof. If T is totally paranormal, then it follows [11, Corollary 4.8] that $\mathcal{H}T(\{\lambda\}) = N(T - \lambda I)$ for every $\lambda \in \mathbb{C}$. Therefore by Theorem 2.4 Weyl's theorem holds for T . But T has totally finite ascent and T is an isoloid, it follows from the proof of Corollary 2.3 that $f(T)$ obeys Weyl's theorem \square

THEOREM 2.6. *Let $T \in B(H)$ be a compact operator. Then*

- (1) *Browder's theorem holds for T ;*
- (2) *Weyl's theorem holds for T if and only if Browder's theorem holds for T ;*
- (3) *The spectrum of T $\sigma(T)$ is a compact set.*

Proof. Let $B(H)/K(H)$ be a Calkin algebra and let $\pi : B(H) \rightarrow B(H)/K(H)$ be the natural map. Then by [19, Definition 2.1] $\sigma_e(T) = \sigma(\pi(T))$. And so if T is compact then $\sigma_e(T) = \sigma(\pi(T)) = \{0\}$.

(1) Since T is compact; $\sigma_e(T) = \{0\}$; $\text{acc}\sigma T = \{0\}$ and $\sigma_e(T) \subseteq w(T) \subseteq \sigma_b(T) = \sigma_e(T) \cup \text{acc}\sigma(T)$, we have $\sigma(T) = \{0\}$ and so $w(T) = \{0\}$. Thus $\text{acc}\sigma(T) \subseteq w(T)$ by [10, Theorem 9] Browder's theorem holds.

(2) Assume that Weyl's theorem holds. Then $\sigma(T) \setminus w(T) = \pi_0^{left}(T)$. But T is compact, $w(T) = \{0\} = \sigma_b(T)$. And so $\sigma(T) \setminus w(T) = \sigma(T) \setminus \sigma_b(T)$. Assume that Browder's theorem holds. Then $\sigma(T) \setminus \sigma_b(T) = \pi_{00}(T)$. Since Browder's theorem holds for T , by [10, Theorem 2] $\sigma(T) = w(T) \cup \pi_0^{left}(T)$. And so $\sigma(T) \setminus w(T) = \pi_0^{left}(T)$.

(3) Since T is a compact operator on H , $\sigma(T)$ is both closed and bounded. And so $\sigma(T)$ is a compact set. \square

DEFINITION 2.7. Let $T \in B(H)$. T is said to be a Riesz Operator if it has the following three properties:

- (1) For every $\lambda \neq 0$ and each positive integer n , the set of solutions of the equations $(\lambda I - T)^n(x) = 0$ forms a finite-dimensional subspace of X , which is independent of n provided that n is sufficiently large.
- (2) For every $\lambda \neq 0$ and each positive integer n , the range of $(\lambda I - T)^n$ is a closed subspace of X which is independent of n provided that n is sufficiently large.
- (3) The eigenvalues of T have at most one cluster point 0.

COROLLARY 2.8. Let $T \in B(H)$ be a Riesz Operator. Then

- (1) Browder's theorem holds for T ;
- (2) Weyl's theorem holds for T if and only if Browder's theorem holds for T .

Proof. By [16, Theorem 3.14] $\sigma(T)$ is countable and has no cluster point except possibly 0. And so Corollary 2.8 is easily proved from Theorem 2.6 \square

Furthermore, if the spectral radius $\nu(T)$ of $T \in B(H)$ is zero number, that is ; $\nu(T) = 0$, then by [16, Proposition 1.10] Weyl's theorem and Browder's theorem hold for T .

References

- [1] S.K. Berberian, An extension of Weyl's theorem to a class of not necessarily normal operators, Michigan Math. J. 16(1969), 273-279.
- [2] S.K. Berberian, The Weyl spectrum of an operator, Indiana Univ. Math. J. 20(1970), 529-544.

- [3] L.Chevreaun, On the spectral picture of an operator, J. Operator Theory 4(1980), 119-132.
- [4] L.A.Coburn, Weyl's theorem for nonnormal operators, Michigan Math. J. 13(1966), 285-288.
- [5] G. Gramsch and D. Lay, Spectral mapping theorems for essential spectra, Math. Ann 192(1971), 17-32.
- [6] K. Gustafson, Doubling Perturbation Size and preservation of operators indices in normed linear spaces, Cambridge Philos. Soc. 66(1969), 281-294.
- [7] K. Gustafson, Necessary and sufficient conditions for Weyl's theorem, Michigan Math. J. 19(1972), 71-81.
- [8] R.E. Harte, Fredholm, Weyl and Browder Theory, Proc. Royal Irish Acad. 85A(2)(1985), 151-176.
- [9] R.E. Harte, Invertibility and Singularity for Bounded Linear Operators, Dekker, New Youk 1988.
- [10] R.E. Harte and W.Y. Lee, Another note on Weyl's theorem, Trans. Amer. Math. Soc. 125(1997), 2115-2124.
- [11] K.B. Laursen, Opeators with finite ascent, Pacific J. Math. 152(1992), 323-336.
- [12] K.B. Laursen, Essential spectra through local spectral theory, Proc. Amer. Math. Soc. 123(1997), 1425-1434.
- [13] W.Y.Lee and S.H.Lee., A spectral mapping theorem for the Weyl spectrum, Glasgow Math. J. 38(1996), 61-64
- [14] C.Schmoeger, On isolated points of the spectrum of a bounded linear operator, Proc. Amer. Math. Soc. 117(1993), 715-719.
- [15] H.Weyl, Über beschra ñkte quadratische Former, deren Differenz Vollsteigist, Rend. Cire. Mat. Dalermo 27(1909), 373-392.
- [16] H.R. Dowson, Spectral theory of linear operators, Academic Press, London, New York, 1978, 53-53.
- [17] Apostol. C., The reduced minimum modulus, Michigan Math. J. 32(1984), 279-282.
- [18] Carl L. Devito, Fuctional Analysis and Linear Operator theory, Addison-Wesley, New York, United Kingdom, 1990, 114-126.
- [19] John B. Conway, A course in Functional Analysis, Springer-Verlag, New Youk, Berlin Heildeberg, 1985, 355-360.

Department of Mathematic Education
Kangwon University
Chuncheon 200-701, Korea
E-mail: dslee@cc.kangwon.ac.kr