

Preventing of Bone Loss in Ovariectomized Rats by *Salvia miltiorrhiza* Radix Root Extract

Sung-Hye Park

Department of Herbal Resources, Professional Graduate School of Oriental Medicine,
Wonkwang University, Iksan 570-749, Korea

Abstract

Salvia miltiorrhiza Radix root extracts (SMR) was evaluated for inhibition of the progress of bone loss induced by ovariectomy (OVX) in rats. We measured body weight and bone histomorphometry in sham, OVX and SMR-administered OVX rats. From light microscopic analyses, porous or erosive appearances were observed on the surface of trabecular bone of tibia in OVX rats, whereas those of the same bone in sham rats and in SMR-administered rats were composed of fine particles. The trabecular bone area and trabecular thickness in OVX rats were decreased by 50% from those in sham rats, and these decreases were completely inhibited by administration of SMR for 7 weeks. In this study, the mechanical strength in femur neck was significantly enhanced by the treatment of SMR for 7 weeks. In OVX rats, free T_3 was normal in all cases, whereas free T_4 was significantly increased. Although there was no difference between OVX and SMR-administered rat in T_3 levels, we have found significant difference between them in T_4 level. These results strongly suggest that SMR may be beneficial for preventing bone loss in OVX rats.

Key words: ovariectomy, *Salvia miltiorrhiza* Radix, bone cell, bone histomorphometry

INTRODUCTION

Bone mass decreases during aging, and low bone mass is the major determinant of all osteoporotic features. The dramatic increase in bone turnover rate and imbalance of bone formation and bone resorption during the first year after the cessation of ovarian function are responsible for the high rate of postmenopausal bone loss in older women. Estrogen therapy is effective in preventing postmenopausal bone loss. In elderly women, their bone loss persists as shown by high rates of bone formation and bone resorption, and estrogen replacement therapy may increase bone mass (1).

The ovariectomized (OVX) rat is a good animal model of estrogen-deficiency-induced osteopenia. Some age-related changes documented in humans, such as the appearance of porosities on the endosteal surface of bone and a loss of trabecular bone, also occur in old rats (2,3).

The annual herb, *Salvia miltiorrhiza* Radix, is widely distributed in China and Korea. *Salvia miltiorrhiza* Radix is a traditional Chinese medicine, which has been well documented for its anti-cancer effects(4). The antifibrotic effects of *Salvia miltiorrhiza* Radix extract was recently studied (5). *Salvia miltiorrhiza*-induced inhibitory effect

on cell hyperplasia induced by balloon angioplasty was also recently studied in rats (6). Therefore, it is reasonable that extracts of *Salvia miltiorrhiza* Radix extracts be investigated for antiosteoporotic activity. In the present study, we have shown that *Salvia miltiorrhiza* Radix extracts prevented the progression of bone loss induced by OVX in rats.

MATERIALS AND METHODS

Preparation of *Salvia miltiorrhiza* Radix extracts

Salvia miltiorrhiza Radix extracts were prepared by steeping the root in distilled water at 60°C for 3 hrs. The extract was filtered with 0.45 μ m filters, lyophilized and kept at 4°C. These materials were obtained from Oriental Medicine Hospital, Wonkwang University (Iksan, Korea). A voucher specimen (No. 99-02-0009) was deposited at the Herbarium at the College of Pharmacy, Wonkwang University.

Experimental animal

Thirty female Sprague-Dawley rats weighting 200~300 g were purchased from Samtako Experimental Animal Center (Osan, Korea), and the animals were maintained under constant temperature ($25 \pm 2^\circ\text{C}$) and humi-

dity ($55 \pm 5\%$), and under 12 h/12 h light-dark cycles. The rats were housed individually in standard cages and provided with a commercial standard diet containing 1.2% calcium and 0.8% phosphorus.

At 15 weeks of age, animals were either OVX or Sham-Operated (Sham). Bilateral ovariectomies were performed under anesthesia with sodium nembutal administered intraperitoneally. Sham operations were performed by exteriorizing the ovaries. OVX rats were divided into two groups based on treatment with either *Salvia miltiorrhiza* extracts or physiological saline by oral feeding. After 7 weeks of treatment, blood was collected and stored at -70°C until biochemical determinations. The rats were sacrificed on the following day. Both femora and tibia were dissected. The right femurs were cleaned from adherent muscles, placed in sterile saline, and stored for 8~16 h at 4°C . Left femurs of 12 and 8 randomly selected animals per group were fixed with Burkhart fixative and used further for histomorphometrical analysis. All procedures using animals were carried out in accordance with the guidelines presented in the Guideline Principles for the Care and Use of Animals in the Field of Physiological Sciences, published by the Physiological Society of Korea, and were approved by the National Institute of Fitness and Sports Animal Care Committee.

Biochemical analysis

Serum calcium, phosphate, and alkaline phosphatase were determined by standard laboratory techniques. Serum free T_4 (FT_4) and free T_3 (FT_3) were measured by the Liso-Phase kits (Technogenetics, Milan, Italy) after chromatographic separation of the hormone by Sephadex LH-20 chromatography.

Bone histomorphometry

The left femurs were fixed for 24 h with Bürkhardt fixative, cut sagittally into 2 equal halves with a diamond saw (Buehler Isomet, low speed saw, Chicago, IL, USA), dehydrated with methanol and embedded in methylmethacrylate. These longitudinal sections of the proximal femur were cut with an AO Autocut/Jung 1150 microtome to a $4\ \mu\text{m}$ thickness. The sections with the widest marrow cavity near the central part of the femoral neck were selected for further histological processing and histomorphometric measurements. The $4\ \mu\text{m}$ thick sections were stained according to the von Kossa method with a tetrachrome counterstain (Polysciences, Warrington, PA) for measurements of cancellous bone volume, osteoblast surface, and osteoclast surface.

All bone measurements were performed with the Bioquant Bone Morphometry System (R & M Biometrics Corp., Nashville, TN) as previously described (7). Cancellous bone measurements were performed in the proximal femur in an area beginning 1 mm distal to the growth plate-metaphyseal junction and extending further distally to the junction of the femoral neck and greater trochanter. Cancellous bone volume as a percentage of bone tissue area and osteoblast and osteoclast surfaces as percentages of total cancellous perimeter were measured at a magnification of $\times 200$. Trabecular number, width, and separation were calculated.

ous bone measurements were performed in the proximal femur in an area beginning 1 mm distal to the growth plate-metaphyseal junction and extending further distally to the junction of the femoral neck and greater trochanter. Cancellous bone volume as a percentage of bone tissue area and osteoblast and osteoclast surfaces as percentages of total cancellous perimeter were measured at a magnification of $\times 200$. Trabecular number, width, and separation were calculated.

Measurement of mechanical strength of the femur neck

The femoral necks were fractured in a hydraulic testing device, using a loading rate of 0.095 radian/sec ($5.43^{\circ}/\text{sec}$). First, the shafts were fractured 19 mm above the knee joint in three-point anterior bending. The necks were then fractured in combined bending and compression and bending, achieving highly reproducible and neck fractures, initiated fracture proximal and stop of the femoral neck and stretching downward in a straight line toward the lesser trochanter in a shearing fashion. The load in the test apparatus was measured with a load cell connected to a microcomputer via amplifier. Load/deflection curves were recorded on-line using Work Bench Mac software (Strawberry Tree, Sunnyvale, CA) (8).

Statistical analysis

All statistical analyses were performed using SAS software (SAS Institute Inc., Cary, NC, USA). Intergroup differences were analyzed by one-way analysis of variance (ANOVA), and Tukey's studentized range test was used to compare pairs of means. These parametric statistical tests could be used because the data were normally distributed. Bone histomorphometry results were analyzed by nonparametric statistics using ANOVA and Wilcoxon test.

RESULTS

Effect of *Salvia miltiorrhiza* extracts on trabecular bone area

The different histological areas of the proximal tibia sections from sham, ovariectomy (OVX), OVX+*Salvia miltiorrhiza* extracts rats are depicted. As expected, the loss of trabecular bone was observed in ovariectomized rats and was most obvious in the central metaphyseal region. The area partially filled with trabecular bone (the thick region) was observed only in the *Salvia miltiorrhiza* extracts-administered rats, which exhibited bone of a constant width. In the extract-treated rats, the trabecular bone in the area between the growth junction and the thick region was maintained, compared with that of the sham rats. Conversely, the loss of trabecular bone

Table 1. Effect of *Salvia miltiorrhiza* extracts on the trabecular bone area of OVX rats

| Groups | No. of animals | Trabecular bone area (%) |
|------------------------|----------------|---------------------------|
| Sham | 10 | 56.95 ± 8.2 ¹⁾ |
| OVX ²⁾ | 10 | 33.43 ± 5.1*** |
| OVX + SM ³⁾ | 10 | 47.16 ± 6.9 [#] |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

***p < 0.001; Statistical significance as compared with sham group.

[#]p < 0.05; Statistical significance as compared with OVX group.

in the area between the growth plate-metaphyseal junction and the thick region was remarkably pliable in the sham rats. The trabecular bone area (%) was measured (Table 1). As expected, the trabecular bone area was significantly lower in the OVX than in the sham group. However, the trabecular bone area in *Salvia miltiorrhiza* extract-administered rats was maintained at the level observed in the sham rats.

Effects of *Salvia miltiorrhiza* extracts on trabecular thickness and separation

At 7 weeks after operation, the trabecular thickness in OVX rats was significantly less than in the sham rats. The decrease in trabecular thickness induced by OVX was inhibited by the administration of *Salvia miltiorrhiza* extracts for 7 weeks (Table 2). The trabecular separation in OVX rats was also significantly inhibited by the administration of *Salvia miltiorrhiza* extracts for 7 weeks (Table 3). In addition, there was no significant change in trabecular number between OVX and *Salvia miltiorrhiza*-administered OVX rats (OVX + *Salvia miltiorrhiza*) (data not shown).

Effect of *Salvia miltiorrhiza* extracts on osteoclast number and osteoblast surface

The osteoclasts located on trabecular bone surfaces

Table 2. Effect of *Salvia miltiorrhiza* extracts on the trabecular thickness of OVX rats

| Groups | No. of animals | Trabecular thickness (µm) |
|------------------------|----------------|-----------------------------|
| Sham | 10 | 95.24 ± 7.8 ¹⁾ |
| OVX ²⁾ | 10 | 46.92 ± 5.1*** |
| OVX + SM ³⁾ | 10 | 74.34 ± 10.0 ^{###} |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

***p < 0.001; Statistical significance as compared with sham group.

^{###}p < 0.01; Statistical significance as compared with OVX group.

Table 3. Effect of *Salvia miltiorrhiza* extracts on the trabecular separation of OVX rats

| Groups | No. of animals | Trabecular separation (µm) |
|------------------------|----------------|------------------------------|
| Sham | 10 | 109.08 ± 23.40 ¹⁾ |
| OVX ²⁾ | 10 | 147.98 ± 50.2 |
| OVX + SM ³⁾ | 10 | 97.85 ± 6.1*** |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

***p < 0.001; Statistical significance as compared with OVX group.

and the mean number of osteoclasts per unit area (mm²) of trabecular bone in the three experimental groups are shown in Table 4. The number of osteoclasts per bone area was remarkably affected by OVX. However, there was no significant change in osteoclast numbers between in OVX rats and in OVX + *Salvia miltiorrhiza* rats. Although the surface of osteoblasts was significantly larger in OVX rats, the extracts-induced effect on osteoblast surface was not significant in these models.

Effect of *Salvia miltiorrhiza* extracts on the mechanical strength of the femur neck

OVX decreased the ash weight of the femur neck at each follow-up timepoint after operation, the respective mean values being 33.86 N in the sham group and 29.89 N in the OVX group at 7 weeks after OVX. The maximum loads of femoral neck also decreased due to estrogen deficiency at each follow-up timepoint after operation (data not shown). The mechanical strength in femur neck was significantly enhanced by the treatment of *Salvia miltiorrhiza* extracts for 7 weeks (Table 5).

Effect of *Salvia miltiorrhiza* extracts on serum biochemical parameters

To assess the contribution of *Salvia miltiorrhiza* extracts to factors involved in anti-osteoporotic reactions, we conducted serum biochemical analysis. At 7 weeks, the serum level of alkaline phosphatase (ALP) and in-

Table 4. Effect of *Salvia miltiorrhiza* extracts on the osteoclast numbers and osteoblast surface of OVX rats

| Groups | No. of animals | Osteoclast numbers (No/mm) | Osteoblast surface (%) |
|------------------------|----------------|----------------------------|------------------------|
| Sham | 10 | 0.58 ± 0.20 ¹⁾ | 7.20 ± 1.6 |
| OVX ²⁾ | 10 | 1.63 ± 0.50** | 21.94 ± 6.7*** |
| OVX + SM ³⁾ | 10 | 1.28 ± 0.41*** | 18.99 ± 4.7*** |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

p < 0.01, *p < 0.001; Statistical significance as compared with sham group.

Table 5. Effect of *Salvia miltiorrhiza* extracts on the mechanical strength of the femur neck of OVX rats

| Groups | No. of animals | Mechanical strength (N) |
|------------------------|----------------|----------------------------|
| Sham | 10 | 33.86 ± 4.25 ¹⁾ |
| OVX ²⁾ | 10 | 25.39 ± 5.89* |
| OVX + SM ³⁾ | 10 | 29.89 ± 3.28 [#] |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

*p < 0.05; Statistical significance as compared with sham group.

[#]p < 0.05; Statistical significance as compared with OVX group.

organic phosphorus (IP) was significantly higher in OVX rats, compared with sham rats. The increase in ALP activity was inhibited by the administration of *Salvia miltiorrhiza* extracts. However, the extract did not prevent the increase in IP in OVX rats. However, the levels of serum calcium were not affected by OVX or the administration of *Salvia miltiorrhiza* extract (Table 6). The data reported in Table 6 also shows the serum T₃ and T₄ level. Thyrotoxicosis is associated with increased bone turnover, resulting in bone loss. Mean T₄ was significantly higher in OVX rats, although the amount of T₃ was not changed in OVX rats compared to sham rats. When all available data on thyroid hormone measurements were evaluated, decreased T₄ was found in *Salvia miltiorrhiza* extract-administered OVX rats.

DISCUSSION

In traditional medicine, there are many natural crude drugs that have the potential for use in the treatment of bone diseases, however, there is little research evaluating the efficacy of most of them. In a search for natural crude drugs having inhibitory activity on bone resorption, we have screened a number of plants widely used in traditional medicine for their inhibitory activity on bone resorption induced by ovariectomy (OVX).

In this study, we evaluated the effects of *Salvia miltiorrhiza* extracts on trabecular bone volume. In the *Sal-*

via miltiorrhiza-treated rats, the trabecular bone in the area of the growth plate-metaphyseal junction was maintained, compared with that of sham rats. *Salvia miltiorrhiza*-treated rats maintained their trabecular bone in the area between the growth plate-metaphyseal junction and the thick region, which resulted in them maintaining the trabecular bone structure.

Humans and rats have many histomorphologic characteristics of the proximal femur in common (9,10). Both exhibit similar changes in bone mass and structure at numerous skeletal locations following loss of estrogen. In this study, OVX promoted cancellous bone remodeling with an increase in bone resorption that exceeded the increase in bone formation resulting in a net loss of bone. This loss of bone was also accompanied by substantial changes in trabecular structure, as indicated by immunohistochemistry. In general, data collected on the rodent model of estrogen deficiency are comparable to other reports (11-13). As shown in Table 5, changes in the maximum loads at the femoral neck revealed a deteriorating effect of OVX after 7 weeks. The loss of trabecular bone breaking force was observed in OVX rats. In *Salvia miltiorrhiza*-administered rats, the trabecular bone force was maintained, compared with that of sham rats. Next, the ovariectomized rats induced an increase in serum phosphorus level, but calcium levels did not change, as reported previously (14,15). Kalu et al. (16) have reported decreased serum calcium after OVX, however, this was detected only at a higher level of dietary calcium. There does not appear to be any evidence for a reduction in serum calcium prior to bone loss in the OVX rats. The increase in bone resorption does not, therefore, appear to occur in response to a detectable change in serum calcium. Serum albumin levels decrease rapidly after ovariectomy, possibly indicating a direct effect of estradiol on hepatic albumin production. This change appears to be due to decreased albumin synthesis rather than albumin leakage from dysfunctional liver cells (17). In addition, we have found that serum phosphate (IP) was increased in OVX rats. However, *Salvia miltiorrhiza* extracts had no regulatory

Table 6. Effect of *Salvia miltiorrhiza* extracts on serum biochemical levels of OVX rats

| Groups | No. of animals | Calcium (mg/dL) | Phosphorus (mg/dL) | Alkaline phosphatase (IU/L) | Tritodothyronine (T ₃ : ng/mL) | Thyroxine (T ₄ : g/dL) |
|------------------------|----------------|----------------------------|--------------------|------------------------------|---|-----------------------------------|
| Sham | 10 | 10.97 ± 0.97 ¹⁾ | 5.60 ± 0.88 | 122.50 ± 29.24 | 1.10 ± 0.19 | 1.02 ± 0.31 |
| OVX ²⁾ | 10 | 10.80 ± 0.70 | 7.31 ± 1.09** | 324.60 ± 43.05** | 1.16 ± 0.11 | 3.23 ± 0.56*** |
| OVX + SM ³⁾ | 10 | 10.64 ± 0.55 | 7.28 ± 1.02 | 214.14 ± 80.28 ^{##} | 1.12 ± 0.17 | 2.64 ± 0.53 [#] |

¹⁾Values are mean ± SD.

²⁾Ovariectomized.

³⁾*Salvia miltiorrhiza*.

p < 0.01, *p < 0.001; Statistical significance as compared with sham group.

[#]p < 0.05, ^{##}p < 0.01; Statistical significance as compared with OVX group.

effect on the increased IP. The OVX-induced increase in IP was related to increased renal phosphate reabsorption and may have been due to a direct effect of estrogen on specific receptors in the kidney (18). In this study, we observed no differences in the IP concentrations between OVX and *Salvia miltiorrhiza*-treated OVX rats (Table 6). Although IP is significantly related to the level of bone resorption, *Salvia miltiorrhiza*-induced regulatory effects on bone resorption is not due to changes in IP.

Thyroid hormones play an important role in bone remodeling (19), and histomorphometric studies have shown that thyroid hormones stimulate osteoblastic and osteoclastic activities in cortical and trabecular bone (20). Thyrotoxicosis is associated with increased bone turnover, and the resorption rate exceeds the formation rate, thus resulting in bone loss (8). The OVX rats had significantly increased serum thyroxine (T₄) levels, but serum T₄ levels were partially reduced in the *Salvia miltiorrhiza* extract-administered rats. Considering that an increased rate of bone turnover is characteristic of subjects with elevated T₄ levels, *Salvia miltiorrhiza* extract inhibition of T₄ levels suggests that the oriental drug has a regulatory effect on bone turnover.

The data presented here support *Salvia miltiorrhiza* extracts-induced regulation on the maintenance of the normal bone remodeling in OVX-operated rats. The results obtained in the present study provide evidence that *Salvia miltiorrhiza* extracts contribute to the prevention of bone loss, at least in OVX rats. The studies on the isolation and characterization of the active chemical constituents are in progress.

REFERENCES

- Riggs BL, Melton LJ. 1986. Involutional osteoporosis. *N Engl J Med* 314: 1676-1686.
- Ettinger B, Genant HK, Cann CE. 1987. Postmenopausal bone loss is prevented by treatment with low-dosage estrogen with calcium. *Ann Intern Med* 106: 40-45.
- Kalu DN. 1991. The ovariectomized rat model of postmenopausal bone loss. *Bone Miner* 15: 175-191.
- Chen XG, Li Y, Yan CH, Li LN, Han R. 2001. Cancer chemopreventive activities of S-3-1, a synthetic derivative of danshinone. *J Asian Nat Prod Res* 3: 63-75.
- Nan JX, Park EJ, Kang HC, Park PH, Kim JY, Sohn DH. 2001. Anti-fibrotic effects of a hot-water extract from *Salvia miltiorrhiza* roots on liver fibrosis induced by biliary obstruction in rats. *J Pharm Pharmacol* 53: 197-204.
- Zhang T, Du L, Cui P. 1998. Experimental study of hirudo and *Salvia miltiorrhiza* on smooth muscle cell hyperplasia after balloon angioplasty. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 18: 159-161.
- Wronski TJ, Yen CF, Qi H, Dann LM. 1993. Parathyroid hormone is more effective than estrogen or bisphosphonates for restoration of lost bone mass in ovariectomized rats. *Endocrinology* 132: 823-831.
- Madsen JE, Aune AK, Falch JA, Hukkanen M, Kontinen YK, Santavirta S, Nordsletten L. 1996. Neural involvement in post-traumatic osteopenia: an experimental study in the rat. *Bone* 18: 411-416.
- Bagi CM, DeLeon E, Rizzoli R, Ammann P, Miller SC. 1996. Histo-anatomical characteristics of the proximal femur in mature rats. Impact of ovariectomy. *Anat Rec* 245: 633-644.
- McGowan JA. 1993. Osteoporosis: assessment of bone loss and remodeling. *Aging* 5: 81-93.
- Ammann P, Rizzoli R, Slosman D, Bonjour JP. 1992. Sequential and precise *in vivo* measurement of bone mineral density in rats using dual-energy x-ray absorptiometry. *J Bone Miner Res* 7: 311-315.
- Ravn P, Hetl ML, Overgaard K, Christiansen C. 1994. Premenopausal and postmenopausal changes in bone mineral density of the proximal femur measured by dual-energy x-ray absorptiometry. *J Bone Miner Res* 9: 1975-1980.
- Griffin MG, Kimble R, Hopfer W, Pacifici R. 1993. Dual energy x-ray absorptiometry of the rat-accuracy, precision, and measurement of bone loss. *J Bone Miner Res* 8: 795-800.
- Morris HA, Porter SJ, Durbridge TC, Moore RJ, Need AG, Nordin BEC. 1992. Effects of ovariectomy on biochemical and bone variables in the rat. *Bone Miner* 18: 133-142.
- Hietala EL. 1993. The effect of ovariectomy on periosteal bone formation and bone resorption in adult rats. *Bone Miner* 20: 57-65.
- Kalu DN, Liu CC, Hardin RR, Hollis BW. 1989. The aged rat model of ovarian hormone deficiency bone loss. *Endocrinology* 124: 7-16.
- Nagae Y, Miyamoto M, Miyamoto H. 1991. Effect of estrogen replacement on liver function in ovariectomized rats. *J Tox Sci* 16: 87-100.
- Stock JL, Coderre JA, Burke EM, Danner DB, Chipman SD, Shapiro RJ. 1992. Identification of estrogen receptor mRNA and the estrogen modulation of parathyroid hormone-stimulated cyclic AMP accumulation in opossum kidney cells. *J Cell Physiol* 150: 517-525.
- Mosekilde L, Eriksen EF, Charles P. 1990. Effect of thyroid hormones on bone and mineral metabolism. *Endocrinol Metab Clin North Am* 19: 35-63.
- Mosekilde L, Melsen F, Bagger JP, Myhre-Jensen O, Schwartz Soresen N. 1977. Bone changes in hyperthyroidism: interrelationships between bone morphometry, thyroid function and calcium-phosphorus metabolism. *Acta Endocrinol* 85: 515-525.