

## **Two New Records of Siphonophores (Hydrozoa: Siphonophora) in Korea**

**Jung Hee Park\* and Jung Hye Won<sup>1</sup>**

(Department of Life Science, College of Natural Sciences,  
The University of Suwon, Gyeonggi-do 445-743, Korea;

<sup>1</sup>Seodaemun Museum of Natural History, Seoul 120-113, Korea)

### **ABSTRACT**

Some siphonophores were collected from the Korea Strait, 33° 59'N-129° 17'E and Yeosu with Issacs-Kidd Midwater trawl net in Sep. 2002. They were identified into *Diphyes chamissonis* Huxley, 1859 and *Abylopsis eschscholtzi* (Huxley, 1859) in the suborder Calycophorae of the order Siphonophora, respectively. They are new to the Korean fauna. As a result of this work, six siphonophores of four families have been reported in Korea.

Key words: taxonomy, Hydrozoa, Siphonophora, Korea

### **INTRODUCTION**

The siphonophore has a polymorphic organization with many types of structure, each of which shows a different function. Unlike most other cnidarians, the siphonophores do not have an alternation of generations consisting of an attached polypoid stage and a free-swimming medusa, but derivatives of both stages are attached to the stem of the free floating or swimming colonies (Kirkpatrick and Pugh, 1984).

Up to date the taxonomic studies on the siphonophores in Korea have not been actively done. Four siphonophore species were reported in Korea by Park (2002), Park and Song (2004) and

---

\*To whom correspondence should be addressed

Tel: 82-220-2480, Fax: 82-31-220-2483, E-mail: jhpark5@suwon.ac.kr

Park and Won (2004).

Some siphonophores were collected from the Korea Strait, 33° 59'N-129° 17' E and Yeosu with Issacs-Kidd Midwater trawl net in Sep. 2002. They were sorted and preserved in about 5% formalin after narcotization with menthol powders. And the observations for identification were done with stereo- and lightmicroscope. The photographs were taken with Nikon stereomicroscope SMG-U with Digital Still Camera DXM 120.

## SYSTEMATIC ACCOUNTS

Order Siphonophora

Suborder Calycothorae

Family Diphyidae

### \**Diphyes chammisonis* Huxley, 1859 (Fig. 1A-D)

*Diphyes chammisonis*: Totton and Bargmann, 1965, p. 156, figs. 93-94; Pugh, 1999, p. 489, fig. 3.70; Gao et al., 2002, p. 114, fig. 58; Namikawa and Soyama, 2004, p. 71, photo.

**Material examined.** Korea Strait (33° 59'N-129° 17'E), 24 Sep. 2002 (J. H. Won); Yeosu, Sep. 2002 (J. H. Won).

**Description.** Anterior nectophore relatively slender, blunt apically, about 11-13 mm long. Five longitudinal ridges, reaching apex. Ventral side twisted a little from apex to right. Hydroecium relatively deep, reaching up to mid height of nectosac, extending well below margin of ostium, with deep basal notch in ventral wall. Somatocyst narrow and short, nearly same wide, not reaching apex. Nectosac slender, gradually narrowing toward apex of nectophore. Ostial teeth three, pointed and equal size. Eudoxid bract roughly conical, about 6-7 mm long from apex to margin of hydroecial flap. Neck-shield large, with a club-shaped phyllocyst. Dorsal and sutural surface well rounded off, meet with dorsal wall at apex. Gonophore resembled with anterior nectophore, with serrated longitudinal ridges, mouth flap and ostial teeth. Eudoxid phase with two gonophores was examined.

**Remarks.** *D. chammisonis* resembles with *D. bojani* (see Park and Won, 2004) in the shape of anterior nectophore. But it is distinguished from later by the deeper hydroecium, the shorter somatocyst and the blunt apex.

**Distributon.** Korea, Japan, East China Sea, Great Barrier Reef, Indo Pacific, South Africa.

Family \*\*Abylidae


### \*\*\**Abylopsis eschscholtzi* (Huxley, 1859) (Fig. 2A-D)

*Agaliomoides eschscholtzi*: Huxley, 1859, p. 59; Lens and van Riemsdijk, 1908, p. 25, pl. III, figs. 29-31; Moser, 1925, p. 334, pl. 2, figs. 5-6 pl. 21, figs. 1-2.

*Abylopsis eschscholtzi*: Totton and Bargmann, 1965, p. 218, pl. 40, figs. 2, 4, 6; Gao et al., 2002, p. 173, fig. 99.

**Material examined.** Korea Strait (33° 59'N-129° 17'E), 24 Sep. 2002 (J. H. Won).


\*카미소투겹관해파리 (신칭), \*\*심해컵해파리과 (신칭), \*\*\*에시슬즈심해컵해파리 (신칭)



**Fig. 1.** *Diphyes chamissonis*. A, lateral view of anterior nectophore in polygastric phase; B, apex of nectophore; C, ostial teeth and mouth flap; D, lateral view of whole body in eudoxid phase. All scale bars = 1 mm.

**Description.** No polygastric phase examined. Eudoxid bract small, distinct pentagonal prismatic shaped in ventral and dorsal views. Upper facet about 1.8 mm long, upper right and left facets about 1.8 mm long, lower right and left facets about 2 mm long. From upper facet to lower junction of lower right and left facets about 3 mm long. All ridges of two facets serrated. Phyllocyst short.

**Remarks.** Not being examined the polygastric phase of *A. eschscholtzi*, it is easy to distinguish this species from related other species by the shape of bract of eudoxid phase. This species is similar to *A. tetragona* (see Totton and Bargmann, 1965) in the shape of eudoxid bract. But the



**Fig. 2.** *Abylopsis eschscholtzi*. A, ventral view of bract in eudoxid phase; B, left lateral view of the same; C, right lateral view of the same; D, frontal view of the same. All scale bars = 1 mm.

lateral ridges of dorsal facet toward apical facet instead of being of vertical as in *tetragona*.

**Distribution.** Korea, Japan, East China Sea, Pacific Ocean, Indian Ocean.

## REFERENCES

- Gao, S., H. Hong and S. Zhang, 2002. Fauna Sinica. Invertebrata 27. Phylum Cnidaria. Science Press, Beijing China, pp. 1-275.
- Huxley, T. H., 1859. The Oceanic Hydrozoa. Lond., pp. 1-141.
- Kirpatrick, P. A. and P. R. Puch, 1984. Siphonophores and Velellids. Linn. Soc. Lond., pp. 1-154.
- Lens, A. D. and T. van Riemsdijk, 1908. The Siphonophora of the Siboga Expedition. Siboga Exped., **9**: 1-230, pls. 1-40.
- Moser, F., 1925. Die Siphonophoren der Deutschen Sudpolar Expedition 1901-03. Dtsch. Sudpol. Exped. 18

- Zool., **9**: 1-541, pls. 1-36, figs. 1-61.
- Nimikawa, H. and I. Soyama, 2004. Jellyfish in Japanese Waters. Tokyo, Japan, pp. 1-118.
- Park, J. H., 2002. Two new records of siphonophores (Hydrozoa) and Semaestomeae (Scyphozoa) in Korea. Korean J. Syst. Zool., **18**(1): 53-58.
- Park, J. H. and J.-I. Song, 2004. Two new records of hydromedusae (Cnidaria: Hydrozoa) in Korea. Korean J. Syst. Zool., **20**(1): 31-37.
- Park, J. H. and J. H. Won, 2004. Three new records of marine hydromedusae (Cnidaria: Hydrozoa) in Korea. Korean J. Syst. Zool., **20**(2): 179-184.
- Pugh, P. R., 1999. Siphonophorae. In Boltovskoy, D., ed., South Atlantic Zooplankton. Backhuys Publisher, Leiden, pp. 467-511.
- Totton, A. K. and H. E. Bargmann, 1965. A Synopsis of the Siphonophora. Br. Mus (Nat. Hist.), Lond., pp. 1-230, pls. 1-11.

RECEIVED: 5 September 2005

ACCEPTED: 25 October 2005

## 한국산 관해파리류 2미기록종(히드라충강: 관해파리목)

박 정 희\* · 원 정 혜<sup>1</sup>

(수원대학교 자연과학대학 생명과학과; <sup>1</sup>서대문자연사박물관)

### 요 약

대한해협 (33°59'N-129°17'E) 과 여수에서 2002년 9월에 Issacs-Kidd Midwater 저인망으로 채집된 관해파리류를 동정·분류한 결과 관해파리목의 두껍관해파리아목에 속하는 카미소두겹관해파리 (*Diphyes chamissonis*) 와 예시솔츠심해겹해파리 (*Abylopsis schscholtzi*) 가 각각 한국 미기록종으로 밝혀져 제기해하여 보고한다. 본 연구의 결과 한국산 관해파리류는 4과 6종이 된다.