

Cloning of Bovine Macrophage Colony-stimulating Factor

Tae-yung Kim, Cheol-ho Kim¹, Sang-gil Lee² and Chung-boo Kang^{2,*}

Animal Health Division, Ministry of Agriculture and Forestry, Kwacheon 427-719, Korea

ABSTRACT : Macrophage colony-stimulating factor (M-CSF) is a growth factor required for growth and differentiation of mononuclear phagocyte lineage. Total and 16 poly (A) mRNA of bovine M-CSF were isolated from healthy bovine peripheral mononuclear cells stimulated by phorbol 12-myristate 13-acetate (TPA). The more compatible cultured mononuclear cells were 5×10^6 /ml for RNA isolation. TPA-activated mononuclear cells increased the level of M-CSF-mRNA more than concanavalin A (Con A) and lipopolysaccharide (LPS). The optimal analysis of reverse transcriptase-polymerase chain reaction (RT-PCR) for M-CSF as a growth factor required for bovine M-CSF was denaturation at 94°C for 1 minute, annealing at 57°C for 1 minute, extension at 72°C for 1 minute for 30 cycles. The size of cDNA of bovine M-CSF by RT-PCR was 774 base pairs. A 774 base pairs cDNA encoding bovine M-CSF was synthesized by reverse transcriptase polymerase chain reaction (RT-PCR). Ligated cDNA was transformed to competent cells and then plasmid isolation and digestion was performed. Molecular cloning and sequencing were performed for cDNA of bovine M-CSF. The size of cloned cDNA of bovine M-CSF was 774 base pairs. The homology of base sequence and amino acid sequence was 88% and 86% compared with known human M-CSF, respectively. From a high degree of sequence similarity, the obtained cDNA of bovine M-CSF is thought to be a specific gene of bovine M-CSF. (*Asian-Aust. J. Anim. Sci.* 2005, Vol 18, No. 6 : 892-897)

Key Words : M-CSF, RT-PCR, Cattle

INTRODUCTION

Macrophage colony stimulating factor (M-CSF) has long been recognized as the most strong immunomodulator (Motoyoshi et al., 1983; Stanley et al., 1983). M-CSF directly affects monocytes and macrophages to produce the granulocyte-CSF (G-CSF), granulocyte/macrophage-colony stimulating factor (GM-CSF) and also affects monocyte to enhance the tumor killing activity. Given these information, it was recognized as useful medicine source (Das and Stanley, 1982; Motoyoshi et al., 1983; Morgan and Stanley, 1984; Met-calf, 1986; Motoyoshi et al., 1986; Horiguchi et al., 1987; Horiguchi et al., 1988; Antczak and Gorman, 1989; Andreani et al., 1991).

Human M-CSF is a lipoprotein with 85 kDa of molecular weight and is known as a hemopoietic factor to activate the monocyte, neutrophil and thrombocyte. It exhibits several other effects such as reduction of blood cholesterol level, placental formation, maintenance of pregnancy, amplification of monocyte antitumor activity etc (Bartocci, 1983). Studies related to cloning, expression and clinical applications of human and mouse M-CSFs were reported (Stanley and Heard, 1977; Guilbert and Stanley,

1980; Hanamura et al., 1980; Ralph et al., 1980; Das et al., 1981; Tushinski et al., 1982; Chen et al., 1983; Stanley et al., 1983; Bartelmez and Stanley, 1985; Rettenmer et al., 1986; Horiguchi et al., 1987; Rambaldi et al., 1987; Rambaldi et al., 1988; Sariban et al., 1988) but research related to the separation, purification, cloning, expression and application of M-CSF of cattle was very few (Oshima et al., 2003; Yoshihara et al., 2003).

Application of bovine M-CSF would be rather broader than that of human M-CSF (hM-CSF). These might be included the analysis of etiology, prevention, and indicator of treatment for mastitis, infectious diseases and metabolic diseases, giving a serious damage to cattle. Hence, in the present study, we cloned bovine M-CSF.

MATERIALS AND METHODS

Isolation and purification of total RNA and mRNA from Cattle M-CSF

Peripheral blood (200 ml) of healthy cattle (Holstein, ♀) was collected and centrifuged (4°C) to isolate total RNA. Mononuclear cells were separated and collected from the buffy coat layer harvested after centrifugation.

These cells were used to determine the appropriate levels of macrophage stimulants and the cell numbers. Concanavalin A (Con A, Sigma-Aldrich Co.), phorbol 12-myristate 13-acetate (TPA, Sigma-Aldrich Co.) and lipopolysaccharide (LPS, Sigma-Aldrich Co.) were used as the stimulants.

Various levels of the mononuclear cells were grown in the medium of 10% FCS RPMI 1640 (w/v) 10% FCS at a

* Corresponding Author: C. B. Kang. Tel: +82-55-751-5814, Fax: +82-55-751-5803, E-mail: cbkang@gsnu.ac.kr

¹ Livestock Promotion Institute, Gyeongsangnamdo, Jinju 660-3601, Korea.

² College of Veterinary Medicine (The Institute of Animal Medicine), Gyeongsang National University, Jinju 660-701, Korea.

Received December 26, 2004; Accepted March 5, 2005

5% CO₂ incubator for 24 h. Only adherent cells were collected, cesium trifluoroacetate (Sigma-Aldrich Co.) treated, and finally purified by ethanol precipitation. The concentration was determined by spectrophotometric method (Manchester, 1995; Wilfinger et al., 1997). mRNA was purified from total RNA using a oligo (dT) cellulose column and then spin column chromatography (Union 3₂R plus, Hanil, Kor.).

Conditions for 5' and 3' primers synthesis

Primers were synthesized from four amino acid sequences (301, 302, 303, 304) of human M-CSF which are identified amino acid of M-CSF sequence. Simultaneously, forward primer 5' GAA CAG TTG AAA GAT CCA and GTG, and reverse primer 5' TCG GAC GCA GGC CTT GTC ATG 3' (Perkin-Elmer Co. USA) were used.

Already purified Total RNA and mRNA were used as templates.

Reverse transcriptase-polymerase chain reaction (RT-PCR)

RT-PCR condition was determined for each primer and product from each synthesized primer (base combinations) as well. Simultaneously, RT-PCR conditions for denaturation, annealing, extension and cycling were also carried out. PCR product being produced was vortexed, spin-downed to remove the mineral oil, added chloroform(99%, 100 µl) and vortexed for 5 min. DNA was collected from this mixture by the low temperature centrifugation (100 g) and stored suspended DNA (60 µl) at -20°C.

Detection of PCR products

PCR products were separated by agarose gel (2%) electrophoresis using 0.5% TBE buffer consisting of tris (hydroxymethyl) aminomethane (>99%), boric acid (>99.5%), and Na₂ EDTA·2H₂O. The products being stained with ethidium bromide (EtBr) were detected under UV. The λ/Hind III was used as a marker.

Southern hybridization

The PCR products (cDNA) marked in the under UV was denatured in a 0.4 N NaOH solution for 15 min. Finally, the marker was southern-transferred onto the Hybond N+membrane (Amersham Co. USA) for 12-16 h. The membrane transferred was neutralized in a 2×SSC solution for one min, and dried with 3 mm filter paper, and then stored, wrapped with a 3 mm filter paper until use. Before hybridization of the product, it was prehybridized in the solution of prehybridizing buffer consisting final concentration of 5×SSC solution, 5×Denhardt's solution, and 0.5% (w/v) SDS solution.

To the sealing bag containing the transferred membrane

was added hybridization buffer and electrically sealed. More than 2 h after incubation of this bag in a water bath (37°C), labelled M-CSF probe DNA was added and continuously shaken in a shaking water bath (37°C) for overnight. After hybridization, the bag was cut and substantial amount of hybridization solution was aspirated. Hybridized membrane was mixed in a solution of 2×SSC/0.1% SDS (250 ml) at 37°C for 10 min to remove excess 32p. This process was repeated until no 32p was detected by Aloka (TGS-121) survey meter (monitor). The hybridized membrane was autoradiographed.

Electrophoresis for cDNA gel extraction

Electrophoresis was performed as mentioned above using 2% agarose gel and 1×Tris-acetate (TAE) buffer except for using a bromophenol blue (BPB) as a dye, 90 V and 150 mA. cDNA band was immediately confirmed and cut under UV after electrophoresis, and put a sterile eppendorf tube (1.5 ml) for extraction of the DNA.

Gel extraction for cDNA fragments

Extraction of the cDNA contained in the gel was performed according to manufacture's instruction (Qiagen Co. Kit. USA). The cDNA was precipitated in ethanol, concentrated and stored (5 µl) at -20°C until use for cloning.

Cloning of M-CSF cDNA

Cloning of M-CSF cDNA was conducted by Cloning kit (Invitrogen Co., USA).

Before cloning, considered the insert DNA and its size, cloning vector of pGEM (3 Kb) was diluted with TE buffer. The cDNA cleaved with EcoRI was ligated into the plasmid at 12°C for overnight in the reaction mixture consisting double distilled water (4.5 µl), 10×ligation buffer 1 (1 µl), diluted vector (2 µl), insert DNA (2.5 µl), T4 DNA ligase (1 µl). Transformation was performed by transfection of the ligated reactants to competent cells in the presence of 2-mercaptoethanol (2-ME).

Transformants were grown on the LB agar plate which consisted of bactotryptone, bacto-yeast extract, NaCl and 5-bromo-4-chloro-3-indolyl-β-D-galactoside(IPTG), and only white colonies transplated onto the new LB agar plates or into sterile Asist tube and incubated in a shaking incubator (37°C). Pure white colonies were collected and plasmids were separated. The white colonies was cultured in 2 ml LB medium containing Kanamycin (50 µg/ml) with shaking.

Separation of plasmids

For Separation of plasmids, 1-1.5 ml of the medium containing transformants was used.

The plasmids separated were identified with 2% agarose gel electrophoresis.




Figure 1. The cDNA blot of bovine M-CSF by immobilization analysis.

Plasmid digestion

Plasmid (insert cDNA) digestion was performed in a 15 μ l volume containing the isolated plasmid (5 μ l), 10 \times digestion buffer (1.5 μ l), EcoR1 enzyme (10 μ g/ μ l) (0.5 μ l), and distilled water (8 μ l).

cDNA sequencing

The cDNA was incubated in a 2 N NaOH/2 mM EDTA solution for 30 min in a water bath (37°C), and denatured by 3 M NaOAc (pH 4.4) solution and absolute ethanol. After these processes, annealing labeling reaction and termination were basically followed the instruction of sequenase Kit (version 2.0, labeled dCTP, USA). After termination, Polyacrylamide gel (7%) electrophoresis was performed for 90 min with loading volume (2.5 μ l) and 1,500 volt and then sequenced by autoradiography.

RESULTS

Optimum number of mononuclear cells

The Optimum number for this experiment was found to be 5×10^6 /ml.

Effects of stimulants

The expression of M-CSF mRNA treated by TPA (40 ng/ml) was larger than the treatment of Con A (10 μ g/ml) and LPS (10 μ g/ml).

Condition for Reverse transcriptase-polymerase chain reaction (RT-PCR)

Optimum condition was found to be 30 cycles with denaturation at 94°C for 1 min, annealing at 57°C for 1 min,

and extension at 72°C for 1 min.

cDNA of PCR products

A PCR product of 774 bp was obtained from 30 cycles operated by denaturation at 94°C for one min, annealing at 57°C for one min and extension at 72°C for 1 min (Figure 1).

cDNA gel extraction

cDNA contained in the specific band obtained by RT-PCR was extracted from the gel and was exactly same as reported (Yoshihara et al., 1998).

cDNA cloning of M-CSF

The cDNA identified by 2% agarose gel electrophoresis was found to be 774 bp.

cDNA gene sequencing

Sequence of the cDNA revealed that the cDNA size was appeared to be 774 bp (Figure 2). Its homology was 88% and 86% in DNA base sequence and amino acid sequence with human M-CSF (Rambaldi et al., 1987; Rambaldi et al., 1988).

DISCUSSION

Cytokines are cellular soluble factors to induce proliferate and differentiate of cells. Variety cytokines including the M-CSF entered into cells via receptors expressed on the cell surface of target cells, and then exerts several biological activities. Several powerful functions of M-CSF were known, so that its application to cattle has important meanings. It is very hot topic to study isolation, purification and expression of human M-CSF (hM-CSF) since it is a hemopoietic factor and involved in the lipid metabolism and embryo development as well. Moreover monocytes and macrophages exhibits antitumor activities so that it is positive for M-CSF to provides antitumor mechanism in the body; thus currently, its isolation, purification and expression are being extensively studied (Ralph et al., 1980; Motoyoshi et al., 1983; Motoyoshi et al., 1986; Horiguchi et al., 1987; Rambaldi et al., 1987; Horiguchi et al., 1988).

It was proven that when monocyte of human peripheral blood was cultured with hM-CSF, it exhibited very high antitumor activity for human leukaemia cell lines K562, 16 HL60 and U937 (Horiguchi et al., 1987).

Rambaldi et al. (Rambaldi et al., 1988) reported that in the expression experiment of human M-CSF isolated 36 from peripheral blood, better expression results was obtained from incubation for 2 h with 1×10^6 cells/ml of mononuclear cell. This result was somewhat different from our result of 5×10^6 /ml, but it is not clear whether this was

			10			20			30			40			50			60		
ATG	ACC	GCG	CGG	GGC	GCC	GCC	GGG	CGC	TGC	CCT	CCC	ACG	ACA	TGG	CTG	GGC	CCT	CTG	CTG	
Met	Thr	Ala	Arg	Gly	Ala	Ala	Gly	Arg	Cys	Pro	Pro	Thr	Thr	Trp	Leu	Gly	Pro	Leu	Leu	
			70			80			90			100			110			120		
CTG	CTG	GCC	TGT	CTC	CTG	GTG	AGC	AAT	GGT	GCT	ACT	GAG	GAG	GTG	TCG	GAG	AAC	TGT	AGC	
Leu	Leu	Ala	Cys	Leu	Leu	Val	Ser	Asn	Gly	Ala	Thr	Glu	Glu	Val	Ser	Glu	Asn	Cys	Ser	
			130			140			150			160			170			180		
CAC	ATG	ATT	GGG	AAC	GGA	CAC	CTG	CTG	TTC	CTG	CAG	CAG	CTG	ATT	GAC	AGT	CAG	ATG	GAG	
His	Met	Ile	Gly	Asn	Gly	His	Leu	Leu	Phe	Leu	Gln	Gln	Leu	Ile	Asp	Ser	Gln	Met	Glu	
			190			200			210			220			230			240		
ACC	TCG	TGC	CAA	ATT	TCC	TTC	GAG	TTT	GTA	GAC	CAG	GAG	CAG	TTG	GAT	GAT	CCC	GTG	TGC	
Thr	Ser	Cys	Gln	Ile	Ser	Phe	Glu	Phe	Val	Ala	Gln	Glu	Gln	Leu	Asp	Asp	Pro	Val	Cys	
			250			260			270			280			290			300		
TAC	CTT	AAG	AAG	GCA	TTT	CTC	CTG	GTG	CAA	GAC	ATA	ATG	GAG	GAT	ACC	ATG	CGC	TTC	AAA	
Thr	Leu	Lys	Lys	Ala	Phe	Leu	Leu	Val	Gln	Asp	Ile	Met	Glu	Asp	Thr	Met	Arg	Leu	Lys	
			310			320			330			340			350			360		
GAC	AAC	ACC	CCC	AAT	GCC	AAA	GRC	ATC	GTC	CAG	CTC	CAG	GAA	CTC	TCT	CTG	AGG	CTG	AAG	
Asp	Asn	Thr	Pro	Asn	Ala	Lys	Val	Ile	Val	Gln	Leu	Gln	Glu	Leu	Ser	Leu	Arg	Leu	Lys	
			370			380			390			400			410			420		
AGC	TGC	TTC	ACC	ATG	GAC	TAT	GAT	GAG	CAG	GAC	AAG	GCT	TGT	GTC	CGA	ACA	TTC	CAT	GAG	
Ser	Cys	Phe	Thr	Met	Ala	Tyr	Asp	Glu	Gln	Glu	Lys	Ala	Cys	Val	Arg	Thr	Phe	His	Glu	
			430			440			450			460			470			480		
CGC	CCT	CTC	CAG	TTG	CTG	GAG	AAG	ATC	AAG	AAT	GTC	TTT	AAT	GAA	ACA	AAA	AAT	CTC	CTT	
Thr	Pro	Leu	Gln	Leu	Leu	Glu	Lys	Ile	Lys	Asn	Val	Phe	Asn	Glu	Thr	Lys	Asn	Leu	Leu	
			490			500			510			520			530			540		
AAA	AAG	GAC	TGG	AAC	ATT	TTC	AGC	AAG	AAC	TGC	AAC	AAC	AGC	TTT	GCT	AAA	TGC	TCC	AGC	
Lys	Lys	Asp	Cys	Asn	Ile	Phe	Ser	Lys	Asn	Cys	Asn	Asn	Ser	Phe	Ala	Lys	Cys	Ser	Ser	
			550			560			570			580			590			600		
CAT	GGC	CAC	GAG	AGG	CAG	CAC	AAG	GAA	GCC	TCC	GAT	CCC	CAG	CTC	CCT	GGT	TTT	GTC	TTC	
His	Gly	His	Glu	Arg	Gln	His	Lys	Glu	Ala	Ser	Asp	Pro	Gln	Leu	Pro	Gly	Phe	Val	Phe	
			610			620			630			640			650			660		
CGC	CTG	CTG	GTG	CCC	AGT	ATC	ATC	CTG	GTC	TTG	CTG	GCT	GTC	GGC	GGG	CTC	CTG	TTC	TAC	
Arg	Leu	Leu	Val	Pro	Ser	Ile	Ile	Leu	Val	Leu	Leu	Ala	Val	Gly	Gly	Leu	Leu	Phe	Tyr	
			670			680			690			700			710			720		
AGG	CGG	CGG	CGT	CGG	AGC	CAT	CAA	GAG	CCA	CAG	ATG	GTG	GAT	TCT	CCC	ATG	GAG	CAA	CCA	
Arg	Arg	Arg	Arg	Arg	Ser	His	Gln	Glu	Pro	Gln	Met	Val	Asp	Ser	Pro	Met	Glu	Gln	Pro	
			730			740			750			760			770			774		
GAG	GGC	AGC	CTC	CTG	ACC	CAG	GAA	GAG	GAC	AGA	CAG	GAG	GAG	CTG	CCA	GTG	TAG			
Glu	Gly	Ser	Leu	Leu	Thr	Gln	Glu	Glu	Asp	Arg	Gln	Glu	Glu	Leu	Pro	Val	***			

Figure 2. Nucleotide sequence of bovine macrophage colony-stimulating factor cDNA gene.

resulted from stimulation differences between macrophages of human and cattle. It might be a reason that Percoll 40 of isolation solution used in this study was different from Ficoll-Conray used by Rambaldi et al. (1998).

Rambaldi et al. (Rambaldi et al., 1987; Rambaldi et al., 1988) reported that Y-interferon (Y-IFN) 500 U/ml or phorbol myristate acetate (PMA) 10^{-10} Mol/L released from activated T cell and NK cell was suitable for macrophage activation. Horiguchi et al. (1987) also reported that maximum RNA was produced in human peripheral blood

M-CSF when stimulated for 6 hrs with TPA. Sariban et al. (Sariban et al., 1988) performed an experiment to stimulate human tumor necrosis factor (TNF) gene expression of monocytes derived from healthy human peripheral bloods with TPA, cycloheximide (CHX), an inhibitor of protein synthesis, and actinomycin D (ACT), an antitumor agent. They found that RNA amount of TNF stimulated by ACT (5 μ g/ml) for one h was rather decreased to 25%, and by CHX (10 μ g/ml), no RNA detected; however, by TPA (32 nM), increased to 50 fold. These results suggested that TPA is a

suitable stimulant for M-CSF and TNF as well.

The M-CSF gene size cloned from peripheral blood of cattle was 774 bp.

No report was appeared in the literature, so it very difficulties to interpret this result at this moment; however even its size is somewhat different from known human M-CSF gene size 4.0 or 4.6 Kb, they exhibited 88% homology in base sequence and 86% homology in amino acid sequence. This suggests that the 774 bp gene could be a bovine specific M-CSF gene (Ralph et al., 1980; Das et al., 1982; Kawasaki et al., 1985; Horiguchi et al., 1987). We believe that even though the 774 bp obtained from this study is not full gene size of bovine M-CSF gene, it is necessary to investigate the expression and functions of M-CSF gene, due to its several important biological functions. New techniques for genomic DNA separation and aquisition are required other than currently proceeded RT-PCR for cDNA aquisition.

Blood activity fo normal human M-CSF (hM-CSF) was turned out 174 ± 76 U/ml (about 40 bM) assayed by radioimmunoassay (RIA) (Das et al., 1981) and 5.5 ± 1.1 ng/ml assayed by enzyme-linked immunosorbent assay (ELISA) (Hanamura et al., 1988).

Simultaneously, blood activity of pregnant women was not consistent: its value was elevated by pregnancy, increased to 2-3 fold at the late stage of pregnancy, decreased by parturition, and finally returned to original value after 2-3 weeks later of parturition (Hanamura et al., 1988). In a mouse experiment, normal mouse treated with gonadotropin has an increased uterus weight as well as M-CSF activity; however in the ovariectomied mouse treated with gonadotropin no such effect was observed, facts that gonadotropin stimulates M-CSF growth in uterus via ovary (Guilbert and Stanley, 1980). Monocytes of human peripheral blood cultured with hM-CSF exhibited an increase antitumor activity against human leukemia cell line (K562, 35 HL60, U937, etc) (Horiguchi et al., 1987).

As mentioned, the hM-CSF acts not only as a hemopoietic growth factor, but is tightly related to lipid metabolism and development of fetal. Furthermore, it induced antitumor activity of monocytes and macrophages for body defence mechanism.

In animals, estrus cycle and spleen macrophage activity of rats during pregnancy. Kang and Kwak (Kang and Kwak, 1994) proved that the macrophage activity was highest ($68 \pm 6.4\%$) at proestrus and lowest ($49 \pm 4.6\%$) at metestrus, and during pregnancy, the activity was less than 5% at 1 first day, but it was slowly increased from days, rapidly increased around 12 days and continued the increase before delivery. Especially, the reason why macrophage activity was low to 5% at day 1 is might due to ovulation which is thaught to be a big injury.

The activity is being used for cure it, after which the activity is increased due to in response to pregnancy

phenomenon (Metcalf, 1986). The activity of spleen macrophage observed from pregnant rats by Kang and Kwok was quite agreed with the fact of increased hM-CSF activity in blood of pregnant during pregnancy investigated by Hanamura et al. (Hanamura et al., 1988) suggesting that M-CSF activity is closely related to macrophage activity. Hence, bovine M-CSF gene expression and its functional tests, macrophage activity could be used to markers for identify causes of mastitis, other infectious disease, metabolic disease and for prognosis of cattle (Newby and Bourne, 1977; Craven, 1986; Colditz and Mass, 1987; Duhamel, 1987; Oshima et al., 2003; Yoshihara et al., 2003).

CONCLUSION

Mononuclear cells were isolated from peripheral blood of cattle for cloning the M-CSF gene of cattle. TPA was best stimulant for mRNA production at mononuclear cell (5×10^6 /ml) among other stimulants (Con A, LPS) tested. RT-PCR with template from total RNA and mRNA of the peripheral blood produced a cDNA fragment with 774 bp long, identified by southern hybridization. After gel extraction of cDNA with a band (774 bp), M-CSF cDNA was cloned. Competent cell was trasfected by the vector ligated with cDNA obtained. After incubation, white colony was collected and digested. Cattle's M-CSF insert DNA (gene) was separated from the digest and sequenced to get 774 bp size of gene. Base sequence of M-CSF gene showed 88% homology with human gene and amino acid sequence was 86% homology with human indicating that this gene might be a specific M-CSF gene of cattle. However, further systemic researches should be performed for the gene expression and the functions of its product.

REFERENCES

- Andreani, C. L., D. W. Payne, J. N. Packman, C. E. Resnick, A. Hurwitz and E. Y. Adashi. 1991. Cytokine-mediated regulation of ovarian function. Tumor necrosis factor α inhibits gonadotropin-supported ovarian androgen. *J Biol Chem.* 266(11):6761-6766.
- Antczak, D. F. and N. T. Gorman. 1989. Cellular interaction in immune responses. In: (Ed. N. T. Gorman and R. E. Halliwell). *Veterinary Clinical Immunology* W.B. Saunders Co. Philadelphia. 107-123.
- Bartelmez, S. H. and E. R. Stanley. 1985. Synergism between hemopoietic growth factors (HCFs) detected by their effects on cells bearing receptors for a lineage specific HGF: Assay of hemopoietin-1. *J. Cell Physiol.* 122:370-378.
- Bartocci, A., J. W. Pollard and E. R. Stanley. 1983. Regulation of colony-stimulating factor during pregnancy. *J. Exp. Med.* 164:207-212.
- Chen, B. D. M., H. S. Lin and S. Hus. 1983. Tumor-promoting phorbol esters inhibits the binding of colony-stimulating factor (CSF-1) to murine peritoneal exudate macrophages. *J. Cell*

- Physiol. 116:207-212.
- Colditz, L. G. and P. J. C. M. Mass. 1987. The inflammatory activity of activated complement in ovine and bovine mammary glands. *Immunol. Cell Biol.* 65:433-436.
- Craven, N. 1986. Chemotactic factor for bovine neutrophils in relation to mastitis. *Comp. Immunol. Microbiol. Infect Dis.* 9:29-36.
- Das, S. K. and E. R. Stanley. 1981. Human colony-stimulating factor radioimmunoassay: Resolution of three subclasses of human colony-stimulating factors. *Blood.* 58(3):630-641.
- Das, S. K. and E. R. Stanley. 1982. Structure-function of a colony stimulating factor (CSF-1). *J. Biol. Chem.* 257(22):13679-13684.
- Duhamel, G. E. 1987. Distribution of T and B lymphocytes in mammary dry secretions. Colostrum and blood of adult dairy cattle. *Vet. Immunol. Immunopathol.* 14:101-122.
- Guilbert, L. J. and E. R. Stanley. 1980. Specific interactions of murine colony-stimulating factor with mononuclear phagocytic cell. *J. Cell Biol.* 85:153-159.
- Hanamura, T., K. Motoyoshi, K. Yoshida, M. Saito, Y. Miura, T. Kawashima, M. Nishida and F. Takaku. 1988. Quantitation and identification of human monocytic colony-stimulating factor in human serum by enzyme-linked immunosorbent assay. *Blood* 72(3):886-892.
- Horiguchi, J., M. K. Warren and D. Kufe. 1987. Expression of the macrophage-specific colony-stimulating factor in human monocytes treated with granulocyte-macrophage colony-stimulating factor. *Blood* 69:1259-1261.
- Horiguchi, J., M. L. Sherman, A. Sampson-Johannes and D. Kufe. 1988. CSF-1 and C-form gene expression in human carcinoma cell lines. *Biochem. Biophys. Res. Commun.* 157(1):395-401.
- Kang, C. B. and S. D. Kwak. 1994. Activity and application of 20 α -hydroxysteroid dehydrogenase in rat. II. Changes in activities of the splenic macrophages and ovarian histological findings. *Kor. J. Vet. Res.* 34(4):873-880.
- Kawasaki, E. S., M. B. Iadner, A. M. Wang, V. Arsdell, M. K. Warren, M. Y. Coyne, V. L. Schweichart, M. T. Lee, K. J. Wilson, A. Boosman, E. R. Stanley, P. Palph and D. F. Mark. 1985. Molecular cloning of a complementary DNA encoding human macrophage-specific colony stimulating factor (CSF-1). *Science (Wash DC).* 230:291-296.
- Manchester, K. L. 1995. Value of A_{260/280} ratio for measurement of purity of nucleic acid. *Biotechniques.* 19:208-219.
- Metcalf, D. 1986. The molecular biology and functions of the granulocyte-molecular biology and function of the granulocyte-macrophage colony-stimulating factors. *Blood* 67:257-263.
- Morgan, C. J. and E. R. Stanley. 1984. Chemical cross-linking of the mononuclear phagocyte specific growth factor CSF-1 to its receptor at the cell surface. *Biochem. Biophys. Res. Commun.* 119:350-41.
- Motoyoshi, K., F. Takaku and Y. Miura. 1983. High serum colony-stimulating activity of leukocytopenic patients after intravenous infusions of human urinary colony-stimulating factor. *Blood* 62:685-691.
- Motoyoshi, K., F. Takaku, T. Maekawa, Y. Miura, K. Kimura, S. Furasawa, M. Hattori, T. Nomura, H. Mizoguchi, M. Ogawa, K. Kinugasa, T. Tomimaga, M. Shimoyama, K. Deura, K. Ohta, T. Taguchi, T. Masaoka and I. Kimura. 1986. Protective effect of partially purified human urinary colony-stimulating factor on granulocytopenia after antitumor chemotherapy. *Exp. Hematol.* 14:1069-1075.
- Newby, T. J. and J. Bourne. 1977. The nature of the local immune system of the bovine mammary gland. *J. Immunol.* 118:461-465.
- Oshima, K., H. Watanabe, K. Yoshihara, T. Kojima, O. Dochi, N. Takenouchi, M. Fukushima and M. Komatsu. 2003. Gene expression of leukemia inhibitory factor (LIF) and macrophage colony stimulating factor (M-CSF) in bovine endometrium during early pregnancy. *Theriogenology* 15:60(7):1217-26.
- Ralph, P., M. K. Warren, M. T. Lee, L. J. Csejtey, J. F. Weaver, H. E. Broxmeyer, D. E. Williams, E. R. Stanley and E. S. Kawasaki. 1980. Inducible production of human macrophage growth factor, CSF-1. *Blood* 68:633-639.
- Rambaldi, A., C. Y. Diane and J. D. Griffin. 1987. Expression of the M-CSF (CSF-1) gene by human monocytes. *Blood* 69:1409-1413.
- Rambaldi, A., N. Wakamiya, E. Vellenga, J. Horiguchi, Warren M-Kim, D. Kufe and J. D. Griffin. 1988. Expression of the macrophage colony-stimulating factor and c-fms genes in human acute myeloblastic leukemia cells. *J. Clin. Invest.* 81:1030-1035.
- Rettenmier, C. W., R. Sacca, W. L. Furman, M. F. Roussel, J. T. Holf, A. W. Nienhuis, E. R. Stanley and C. J. Sherr. 1986. Expression of the human c-fms proto-oncogene product (colony-stimulating factor-1 receptor) on peripheral blood mononuclear cells and choriocarcinoma cell lines. *J. Clin. Invest.* 77:1740-1746.
- Sariban, E., T. Mitchell and D. Kufe. 1985. Expression of the c-fms proto-oncogene during human monocytic differentiation. *Nature (Lond).* 316:64-66.
- Sariban, E., K. Imamura, L. Robert and D. Kufe. 1988. Transcriptional and post transcriptional regulation of tumor necrosis factor gene expression in human monocytes. *J. Clin. Invest.* 81:1506-1510.
- Stanley, E. R. and P. M. Heard. 1977. Factors regulating macrophage production and growth. *J. Biol. Chem.* 252(12):4305-4312.
- Stanley, E. R., L. J. Guilbert, R. J. Tushinski and S. H. Bartelmez. 1983. CSF-1 A mononuclear phagocytic lineage-specific hemopoietic growth factor. *J. Cell Biochem.* 21:151-159.
- Tushinski, R. J., I. T. Oliver, L. J. Guilbert, P. W. Tynan, J. R. Warner and E. R. Stanley. 1982. Survival of mononuclear phagocytes depends on lineage-specific growth factor that the differentiated cells selectively destroy. *Cell* 28:71-81.
- Wilfinger, W. W. and R. Cortini. 1997. Effect of pH and ionic strength on the spectrophotometric assessment of nucleic acid purity. *Biotechniques* 22:474-481.
- Yoshihara, K., S. Inumaru, Y. Hirota and E. Momotani. 1998. Cloning and sequencing of cDNA encoding bovine macrophage colony-stimulating factor (bM-CSF) and expression of recombinant bM-CSF using baculovirus. *Vet. Immunol Immunopathol.* 12:63(4):381-91.
- Yoshihara, K., K. Oshima, Y. Muneta, R. Kikuma, C. Yayota, T. Hirai, N. Satoh, S. Matsuura, Y. Kikyo, M. Satoh, C. Kubota, S. Inumaru, Y. Yokomizo and Y. Mori. 2003. Quantitation of bovine macrophage colony-stimulating factor in bovine serum by ELISA. *Vet. Immunol. Immunopathol.* 15:95(3-4):103-111.