

## Genetic Variability and Relationships of Native Japanese Chickens Assessed by Microsatellite DNA Profiling - Focusing on the Breeds Established in Kochi Prefecture, Japan -

S. A.-M. Osman, M. Sekino<sup>1</sup>, M. Nishibori, Y. Yamamoto and M. Tsudzuki\*

Laboratory of Animal Breeding and Genetics, Graduate School of Biosphere Science, Hiroshima University  
Higashi-Hiroshima, Hiroshima 739-8528, Japan

**ABSTRACT** : Blood samples were collected from eight native Japanese breeds of chickens (Miyadi-dori, Ohiki, Onaga-dori, Shoukoku, Tosa-Jidori, Tosa-Kukin, Toutenkou and Uzurae) and two foreign breeds of chickens (White Leghorn and Rhode Island Red) to examine the genetic variability and relationships among the breeds by using a microsatellite DNA technique. Except for the Shoukoku breed, the other Japanese chicken breeds all originate from Kochi Prefecture. Ohiki, Onaga-dori, Tosa-Jidori, Toutenkou and Uzurae are fancy fowl, and Miyadi-dori and Tosa-Kukin are utility fowl. Among the fancy fowl, Ohiki, Onaga-dori, and Toutenkou males have thick and long feathers in the saddle and tail. Genetic variabilities of the 20 microsatellites examined, varied depending on the breed: the mean number of alleles per locus ranged from 2.05 (Miyadi-dori) to 3.90 (Rhode Island Red); proportion of polymorphic loci ranged from 0.75 (Miyadi-dori) to 1.00 (Rhode Island Red, Shoukoku and Uzurae); and mean expected heterozygosity ranged from 0.330 (Miyadi-dori) to 0.607 (Rhode Island Red). Unique microsatellite alleles were detected in each breed. Using the neighbour-joining method, phylogenetic trees were constructed based on the genetic distances of  $D_J$  and  $D_{ST}$ . Among the breeds originating from Kochi Prefecture, fancy and utility breeds belonged to different clusters. Among the fancy breeds, those having thick and long feathers in the tail and saddle showed a close genetic relationship to the Shoukoku breed, which also has thick and long feathers in the tail and saddle. (*Asian-Aust. J. Anim. Sci.* 2005, Vol 18, No. 6 : 755-761)

**Key Words** : Genetic Relationship, Genetic Variability, Japanese Native Japanese Chicken, Microsatellite, Kochi Prefecture

### INTRODUCTION

In Japan, there are approximately 50 breeds of native chickens (Tsudzuki, 2003). Most of them were bred by the end of the Edo Era (1603 to 1867) for their special plumage, crowing ability, and cockfighting ability. Others were bred to produce eggs and/or meat through the Meiji Era (1868 to 1912) to the Taishou Era (1912 to 1926).

Among the Japanese breeds, seven originated from Kochi Prefecture: they are Tosa-Jidori, Onaga-dori (Tosano-Onaga-dori), Ohiki (Minohiki Chabo), Toutenkou, Uzurae (Uzura Chabo), Tosa-Kukin, and Miyadi-dori. This is the largest number of breeds that were established in any one prefecture. The former five are fancy (ornamental) breeds that were established by the end of the Edo Era, and the latter two are utility breeds developed during and after the Meiji Era (Oana, 1951; Tsudzuki, 2003). Kochi Prefecture is located on the Pacific Ocean side of Japan's Shikoku Island (<http://www.jetro.go.jp/investjapan/>).

The earliest study of genetic relationships among native Japanese chickens was performed based on external characteristics and old literature (Oana, 1951). Later,

osteometrical and somatometrical studies were carried out (Nishida et al., 1985 a,b). In addition, several phylogenetic and variability studies based on blood groups and/or blood protein polymorphisms have been performed (Hashiguchi et al., 1981; Okabayashi et al., 1998; Okada et al., 1980, 1984, 1989; Tanabe and Mizutani, 1980; Tanabe et al., 1991). However, these investigations detected only a limited number of polymorphic loci and alleles per locus. In addition, the genetic relationships of native Japanese chicken breeds reported in these studies were not always in accordance with those from the morphological and literature studies. As a result, the genetic relationships among native Japanese chicken breeds remain unclear.

Recently, microsatellites have become the preferred type of genetic markers for several reasons, including their abundance, random distribution, codominant inheritance, high variability, and possibility of automated detection. Additionally, microsatellites have the ability to show genetic polymorphisms even in species in which classical genetic markers have been proven to be unsuccessful (Goldstein and Pollock, 1997; Petit et al., 1997). Accordingly, microsatellite markers have proven useful in assessing genetic variation and diversity in livestock (Buchanan et al., 1994; MacHugh et al., 1994; Martinez et al., 2000; Pandey et al., 2002).

A large number of chicken microsatellite markers have been determined (Groenen et al., 2000) and have been

\* Corresponding Author: M. Tsudzuki. Tel: +81-82-424-7950, Fax: +81-82-424-7950, E-mail: [tsudzuki@hiroshima-u.ac.jp](mailto:tsudzuki@hiroshima-u.ac.jp)

<sup>1</sup> Tohoku National Fisheries Research Institute, Shinjima, Shiogama, Miyagi 985-0001, Japan.

Received July 12, 2004; Accepted December 27, 2004

**Table 1.** List of chicken breeds used in this study

Breeds	Breed abbreviations	No. of individuals examined	Source of samples
Miyadi-dori	MYA	24	Kochi Prefecture (fancier)
Ohiki	OHK	28	Kochi Prefecture (fanciers)
Onaga-dori	ONA	27	Kochi Prefecture (fanciers)
Rhode Island Red	RIR	27	HS-NLBC <sup>1</sup>
Shoukoku	SHO	24	Mie Prefecture (fancier)
Tosa-Jidori	TJI	24	KPLES <sup>2</sup>
Tosa-Kukin	TKU	26	Kochi Prefecture (fanciers)
Toutenkou	TOT	37	Kochi Prefecture (fancier)
Uzurao	UZU	23	Kochi Prefecture (fanciers)
White Leghorn	WL	24	OS-NLBC <sup>3</sup>

<sup>1</sup> Hyogo Station, National Livestock Breeding Center.

<sup>2</sup> Kochi Prefectural Livestock Experiment Station.

<sup>3</sup> Okazaki Station, National Livestock Breeding Center.

employed to evaluate the genetic diversity of chicken populations (Groen et al., 1994; Crooijmans et al., 1996a; Ponsuksili et al., 1996; Vanhala et al., 1998; van Marle-Köster and Nel, 2000; Emara et al., 2002). Recently, two studies revealed the genetic relationships of native Japanese chickens based on microsatellite DNA polymorphisms (Takahashi et al., 1998; Osman et al., 2004). However, these studies did not focus on the breeds originating from Kochi Prefecture, which are the main breeds in Japan.

This paper describes the genetic variability and relationships of native chickens originating from Kochi Prefecture, Japan. The results will be useful in order to support decisions regarding conservation and further use of the breeds.

## MATERIALS AND METHODS

### Chicken breeds

Table 1 shows the breeds and sampling sites of blood samples used in this study. We assessed eight native Japanese breeds [Miyadi-dori (MYA), Ohiki (OHK), Onaga-dori (ONA), Shoukoku (SHO), Tosa-Jidori (TJI), Tosa-Kukin (TKU), Toutenkou (TOT), and Uzurao (UZU)] and two foreign breeds [White Leghorn (WL) and Rhode Island Red (RIR)]. Details of breed features are described elsewhere (Mitsui, 1979; Hawksworth, 1982; Kuroda et al., 1987; Tsudzuki, 2003). The following is a brief description of the native Japanese breeds screened in this study. TJI has a small body size (600 to 675 g), and its ancestor is thought to have been introduced to Japan more than 2,000 years ago. The overall appearance of this breed is similar to that of the Red Jungle Fowl. UZU is similar to TJI in its body size, but this breed shows a sloping back and has no tail feathers. ONA is an especially distinguished-looking breed. Some of the tail feathers and all of the saddle hackles in the males show no molting during their lives, continue to extend, and sometimes exceed 10 m in length. OHK has a small body size, but is somewhat larger than TJI. In spite of this small

body size, the tail feathers and saddle hackles of males are considerably thick and long. TOT is characterized by prolonged crowing (15 sec in average). Excepting the plumage color, TOT has great similarity to SHO in the body shape. SHO is one of the oldest breeds in Japan. It is believed to have an ancestor of Chinese origin. Tail feathers and saddle hackles of males are thick and long. TKU is a utility breed having a large (3,300 to 4,500 g), Cochun-type body. MYA is also a utility fowl with black plumage and short legs the color of lead. This breed has a single comb and white ear lobes. The whole body shape is similar to that of the Creeper in Germany (van Wulfften Palthe, 1992).

### DNA extraction and microsatellite amplification

The blood samples were collected from the ulnar vein. Genomic DNA was extracted from the whole blood by using a DNeasy Tissue Kit (QIAGEN, Tokyo). Prior to the PCR amplification, DNA concentration of individual samples was adjusted to 20 ng/ $\mu$ l.

A number of chicken microsatellite markers are available at the Web Site of the Chicken Genome Project (<http://poultry.mph.msu.edu/>). In this study, we used 20 microsatellite primer pairs, taking the genome coverage of the loci into consideration. Details of the microsatellite markers used are described in Table 2. Polymerase chain reactions (PCR) were carried out following the protocols suggested by Cheng and Crittenden (1994) and Crooijmans et al. (1996b), with some modifications. The protocol is as follows. PCR amplifications were carried out in a 20  $\mu$ l reaction mixture containing 20 ng DNA, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 0.001% gelatin, 100  $\mu$ M dNTPs, 0.5U of AmpliTaq Gold (Applied Biosystems, Foster City, USA) and 10 pM of each primer. Each reaction was performed according to the following conditions: first there were an initial denaturation at 95°C for 9 min and 43 cycles of denaturation at 95°C for 1 min. Next, there was an annealing for 1 min at an appropriate temperature (50 to

**Table 2.** Details of 20 microsatellites screened in this study

Loci <sup>1</sup>	Chro. No.	Repeat motif (5'-3')	Forward primer (5'-3')	Reverse primer (5'-3')	Annealing (°C)
ADL 262	23	(TG) <sub>9</sub>	GTGCAGACACAGAGGGAAAG	TCACATGCACACAGAGATGC	55
LEI 092	6	(TG) <sub>9</sub>	GATCTACATTTGTGCAGTGT	TCCTTGGTCTGACTCTCCATG	50
LEI 096	2	(TG) <sub>9</sub>	GATCAGATTGGTTCCCTGTG	TGGGTGAAGTTTCCCTCGTAG	55
LEI 099	12	(TG) <sub>9</sub>	GATCTGGCAGAACAGAAACAG	ATATTTCACACCTGACCTGCG	55
LEI 135	28	(TG) <sub>9</sub>	CACAATGAAGGATGAATAGTGC	AATTCACAGTTACACCTGAGG	55
LEI 209	1	(AC) <sub>11</sub> -AT(AC) <sub>4</sub> (GC) <sub>4</sub>	AATTTGGTGTGCATACCTCTCC	GACTTTCAGTGTCTCGTTTAG	55
MCW 067	10	(TG) <sub>9</sub>	GCACTACTGTGTGCTGCAGTTT	GAGATGTAGTGCCACATTCCGAC	55
MCW 145	1	(TG) <sub>10</sub>	ACTTT ATTCT CCAA TTTGG CT	AAACA CAATG GCAAC GGAAA C	55
MCW 183	7	(TG) <sub>10</sub>	ATCCCAGTGTCCGAGTATCCGA	TGAGATTTACTGGAGCCTGCC	55
MCW 193	5	(TG) <sub>10</sub>	TATTC AATAGAGTTACGCTGTC	ATTACGTCTGCACCAGTACAG	50
MCW 214	5	(TG) <sub>10</sub>	CAACAGTAACCATAACATCTGC	TACCTGGATTCTTTCATCAGG	50
MCW 217	18	(TG) <sub>10</sub>	GATCTTTCTGGAACAGATTTT	CTGCACTTGGTTCAGGTTCTG	55
MCW 222	3	(TG) <sub>10</sub>	GCAGTTACATTGAAATGATTCC	TTCTCAAAAACACCTAGAAGAC	55
MCW 233	27	(TG) <sub>10</sub>	TCCAGCAGTAAGTATAGCTGC	TGTTAGCTGCAGGGTATTAGC	55
MCW 240	4	(TG) <sub>10</sub>	CAAAACCGGTGTCACCTACTG	GGTTATTTCTTCAGTGACTTCC	50
MCW 252	3	(TG) <sub>10</sub>	CTGCTCAAGCCCATCAAATGG	CGATAACATCTGACACTGCC	55
MCW 295	4	(TG) <sub>10</sub>	ATCACTACAGAACCCTCTC	TATGTATGCACGCAGATATCC	55
MCW 301	24	(TG) <sub>10</sub>	GGAGAGGAGACAACCTGTATTC	AGGGTGAGAGGTAACAAGTGC	55
MCW 322	13	(TG) <sub>10</sub>	GATCTCCCTAGCTACAAACC	CTCCGCTTCTTGAGAGTC	55
MCW 330	17	(AC) <sub>11</sub> -AT(AC) <sub>4</sub> (GC) <sub>4</sub>	TGGACCTCATCAGTCTGACAG	AATGTTCTCATAGAGTTCTCGC	55

<sup>1</sup>The Chicken Genome Project Web Site (<http://poultry.mph.msu.edu/>) provides PCR primer information to amplify each locus.

55°C) for each marker. This was followed by an extension at 72°C for 1 min, using a Perkin-Elmer Cetus 9600 thermalcycler (Applied Biosystems). Finally, there was a final extension at 72°C for 10 min.

The PCR fragments were electrophoresed in a 5% polyacrylamide gel using an automated ABI 377 DNA sequencer with the internal size standard Genescan 350-TAMRA (Applied Biosystems). Data were collected with the ABI PRISM 377 (version 2.5) software (Applied Biosystems). Microsatellites were sized and genotyped using the GENESCAN (version 3.1) and GENOTYPER (version 2.5) software programs (Applied Biosystems).

### Statistical analysis

Allele frequency of each locus and breed was directly calculated from the observed genotypes. Genetic variability of each breed was evaluated by calculating the mean number of alleles per locus (*MNA*), proportion of polymorphic loci ( $P_{poly}$ ; Lewontin and Hubby, 1966), and unbiased expected heterozygosity ( $H_e$ ; Nei, 1978).

Two genetic distances ( $D_A$ ; Nei et al., 1983 and  $D_{ST}$ ; Nei, 1972) were calculated between all pairs of breeds, and two phylogenetic trees were constructed based on the neighbor-joining (NJ) method (Saitou and Nei, 1987). The reliability of the tree obtained was examined by a bootstrap test with 1,000 replicate re-sampling of loci with replacement. These procedures were conducted by using the POPTREE program (<http://www.bio.psu.edu/People/Faculty/Nei/softw are.htm>), and the tree was visualized on the TREEVIEW program (<http://taxonomy.zoology.gla.ac.uk/rod>).

## RESULTS

### Genetic variability

Genetic variabilities estimated for each breed are summarized in Table 3. A total of 155 alleles were observed at the 20 loci in 264 individuals from 10 chicken breeds. The number of alleles per locus across breeds ranged from 3 (ADL 262) to 15 (LEI 209) with the *MNA* of 7.75. The *MNA* in each breed ranged from 2.05 (MYA) to 3.90 (RIR). The lowest value of the  $P_{poly}$  (0.75) was found in the MYA breed, while three breeds (RIR, SHO and UZU) showed polymorphisms in all the 20 microsatellites (i.e.  $P_{poly} = 1.00$ ). The  $H_e$  ranged from 0.330 (MYA) to 0.607 (RIR). In other words, the MYA had the lowest genetic variability and the RIR showed the highest variability.

All breeds had two or more private alleles (Table 4). There were seven private alleles in WL, five in ONA and SHO, four in TKU and RIR, three in UZU, and two in each of MYA, OHK, TJI and TOT.

### Genetic distances

Matrices of genetic distances ( $D_A$  and  $D_{ST}$ ) between every pair of breeds were calculated from the allele frequencies at the 20 loci and are presented in Table 5. Two dendrograms drawn from the genetic distance matrices are shown in Figure 1A and B.

$D_A$  distance: The average genetic distance among all breeds was  $0.485 \pm 0.095$ . The lowest distance (0.275) was observed between the ONA and SHO breeds, and the highest distance (0.645) between the UZU and WL breeds. The  $D_A$  values measured between the WL and Japanese

**Table 3.** Statistics of microsatellite variability in terms of the number of different alleles at each locus, the mean number of alleles per locus ( $MNA$ ), proportion of polymorphic loci ( $P_{poly}$ ), and mean expected heterozygosity ( $He$ ), estimated for 10 chicken breeds

Loci	Breeds <sup>2</sup>										
	All <sup>1</sup>	MYA	OHK	ONA	RIR	SHO	TJI	TKU	TOT	UZU	WL
ADL 262	3	1	3	1	3	2	2	2	2	3	3
LEI 092	12	3	4	7	3	2	1	3	2	4	3
LEI 096	7	2	4	3	4	2	3	3	2	5	6
LEI 099	7	1	1	2	4	3	1	3	1	3	4
LEI 135	8	4	3	5	4	3	3	4	3	5	1
LEI 209	15	2	3	6	7	6	4	6	2	5	5
MCW 067	4	1	3	3	3	3	2	1	3	4	3
MCW 145	7	1	3	3	5	4	2	2	1	3	3
MCW 183	9	1	3	2	4	2	2	6	2	4	3
MCW 193	9	3	4	2	2	3	3	2	5	2	5
MCW 214	11	3	2	5	2	4	3	4	3	5	4
MCW 217	7	2	3	3	5	5	3	4	2	3	1
MCW 222	5	2	3	3	3	2	2	2	2	2	2
MCW 233	5	2	5	3	4	3	3	3	3	4	1
MCW 240	11	2	5	6	5	5	3	4	4	5	2
MCW 252	8	2	5	4	4	4	3	2	4	3	3
MCW 295	7	2	2	2	4	4	4	3	2	3	3
MCW 301	11	2	3	4	5	4	3	4	2	4	3
MCW 322	3	3	3	3	3	3	3	3	2	2	2
MCW 330	6	2	2	2	4	2	4	2	3	4	1
$MNA$	7.75	2.05	3.20	3.45	3.90	3.30	2.70	3.15	2.50	3.65	2.90
$SE$ of $MNA$		0.19	0.24	0.36	0.26	0.26	0.19	0.29	0.22	0.23	0.32
$P_{poly}$		0.75	0.95	0.95	1.00	1.00	0.90	0.95	0.90	1.00	0.80
$He$		0.330	0.486	0.403	0.607	0.461	0.362	0.473	0.351	0.566	0.459
$SE$ of $He$		0.055	0.045	0.045	0.027	0.045	0.049	0.047	0.052	0.031	0.056

<sup>1</sup> Number of different alleles per locus across all breeds.<sup>2</sup> See Table 1 for the abbreviation.**Table 4.** Microsatellite alleles specific to each breed

Breeds <sup>1</sup>	No. of alleles	Locus name and allele sizes (bp)
MYA	2	LEI 135: (140), MCW 217: (147)
OHK	2	MCW 183: (298), MCW 301: (285)
ONA	5	LEI 092: (265, 273, 281), LEI 209: (166), MCW 240: (175)
RIR	4	LEI 099: (123), MCW 145: (204), MCW 183: (306), MCW 240: (197)
SHO	5	LEI 099: (117), LEI 092: (243), MCW 217: (163), MCW 295: (93), MCW 301: (267)
TJI	2	MCW 193: (311), MCW 330: (258)
TKU	4	MCW 183: (296), MCW 222: (224), MCW 240: (187), MCW 295: (99)
TOT	2	LEI 135: (152), LEI 209: (138)
UZU	3	LEI 099: (135), MCW 301: (305), MCW 330: (266)
WL	7	LEI 096: (242), LEI 209: (164, 172), MCW 183: (310), MCW 214: (277, 289), MCW 252: (297)

<sup>1</sup> See Table 1 for the abbreviation.

breeds were generally high (0.531 to 0.645), and constantly greater than the average value (0.485) of 45 breed pairs. While the  $D_A$  values measured between RIR and Japanese breeds ranged from 0.393 to 0.538, the mean value (0.466) is less than the average value (0.485) of 45 breed pairs.

**$D_{ST}$  distance :** The average genetic distance among all breeds was  $0.847 \pm 0.248$ . The lowest distance (0.403) was observed between the ONA and SHO breeds, and the highest distance (1.413) between the UZU and WL breeds. The  $D_{ST}$  values determined between the WL and Japanese breeds were generally high (1.024 to 1.413), and constantly greater than the average value (0.847) of 45 breed pairs.

While the  $D_{ST}$  values determined between RIR and Japanese breeds ranged from 0.602 to 1.009, the mean value (0.801) is less than the average value (0.847) of 45 breed pairs.

#### NJ tree


Figure 1 A illustrates the genetic relationships among breeds as the NJ tree was reconstructed based on the  $D_A$  distance matrix. Chicken breeds were divided into three major groups. The first group consisted of TOT, ONA, OHK, SHO and UZU; the second group comprised WL, RIR, TKU, and MYA; and the third group included only one breed, TJI. Figure 1B illustrates the genetic

**Table 5.** Matrix of  $D_J$  (below diagonal) and  $D_{NJ}$  (above diagonal) genetic distances estimated between 10 chicken breeds based on 20 microsatellites<sup>1</sup>

	MYA	OHK	ONA	RIR	SHO	TJI	TKU	TOT	UZU	WL
MYA	0	0.959	0.717	0.610	0.835	0.916	1.025	0.781	0.708	1.024
OHK	0.594	0	0.476	0.977	0.413	0.672	1.080	0.505	0.549	1.257
ONA	0.463	0.319	0	0.809	0.403	0.868	1.062	0.607	0.714	1.362
RIR	0.425	0.518	0.468	0	0.851	0.766	0.602	1.009	0.846	0.739
SHO	0.474	0.300	0.275	0.473	0	0.733	0.839	0.585	0.520	1.063
TJI	0.542	0.447	0.491	0.418	0.423	0	1.026	1.050	0.740	1.234
TKU	0.580	0.594	0.571	0.393	0.498	0.566	0	1.094	0.841	1.056
TOT	0.519	0.379	0.356	0.538	0.411	0.547	0.582	0	0.613	1.152
UZU	0.516	0.352	0.407	0.442	0.353	0.445	0.504	0.474	0	1.413
WL	0.547	0.603	0.638	0.418	0.560	0.623	0.531	0.614	0.645	0

<sup>1</sup> Calculation of genetic distances  $D_J$  and  $D_{NJ}$  was performed based on Nei et al. (1983) and Nei (1972), respectively.

See Table 1 for the abbreviation of breed names.



**Figure 1.** Neighbor-joining trees showing the genetic relationships among eight breeds of native Japanese chickens and two foreign breeds (White Leghorn and Rhode Island Red) using  $D_J$  (A) and  $D_{NJ}$  (B) genetic distances calculated from 20 microsatellites. The numbers at the node indicate bootstrap values in percentage (1,000 replicates). MYA: Miyadi-dori, OHK: Ohiki, ONA: Onaga-dori, SHO: Shoukoku, TJI: Tosa-Jidori, TKU: Tosa-Kukin, TOT: Toutenkou, UZU: Uzuraou, RIR: Rhode Island Red, WL: White Leghorn.

relationships between breeds as the NJ tree was reconstructed based on the  $D_{NJ}$  distance matrix. Breeds were clustered into three major groups. The first group consisted of ONA, SHO, OHK, TOT, UZU and TJI; the second group was composed of WL, RIR and TKU; and the third group included only one breed, MYA.

## DISCUSSION

### Genetic variability

Genotyping at various microsatellite loci across the genome indicated that the chicken breeds examined were genetically different from each other. The microsatellite allelic composition and frequencies differed among the 10

breeds. While several alleles were shared among the 10 breeds, some alleles were observed to be breed-specific. Recently in Japan, native Japanese chickens are being mated to foreign breeds to produce better tasting, high-quality meat (Japan Chicken Association, 2003). Breed-specific alleles may be a diagnostic marker to trace the origin of meat, when ordinary chicken meat has been attempted to be passed off as high-quality meat.

In our study, the number of alleles per locus across all breeds examined was three to 15 (Table 3). This result is similar to those of Crooijmans et al. (1993, 1996a), Cheng et al. (1995), and Ponsuksili et al. (1996). As well, the *MNA* in each breed varied from 2.05 to 3.90. The Kochi breeds varied from 2.05 to 3.65. This result, also, is similar to that of van Marle-Köster and Nel (2000), in which the *MNA* varied from 2.3 to 4.3 in five chicken lines from South Africa (Koekoek, New Hampshire Red, Naked-Neck, Lebowa-Venda, and Ovambo) and two chicken populations each from Mozambique and Botswana. Furthermore, Groen et al. (1994) and Emara et al. (2002) also observed similar values of *MNA* in their studies.

In the present study, the  $H_e$  value ranged from 0.330 to 0.607 with the values of 0.330 to 0.566 in the Kochi breeds. This result is generally similar to that of van Marle-Köster and Nel (2000) mentioned above, in which the  $H_e$  values ranged from 0.314 to 0.612. Also, using microsatellite markers, Vanhala et al. (1998) reported similar  $H_e$  values ranging from 0.290 to 0.670 in eight chicken lines (three White Leghorn hybrids, three Finish Landrace lines, a Rhode Island Red line, and a broiler hybrid line). Judging from the values of *MNA* and  $H_e$ , the genetic variability displayed by the Kochi Prefecture chicken breeds is generally similar to that of chickens reared elsewhere in the world.

The  $P_{poly}$  and  $H_e$  values of native Japanese breeds in the present study were 0.75 to 1.00 and 0.330 to 0.566, respectively (Table 3). These values are higher than those reported by Hashiguchi et al. (1981) and Okada et al. (1989). These discrepancies are thought to result from the

differences of genetic markers used. We used microsatellite DNA polymorphisms, while both Hashiguchi et al. (1981) and Okada et al. (1989) reported on the  $P_{poly}$  and  $H_e$  based on blood groups and/or protein polymorphisms.

### Genetic relationships

The present study demonstrated for the first time the relationships between chicken breeds originating from Kochi Prefecture using microsatellite DNA polymorphisms. According to the NJ trees (Figure 1A and B), the utility breeds (MYA and TKU) and the fancy breeds (OHK, ONA, TJI, TOT, and UZU) originating from Kochi Prefecture, belonged to different clusters. This result is thought to accurately reflect their breeding history. The breeding of the MYA started late in the Meiji Era (1868 to 1912) based on the mating of a Black Minorca male and a Kamochi-dori female (Kamimoto, 1951). The Kamochi-dori seems to have been a small chicken with a bearded face and feathered short shanks, native to Kochi Prefecture (Sawada, 1978). However, this chicken has become extinct, and details are unknown; for example, it is unknown whether it was a pure breed or a hybrid. The TKU breed was established during and after the Meiji Era by mating some Japanese chicken with the Cochin breed, which has Chinese origin and was brought to Japan via England (Mitsui, 1979). Whereas, OHK, ONA, TJI, TOT, and UZU had already been in Kochi Prefecture by the end of the Edo Era (1603 to 1867) many years earlier (Oana, 1951). It is noteworthy that among the Kochi fancy breeds, those having thick and long tail feathers and saddle hackles (OHK, ONA, and TOT) showed a close relationship to the SHO breed, which is one of the oldest Japanese breeds and also has thick and long feathers in the tail and saddle.

There are few reports that dealt with genetic relationships between the chicken breeds originating from Kochi Prefecture. Okada et al. (1984) and Okabayashi et al. (1998) reported on the basis of blood groups and/or blood protein polymorphisms that ONA is genetically close to TOT and apparently far from SHO. This contrasts with our result in which ONA showed a closer genetic distance to SHO than to TOT (Table 5). Oana (1951) purported, based on morphology and old literature, that ONA directly descended from SHO. Although our study revealed that ONA is genetically closer to SHO than to TOT, there is no evidence from the present study to show SHO is a direct ancestor of ONA. The present study only showed that the genetic distance between ONA and SHO is slightly smaller than that between ONA and TOT.

According to Oana (1951), UZU was thought to be directly derived from TJI by a single mutation, because the two breeds are very similar in body shape and size, except for tail morphology. UZU looks like a type of TJI, lacking a rump. In the NJ tree, based on  $D_A$  (Figure 1A), TJI was

located far from UZU, differing from the hypothesis of Oana (1951). However, in the NJ tree based on  $D_{ST}$  (Figure 1B), TJI and UZU showed a close relationship, supporting the hypothesis. Further studies are necessary to reveal the genetic relationship between the TJI and UZU breeds.

### ACKNOWLEDGMENTS

This study was supported in part by a grant-in-aid from Nankoku City, Kochi Prefecture, Japan, and that from the Integrated Research Program for Functionality and Safety of Food Toward the Establishment of a Healthy Diet. The Ministry of Agriculture, Forestry and Fisheries of Japan. We are grateful to the members of Kochi Prefectural Livestock Experimental Station, and Okazaki and Hyogo Stations, National Livestock Breeding Center, for their kind help in collecting blood samples. We are also grateful to fanciers M. Tajima, T. Ikemoto, Y. Sawada, H. Nakahira, Y. Hamada, K. Nishiguchi, T. Matsuoka, and M. Komatsu (Kochi Prefecture) and S. Nishikawa and T. Nakanishi (Mie Prefecture) for their kind help in the collection of blood samples. We wish to thank Dr. H. H. Cheng, USDA, for his kind supply of primer sets of microsatellites.

### REFERENCES

- Buchanan, F. C., L. J. Adams, R. P. Littlejohn, J. F. Maddox and A. M. Crawford. 1994. Determination of evolutionary relationships among sheep breeds using microsatellites. *Genomics* 22:397-403.
- Cheng, H. H. and L. B. Crittenden. 1994. Microsatellite markers for genetic mapping in the chicken. *Poult. Sci.* 73:539-546.
- Cheng, H. H., I. Levin, R. L. Vallejo, H. Khatib, J. B. Dodgson, L. B. Crittenden and J. Hillel. 1995. Development of a genetic map of the chicken with markers of high utility. *Poult. Sci.* 74:1855-1874.
- Crooijmans, R. P. M. A., A. B. F. Groen, A. J. A. van Kampen, Van der S. Beek, J. J. van Der Poel and M. A. M. Groenen. 1996a. Microsatellite polymorphism in commercial broiler and layer lines estimated using pooled blood samples. *Poult. Sci.* 75:904-909.
- Crooijmans, R. P. M. A., A. J. A. van Kampen, J. J. van der Poel and M. A. M. Groenen. 1993. Highly polymorphic microsatellite markers in poultry. *Anim. Genet.* 24: 441-443.
- Crooijmans, R. P. M. A., P. A. M. van Oers, J. A. Strijk, J. J. van der Poel and M. A. M. Groenen. 1996b. Preliminary linkage map of the chicken (*Gallus domesticus*) genome based on microsatellite markers: 77 New Markers Mapped. *Poult. Sci.* 75:746-752.
- Emara, M. G., H. Kim, J. J. Zhu, R. R. Lapierre, N. Lakshmanan and H. S. Lillehoj. 2002. Genetic diversity at the major histocompatibility complex (B) and microsatellite loci in three commercial broiler pure lines. *Poult. Sci.* 81:1609-1617.
- Goldstein, D. B. and D. D. Pollock. 1997. Launching microsatellites: A review of mutation processes and methods of

- phylogenetic inference. *J. Hered.* 88:335-342.
- Groen, A. F., R. P. M. A. Crooijmans, A. J. A. van Kampen, S. van der Beek, J. J. van der Poel and M. A. M. Groenen. 1994. Microsatellite polymorphism in commercial broilers and layers. In *Proceedings of the 5th World Congress on Genetics Applied to Livestock Production*, Guelph, Ontario, pp. 94-97.
- Groenen, M. A. M., H. H. Cheng, N. Bumstead, B. F. Benkel, W. E. Briles, T. Burke, D. W. Burt, L. B. Crittenden, J. Dodgson, J. Hillel, S. Lamont, A. P. de Leon, M. Soller, H. Takahashi and A. Vignal. 2000. A consensus linkage map of the chicken genome. *Genome Res.* 10:137-47.
- Hashiguchi, T., M. Tsuneyoshi, T. Nishida, H. Higashiawatoko and E. Hiraoka. 1981. Phylogenetic relationships determined by the blood protein types of fowls (in Japanese with English summary). *Jpn. J. Zootech. Sci.* 52:713-729.
- Hawksworth, D. 1982. *British Poultry Standards*. 4th edn. Butterworth Scientific, London.
- Japan Chicken Association. 2003. *The Guidebook of Special Meat-Type Chickens* (in Japanese). Zenkoku-Shokuchou-Shinbun Co. Ltd., Tokyo.
- Kamimoto, H. 1951. The tale of Miyadi-dori (in Japanese). *Niwatori-no-Kenkyu* 26:38-40.
- Kuroda, N., T. Yamaguchi, Y. Tanabe and K. Ichinoe. 1987. Japanese Chickens as Natural Monuments (in Japanese). Kyoikusya, Tokyo.
- Lewontin, R. C. and J. L. Hubby. 1966. A molecular approach to the study of genetic heterozygosity in natural populations of *Drosophila pseudoobscura*. *Genetics* 54:595-609.
- MacHugh, D. E., R. T. Loftus, D. G. Barley, P. M. Sharp and P. Cunningham. 1994. Microsatellite DNA variation within and among European cattle breeds. *Proc. Royal Soc. Lond. Series B.* 256:25-31.
- Martinez, A. M., J. V. Delgado, A. Rodero and J. L. Vega-pla. 2000. Genetic structure of the Iberian pig breed using microsatellite. *Anim. Genet.* 31:295-301.
- Mitsui, T. 1979. *General View of Japanese Native Chickens* (in Japanese). Petlife-Sha, Tokyo.
- Nei, M. 1972. Genetic distance between populations. *Am. Natul.* 106:283-291.
- Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics.* 89:583-590.
- Nei, M., F. Tajima and Y. Tateno. 1983. Accuracy of estimated phylogenetic trees from molecular data: 2. Gene frequency data. *J. Mol. Evol.* 19:153-170.
- Nishida, T., Y. Hayashi, T. Fujioka, I. Tsugiyama and K. Mochizuki. 1985a. Osteometric studies on the phylogenetic relationships of Japanese native fowls. *Jpn. J. Vet. Sci.* 47:25-37.
- Nishida, T., Y. Hayashi and T. Hashiguchi. 1985b. Somatometrical studies on the morphological relationships of Japanese native fowls. *Jpn. J. Zootech. Sci.* 56:645-657.
- Oana, H. 1951. *The Histories of Japanese Native Chickens* (in Japanese). Nihonkei-Kenkyusha, Tokyo.
- Okabayashi, H., S. Kamiya and Y. Tanabe. 1998. Phylogenetic relationships among Japanese native chicken breeds based on blood protein polymorphism (in Japanese with English summary). *Jpn. Poult. Sci.* 35:173-181.
- Okada, I., K. Toyokawa and I. Takayasu. 1980. Genetic relationships of some native chicken breeds in the northern Tohoku district of Japan (in Japanese with English summary). *Jpn. Poult. Sci.* 17:337-343.
- Okada, I., Y. Yamamoto, T. Hashiguchi and S. Ito. 1984. Phylogenetic studies on the Japanese native breeds of chickens. *Jpn. Poult. Sci.* 21:318-329.
- Okada, I., Y. Yamamoto, A. Shinjo, S. Kimura and H. Hiraoka. 1989. Studies of genetic differentiation within breeds of Japanese indigenous chickens. *Jpn. Poult. Sci.* 26:207-215.
- Osman, A.-M. S., M. Sekino, M. Nishibori, Y. Kawamoto, K. Kinoshita, Y. Yamamoto and M. Tsudzuki. 2004. Genetic variability and relationships of Japanese native chickens assessed by means of microsatellite profiling approach - Focusing on the Oh-Shamo (Japanese Large Game) and its related breeds. *J. Poult. Sci.* 41:94-109.
- Pandey, A. K., M. S. Tandia, D. Kumar, B. Misra, P. Choudhary and R. K. Vijh. 2002. Microsatellite analysis of three poultry breeds of India. *Asian-Aust. J. Anim. Sci.* 15:1536-1542.
- Petit, E., S. Aulagnier, D. Vaiman, C. Bouissou and B. Crouauroy. 1997. Microsatellite variation in a introduced mouflon population. *J. Hered.* 88:517-520.
- Ponsuksili, S., K. Wimmers and P. Horst. 1996. Genetic variability in chickens using polymorphic microsatellite markers. *Thai J. Agr. Sci.* 29:571-580.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406-425.
- Sawada, Y. 1978. My grandfather, I, and Miyadi-dori (in Japanese). *Nippon-no-Aigankei* 55:8-9.
- Takahashi, H., Y. Nirasawa, M. Nagamine, M. Tsudzuki and Y. Yamamoto. 1998. Genetic relationships among Japanese native breeds of chicken based on microsatellite DNA polymorphisms. *J. Hered.* 89:543-546.
- Tanabe, Y. and M. Mizutani. 1980. Studies on the phylogenetic relationships of the Japanese native fowl breeds: 3. Genetic distances among the 16 breeds based on 16 loci, and their dendrogram (in Japanese with English summary). *Jpn. Poult. Sci.* 17:116-121.
- Tanabe, Y., T. Iida, H. Yoshino, A. Shinjo and S. Muramatsu. 1991. Studies on the phylogenetic relationships of the Japanese native fowl breeds: 5. The comparisons among native fowl breeds in Japan and its adjacent areas and European and American fowl breeds (in Japanese with English summary). *Jpn. Poult. Sci.* 28:266-277.
- Tsudzuki, M. 2003. Japanese native chickens. In *The Relationship between Indigenous Animals and Humans in APEC Region*. (Ed. H. L. Chang and Y. C. Huang). The Chinese Society of Animal Science, Taiwan.
- Vanhala, T., M. Tuiskula-Haavisto, K. Elo, J. Vilkki and A. A. Mäki-Tanil. 1998. Evaluation of genetic variability and genetic distances between eight chicken lines using microsatellite markers. *Poult. Sci.* 77:783-790.
- van Marle-Köster, E. and L. H. Nel. 2000. Genetic characterization of native southern African chicken populations: Evaluation and selection of polymorphic microsatellite markers. *South Afric. J. Anim. Sci.* 30:1-6.
- van Wulfften Palthe, A. W. 1992. C. S. Th. van Gink's Poultry Paintings. Dutch Branch of the World's Poultry Science Association, Beekbergen, the Netherlands.