

Morphological Correlation and Chronology of Lower Terrace Formations of Southeastern Coast of Korea

Kim Ju-Yong^{1*} and Choi Weon-Hak²

Corresponding author

¹*Geological and Environmental Hazard Research Division, Korea Institute of Geoscience and Mineral Resources*

²*Site Assessment Group, Environment and Structure Lab.*

Korea Electric Power Research Institute (KEPRI)

1. Introduction

Terrace stratigraphy of the southeastern coastal areas of Korea has been studied with the help of terrace mapping and chronology. Generally coastal terraces are regarded to be formed in response to coastal base level changes. Coastal terraces are conspicuously distributed along the southeastern coastal area (Kim, 1973; Oh, 1977; Jo, 1976, 1980; Oh, 1981; Lee, 1985, 1987; Kim, 1990; Kim et al, 1990; Lee and Kim, 1992a, 1992b; Choi, 2001; Kim, 2001; Kim, et al, 2004a, 2004b). In this study coastal terraces are divided into LT (above 8m~25m), MT (36m~55m), HT(63m~86m), and uHT(90m~130)m.

This study aims to suggesting the distribution of coastal lower terraces with the help of key tephra production, particularly the ages of the LT2 and LT3 Formations (Kim et al, 2004a). This research covers Lower Terrace distributed from Bonggil site to Jeongja site in SE coast of Korea.

2. Coastal terrace mapping

The longitudinal and transversal distributions of coastal terrace formations are important not only for characterizing coastal landform, but also to unravel the uplift history of coastal landscape. Coastal terrace is characterized by spherical and round beach pebbles distributed

at different altitudes along the coastline (Kim et al, 1990).

Coastal terraces can be divided into 5 main surfaces, including beach and coastal alluvial plain(AP, 4~5m), Low Terrace(LT, 8~25m), Middle Terrace(MT, 36~55m), High Terrace(HT, 63~86m) and upper High Terrace(uHT, 90~130)m. Successive steps of coastal terraces can be mapped by altitude, landscape dissection feature (back edge and fore edge) and continuity of terrace surfaces or formations distributed along or vertical to present shoreline. The chronology of coastal terrace formations can be illuminated by the carbon radiometric age, OSL and tephra production, as well as their morphostratigraphic correlations.

3. Terrace chronology

Age dating of terrace sediments and terrace formation correlation to marine isotope stages are well known methods for terrace chronology (Muhs, 2000; Martinson, et al, 1987). Coastal terrace chronology in Korea has been reviewed by many Korean researchers, even though they are in some controversy with morphostratigraphical correlation (Cheong, 2002). Holocene alluvial plain is a lowermost base level in terrestrial part and distributed at about 4m to 5m in altitude. This Holocene Coastal Alluvial Plain (AP) is assumed to be formed initially during the Holocene Climatic Optimum (HCO) or Transgression-I (Flandrian transgression, ca 6,000-7,000yrsBP) (Fig.1).

Below 25m in altitude, Lower Terrace are distributed at the altitude of about 8m to 25m (Kim et al, 2004a). In particular Lower Terrace 3 is distributed as low as 8m, and shows a sharp gradient change in fore edges in many sites. Lower terrace 2 is distributed in altitude between 14m and 18m, and Lower Terrace 1 at the level about 25m. Some horizons of sedimentary deposits of Lower Terrace 2 and 3 Formations have been numerically dated by means of OSL methods, and also examined by their tephra contents (volcanic glass). The OSL age of Lower Terrace 2 Formation, distributed at the level of about 15-18m in Yonghanri site, indicates 111-112ka, which in turn hints the formation ages of basal beach pebbles, or of wave-cut platforms below OSL-dated sand deposits. A number of tephra evidences also support the age of Lower Terrace-II Formation. It is generally accepted that the age of Aso-4 is MIS 5c(ca 90ky), that of the Ata (100-120ky) is MIS 5d or 5e, and that of much younger AT tephra (ca 25ka) is marking the boundary MIS 2 and MIS 3. The Aso-4 tephra and the Ata tephra were found in Lower Terrace 2 of Yonghan site(Inoue et al., 2004). The former tephra (Aso-4) was also found near the Wolsong Nuclear Power Plant site, while the latter tephra (Ata) was found at many sites such as Wolsong gate (Wolsong), Yonghanri(Pohang) and Shinchangri (Janggi), Jeongja (Gangdong) (Kim et al, 2004a; unpublished data of the KEPRI). Each of these tephra was identified by the proper refraction index and chemical composition of glass and pyroxene. The amount of glass in those tephra is very small, and the tephra is regarded as having been experienced some reworking in general. Based on both OSL dating and the tephra evidences, the formation age of LT 2 are correlated to MIS 5d, or MIS 5e (Kim et al, 2004a; Shackleton, et al, 1973).

The age of LT 3 can also be explained by tephra identification. First of all AT tephra was found in the upper part of both LT2 and LT 3 Formations, not in the form of any stratigraphical meanings, but produced as scattered fragments, particularly in late Pleistocene paleosol layer, or as reworked materials in pedo-sedi-

mentary layers. LT 3 is commonly superjacent by slope deposits, or old sand and gravel with some intercalations of muds as old as 30 to 40 ka, the top of which is frequently covered by patterned grounds formed under cold (boreal) climate regime during the last glacial maximum period (Kim et al, 2004b). Although a number of OSL dating have been reported and been in controversy with morpho-stratigraphy for last several years, Hataya's recent OSL result in Yonghan site of the northern part of Pohang City is remarkably promising in matching morphostratigraphical meaning with numerical datings of the LT 3 formation situated at the level of about 10m to 12m. Hataya had attempted to relate terrace stratigraphy to numerical dating, and shows OSL ages of LT 3 formation range 58ka to 63ka. The formation age of basal beach pebbles, or wave-cut platforms of LT 3 is interpreted most probably as MIS 5a (Kim, et al, 2004a).

4. Conclusion

Terrace morphology is so conspicuous in the south eastern coastal areas. Lower Terrace Formation above 8m in altitude were formed during the Late Pleistocene. Tephra deposited just above the beach pebble layers of LT2 formation and OSL dating of LT 2 and LT 3 Formations support the age of the LT 2 formation is MIS 5d or 5e, most probably 5e. The age of LT 3 is interpreted MIS 5a, based on tephra production and OSL dating of sand deposits just above the beach pebbles in the LT 3 in Yonghan area. Particularly the LT 2 Formation was formed in association with second major transgression, possibly equivalent to the well-known Monastirian episode in the Mediterranean Sea

References

- Cheong, C.S., 2002. Dating of marine terraces. *Jour. Geol. Soc. Korea*, v. 38(2), p. 279-291.
- Choi, S.G., 2001, Tectonic Movement indicated by the Pleistocene Palaeo-shoreline in the Eastern Coast of Korea. *Transactions, Japanese Geomorphologist*

- Union, v.22(3), 265-275.
- Jo,W.R., 1976. Geomorphic development of the Pohang coastal plain. *Annals. Tohoku Geographical Associations*, v.30, p.152-160.
- Jo, W.R., 1980. Holocene sea-level changes on the east coast of Korean peninsula. *Geogr. Rev. of Japan*, v. 53-5, p.317-328.
- Inoue, D., Choi, W.H., Yanagida, M., Sasaki., T., 2004, Pleistocene uplift rate of eastern Korean Peninsula using Japanese key tephra - Joint Research between Japan (CRIEPI) and Korea(KEPRI), 32nd IGC – Florence 2004, Scientific session abstracts 198-25.
- Kim, J.Y., 1990. Quaternary stratigraphy of the terrace gravel sequences in the Pohang area (Korea). Ph.D. thesis, Seoul National University. p. 1-203
- Kim, J.Y.,Choi, D.K. and Lee, D.Y., 1990. Paleoenvironmental implications of the Quaternary gravel sequences on the basis of gravel shape. *The Korean Jour. Quaternary Res.* v. 4, p. 41-57.
- Kim, J.Y., 2001. Quaternary geology and assessment of aggregate resources of Korea for the national industrial resources exploration and development.*Quaternary International*, v. 82, p. 87-100.
- Kim, J.Y., Chang, S.B., Choi, W.H, Jang, H., Choi, S,G., 2004a. Geohazard potential and paleoseismology based on coastal geomorphology and structural development. *Field guidebook in 2004-Special Lecture Course of Geohazard and Paleoseismology* (edited by McCalpin, et al), p. 52-85.
- Kim, J.Y., Lee, G.G., Yang D.Y,Hong S.S., Nahm,W.H, Lee, J.Y., 2004b. Research on the distribution and Formation Process of Quaternary Deposits of South Korea. *the Korean Pleacolithic Soc.*, v. 10, p. 11-22.
- Kim, S.W., 1973, A study of the terraces along the south-eastern coast (Bangeojin-Pohang) of the Korean peninsula.*Jour. Geol. Soc. Korea*, 9, 89-121.
- Lee, D.Y., 1985. Quaternary Deposits in the Coastal Fringe of the Korean Peninsula. Dissertation to the Vrije Universiteit, Brussel, p. 1-315.
- Lee,D.Y., 1987. Stratigraphic research of the Quaternary deposits in the Korean Peninsula: Progress in Quaternary geology of east and southeast Asia. *CCOP/TP 18*, p. 227-242.
- Lee, D.Y.and Kim, J. Y., 1992(a). Quaternary Terrace Deposits and Their Stratigraphy along the East Coast of the Korean Peninsula, *The Geological Society of Korea. field guide book*
- Lee,D.Y.and Kim, J. Y., 1992(b). Review on the Quaternary Stratigraphy of the Korean Peninsula, *Sino-Korean Prehistory Workshop*, p. 69-99.
- Martinson, D.G., Pisias, N.G., Hays, J.D., Imbrie, J., Moore Jr., T.C. and Shackleton, N.J., 1987. Age dating and orbital theory of the ice ages : Development of a high-resolution 0 to 300,000-year chronostratigraphy. *Quart. Res.*, v.27, 1-29.
- Muhs, D.R., 2000. Dating marine terraces with relative-age and correlated-age methods. In : Noller, J.S., Sower, J.M., and Lettis, W.R. (eds.) *Quaternary : Methods and applications*, AGU Reference Shelf 4, American Geophysical Union, 434-446.
- Oh, G.H., 1977. The geomorphic history of the south-eastern coast of the Korean peninsula.*Geogr. Rev. of Japan*, v.50-12, p.689-699.
- Oh G. H., 1981. Marine terrace and their tectonic deformation on the coast of southern part of the Korean peninsula. *Bull. Department of Geography, Univ. Tokyo, Japan*, v.13, 11-61.