

Managing Within-Field Spatial Yield Variation of Rice by Site-Specific Prescription of Panicle Nitrogen Fertilizer

Nguyen Tuan Ahn*, Jin Chul Shin**, and Byun-Woo Lee*†

*Department of Plant Science, Seoul National University, Seoul 151-921, Korea

**National Institute of Crop Science, R D A, Suwon 441-857, Korea

ABSTRACT: Rice yield and protein content have been shown to be highly variable across paddy fields. In order to characterize this spatial variability of rice within a field, two-year experiments were conducted in 2002 and 2003 in a large-scale rice field of 6,600 m². In year 2004, an experiment was conducted to know if variable rate treatment (VRT) of N fertilizer, that was prescribed for site-specific management at panicle initiation stage, could reduce spatial variation in yield and protein content of rice while increasing yield compared to conventional uniform N topdressing (UN, 33 kg N/ha at PIS) method. VRT nitrogen prescription for each grid was calculated based on the nitrogen (N) uptake (from panicle initiation to harvest) required for target rice protein content of 6.8%, natural soil N supply, and recovery of top-dressed N fertilizer. The required N uptake for target rice protein content was calculated from the equations to predict rice yield and protein content from plant growth parameters at panicle initiation stage (PIS) and N uptake from PIS to harvest. This model equations were developed from the data obtained from the previous two-year experiments. The plant growth parameters for the calculation of the required N were predicted non-destructively by canopy reflectance measurement. Soil N supply for each grid was obtained from the experiment of year 2003, and N recovery was assumed to be 60% according to the previous reports. The prescribed VRT N ranged from 0 to 110 kg N/ha with an average of 57 kg/ha that was higher than 33 kg/ha of UN. The results showed that VRT application successfully worked not only to reduce spatial variability of rice yield and protein content but also to increase rough rice yield by 960 kg/ha. The coefficient of variation (CV) for rice yield and protein content was reduced significantly to 8.1% and 7.1% in VRT from 14.6% and 13.0% in UN, respectively. And also the average protein content of milled rice in VRT showed very similar value of target protein content of 6.8%. In conclusion the procedure used in this paper was believed to be reliable and promising method for reducing within-field spatial variability of rice yield and protein content. However, inexpensive, reliable, and fast estimation methods of natural N supply and plant growth and nutrition status should be prepared before this method could be practically used for site-specific crop management in large-scale rice field.

Keywords: spatial variation, site-specific, variable rate, prescription, nitrogen, yield, rice, precision farming

Significant spatial variability of plant growth, and grain yield and quality has been visualized anywhere although uniform management is practiced. Management of this spatial variability is a fundamental objective of site-specific management (SSM) with expectations of improving profits while reducing environmental pollution caused by agricultural production. Significant crop yield variability and their causal factors in upland fields have been reported by numerous researchers (Paz, 2000; Beckett *et al.*, 1971; McBratney *et al.*, 1997; Miller *et al.*, 1995) and this information enabled site-specific crop management to be justified (Machado *et al.*, 2000; Bahman & Varvel, 1997; Cahn *et al.*, 1994; Jin & Jiang, 2002). In lowland rice, within-field spatial variation in crop growth and yield in relation to soil properties has been reported by Dobermann (1994), Dobermann *et al.* (1995), Nguyen *et al.* (2004), and Yanai *et al.* (2000). Nevertheless, fertilizer has been uniformly applied in rice field without regard to the spatial variability of soil properties. This uniform management of fertilizer might have resulted in under-fertilizer application in a location but over-fertilizer application in the other location. Under-application in some locations restricted crop production whereas over-application resulted in high risk of pollution of environment and also increased cost for production (Verhagen *et al.*, 1995; Booltink *et al.*, 2001; Verhagen, 1997; Bouma, 1997).

To prescribe a site-specific N fertilizer amount for rice crop field, it is important to understand the response of grain yield and quality to rates of nitrogen application in combination with spatial variation in soil properties, plant growth status before N application and other factors. Spatial variation in crop yield response to fertilizer nitrogen under variable soil conditions for winter wheat, spring barley (Delin *et al.*, 2002), and for maize (Kahabka *et al.*, 2004) have been reported. Although variation in plant growth and N nutrition status of rice crop was reported to be an important factor determining rice yield and its response to N application at panicle initiation stage (PIS) (Cui & Lee, 2002; Peng *et al.*,

†Corresponding author (Phone) +82-02-880-4544 (E-mail) leebw@snu.ac.kr

<Received August 26, 2005>

1996), consideration of variation in plant growth and N nutrition status for variable N rate prescription was rare or not available in Korea and other countries.

The fertilizer prescription based on the measurement of plant growth status is usually laborious, expensive and cannot meet the time requirement. However, recently a fast and inexpensive *in-situ* measurement for crop growth variables such as biomass, nitrogen concentration, and uptake using advanced remote sensing technique have been reported (Casanova *et al.*, 1998; Hansen & Schjoerring, 2003; Nguyen & Lee, 2004), promising the use of crop growth status for site specific nitrogen management.

Information of spatial variability of rice yield and its causal factors obtained previously at the two trial fields in 2002 and 2003 indicated that plant growth, soil properties, and grain yield were highly variable within the present trial field (Nguyen *et al.*, 2004). However, a question of how to manage within-field spatial rice yield variability is still remained. An experiment was carried out to test whether variation in plant growth status may be used for the prescription of variable N rate top-dressing at PIS. The objectives of the study, therefore, were (1) to formulate the site-specific prescription rule for prescribing nitrogen topdressing amount at PIS using available spatial plant variability information and (2) to test if the prescription rule significantly reduces spatial variability of rice yield and protein content as well as improving rice yield.

MATERIALS AND METHODS

Site description and Experimental design

A paddy field located in experimental farm of National Institute of Crop Science, Rural Development Administration (37°16'N), Korea was designed as a trial field for testing a possibility of management of spatial yield variability of rice in 2004. The field was used to investigate spatial variation of plant growth and yield in relation to soil properties under direct-seeded rice and the uniform fertilizer application of 110 N: 70 P: 80 K (kg/ha) in 2002, and treated with no N fertilizer under transplanting rice culture in 2003.

The field with areas of 6,600 m² was divided schematically into 66 grids (10 m × 10 m) (Fig. 1). Whole field was managed uniformly in 2002 and 2003 throughout the rice growing season. However, An upper part consisting of 33 grids was applied with prescribed N topdressing amount (VRT) while lower part consisting of 33 grids was applied with conventional uniform nitrogen top-dressing (UN) of 33 kg N/ha at PIS in 2004. The whole field was applied with 44 and 33 kg N/ha as basal and tillering fertilizer, respectively. Potassium fertilizer (80 kg/ha) was applied in two splits of




Fig. 1. Design of experimental field for site-specific N topdressing at panicle initiation stage in 2004 (upper 30 grids were used for variable rates of N topdressing (VRT) and lower 30 grids for conventional uniform management (UN) with 110 N kg/ha)

70-30% as basal-panicle fertilizer while phosphorus fertilizer (70 kg/ha) in one split (100%) as basal fertilizer.

Crop cultivation and Plant growth measurements

Japonica rice (*Oryza sativa* L.) variety “Daeanbyeon” was machine-transplanted. Field management practices followed a standard practices typical for Korean farmer’s field. Plant growth parameters such as shoot dry weight (DW), number of tiller (Til), shoot nitrogen concentration (SN), shoot nitrogen uptake (Nup) and SPAD reading (Minolta SPAD 502), etc. were measured at panicle initiation stage (PIS), heading stage (HD) and harvest stages (HA) of rice. Nitrogen nutrition index (NNI) was calculated with measured SN and DW according to Cui *et al.* (2000). Tiller number was counted for 20 hills at the center part of each grid, and the five hills were randomly sampled for shoot dry weight and nitrogen concentration measurement. Sampled plants were dried at 70 °C for two days, weighed for DW, and ground through 40-mesh sieve for nitrogen analysis. Nitrogen concentration was analyzed by CNS analyzer (Leco CNS 2000). Grain yields were measured with sample harvested from the area of 6 m² for each grid. Final yield of rough rice was adjusted to 14% of water content.

Prescription of site-specific nitrogen topdressing at panicle initiation stage

Prescription formula of topdressing N amount

Nitrogen topdressing amount at panicle initiation stage was basically calculated based on the following formula:

$$P_{SFN} = \frac{N \text{ uptake} - \text{Natural N supply}}{N \text{ recovery from PIS to harvest}} \quad (\text{Eq.1})$$

Where,

P_{SFN} : N amount prescribed for site-specific N fertilizer

Natural N supply: Natural N supply from PIS to harvest (estimated from 2003 experiment in the same field where no N fertilizer application was applied).

N recovery from PIS to harvest: recovery of N applied at panicle initiation stage (60% was used as reported by Kim (2004).

N uptake: uptake from PIS to harvest required for target protein content of milled rice (P).

The calculation of N uptake from PIS to harvest for target protein content of milled rice was presented in the next section.

Estimation of N uptake requirement for targeting grain yield and protein content in relation to plant growth parameters at panicle initiation stage

In order to estimate nitrogen uptake amount (from panicle initiation to harvest stage) required for target rice yield and protein content, we firstly formulated the regression equation of rice yield (Y) and protein content (P) response to plant

Table 1a. Regression equations of gram yield (Y) and protein content (P) response to plant growth parameters.

Items	Regression equations	R ²
Protein content (P)	$P=4.69+0.0144Nup+0.639NNI$	0.950
	$P=5.00+0.0144Nup-0.238DW+0.081DW^2$	0.956
	$P=4.433+0.0186Nup-0.000Nup^2+0.235SN$	0.942
Rice yield (Y)	$Y=2266+27.9Nup+1517NNI$	0.984
	$Y=5.00+0.0144Nup-0.23SDW+0.081DW^2$	0.989
	$Y=2304+28.1Nup+77.9SN^2$	0.979

P: protein of rice (%)

Y: rice yield (kg/ha)

Nup: N uptake from panicle initiation stage (PIS) to harvest (kg/ha)

NNI: Nitrogen nutrition index (Cui *et al.*, 2002) at PIS

DW: shoot dry weight at PIS (ton/ha)

SN: Shoot nitrogen concentration at PIS (%)

Table 1b. Equations to estimate N uptake amount (Nup) from panicle initiation stage (PIS) to harvest stage required for target rice yield (Y) and grain protein content (P) in relation to plant growth parameter at PIS.

Item	N uptake (Nup) required for targeting
Protein content of grain (P)	$Nup=(P-4.691-0.639NNI)/0.0144$
	$Nup=(P-5.008+0.238DW-0.081SDW^2)/0.0144$
	$Nup=(P-4.433+0.000Nup^2-0.235SN)/0.0186$
Rice yield (Y)	$Nup=(Y-2266-1517NNI)/27.9$
	$Nup=(Y-5.00+0.23DW-0.081DW^2)/0.0144$
	$Nup=(Y-2304-77.9SN^2)/28.1$

growth parameters such as shoot dry weight (DW), shoot nitrogen concentration (SN), nitrogen nutrition index (NNI) at PIS and nitrogen uptake (Nup) from PIS to harvest with the data obtained from the previous two-year experiments (Table 1a). Equations from Table 1a showed that crop parameters at PIS in combination with Nup successfully explained rice yield and protein content of milled rice ($R^2>0.94$). Thus, we converted these equations to get Nup for target yield and protein content (Table 1b).

To estimate N top-dressing amount at PIS using Eq.1, Nup for each grid has to be calculated from the equations in Table 1b with the target rice yield/protein content and the measured or predicted plant growth parameters prior to N topdressing at PIS. In this study, we applied canopy reflectance method using hand-held spectroradiometer (GER Inc. USA; GER 1500) for in-situ determination of crop growth parameters at PIS. The detailed information of the method may be found in reports from Nguyen & Lee (2004), Nguyen *et al.* (2004), and Hansen & Schioerring (2003), and the details of the applied method for determination of plant growth variables at PIS in this study were presented in the next section.

Prediction of plant parameters using canopy reflectance at panicle initiation stage

To estimate Nup for target yield and protein content in the present experiment, canopy reflectance of rice before applying N fertilizer at PIS was measured by GER 1500. The procedure of canopy measurement using GER 1500 for the prediction of rice crop growth variables using canopy reflectance data were described in detail by Nguyen & Lee (2004) and Nguyen *et al.* (2005). The measured DW was presented in Fig. 2, indicating that rice growth is spatially variable prior to nitrogen top-dressing at PIS while plant parameters such as DW, NNI and SN predicted from canopy




Fig. 2. Kriged map of spatial variation of shoot dry weight (DW) at panicle initiation stage prior to nitrogen topdressing.




Fig. 3. Comparison of nitrogen nutrition index (NNI) and shoot dry weight (DW) at panicle initiation stage between measured and predicted by canopy reflectance for N topdressing prescription.




Fig. 4. Comparison of N topdressing amount prescribed based on plant growth variables (NNI, SN and SDW) estimated by canopy reflectance at panicle initiation stages.

reflectance data by partial least square method were presented in comparison with the observed values in Fig. 3. The significant close correlation between predicted and observed values of dry weight and NNI in Fig. 3 ($R^2 > 0.73$, $n=66$) suggested that partial least square using canopy reflectance of rice is reliable method for in-situ rice growth status measurement.

Prescribed N topdressing amount

Nitrogen topdressing amount were prescribed for targeting rice protein content of 6.8 % instead of targeting rice grain yield as the Nup for maximizing rice grain yield is too high because of the decreasing internal N use efficiency at high N level (Witt *et al*, 1999) and protein content is a good criterion of rice quality in terms of its palatability.

Nitrogen topdressing amounts prescribed based on predicted NNI, DW, and SN were compared in Fig. 4 and they showed good agreement each other with $R^2 > 0.99$. The prescribed nitrogen amounts ranged from 0 to 110 kg N/ha, depending on each grid of the field and the average was 57 kg/ha. The spatial variation of protein content in year 2003 without fertilizer application (Fig. 5a) was presented in comparison with the prescribed N topdressing amount in this study in year 2004 (Fig. 5b). The grid that higher protein content occurred was prescribed with less nitrogen topdressing amount.

Data analysis

Descriptive statistical analysis was used for presenting the




Fig. 5. Kriged map of spatial variation of grain protein content at the experimental field in 2003 (A) and prescribed N topdressing amount (B) (kg/ha) for targeting protein content of 6.6% (VRT: 33 grids treated with prescribed N amount and UN: 33 grids with uniform application of (33 N kg/ha at panicle initiation stage).

spatial variability of rice yield and protein content: mean, minimum, maximum, range, and coefficient of variation (CV). Multiple linear regression was used to formulate regression equation of rice yield and protein content response to plant growth variables, including NNI, SN and DW. Partial least square (PLS) procedure (Nguyen *et al.*, 2004) was used to predict NNI, SN and DW from canopy reflectance data measured by GER 1500 at PIS. Kriging map of spatial variation of yield and protein content and other related parameters was performed using Acview GIS. Data analysis and processing were carried out using SAS (SAS institute inc. 8.12).

RESULTS

Comparison of spatial plant growth variability between VRT and UN

Variation of plant growth at panicle initiation stage (PIS) and heading stages (HD) of VRT and UN in comparison with no N treatment in year 2003 were presented in Table 2. Most of plant parameters were highly variable in prescribed nitrogen application (VRT) and conventional uniform nitrogen application (UN) at PIS. Coefficient of variation (CV) ranged from 5.3% for SPAD reading to 22.4% for shoot

Table 2. Comparison of spatial variation of growth parameters between the prescribed and uniform N topdressing (33 grids).

Parameter		Panicle initiation stage		Heading stage		Harvest stage	
		Prescribed Nitrogen	Uniform Nitrogen	Prescribed Nitrogen	Uniform Nitrogen	Prescribed Nitrogen	Uniform Nitrogen
DW	Range	1.7-4.1	2.3-3.7	5.2-10.3	5.2-9.4	8.0-14.6	7.1-15.8
	C.V	22.1	14.0	15.5	15.1	17.1	22.4
SN	Range	1.3-2.0	1.5-2.0	1.0-1.8	1.0-1.7	0.7-1.2	0.8-1.2
	C.V	11.4	6.98	10.8	13.1	13.1	14.2
Nup	Range	26-66	41-79	54-176	55-152	67-147	62-187
	C.V	22.4	18.7	20.4	25.5	20.1	30.2
NNI	Range	0.3-0.6	0.4-0.6	0.4-0.9	0.4-0.8	ND	ND
	C.V	15.7	8.3	19.7	13.1	ND	ND
SPAD	Range	29-38	32-38	31-45	30-43	ND	ND
	C.V	9.99	5.33	8.0	8.31	ND	ND

DW: Shoot dry weight (ton/ha)

SN: Shoot nitrogen concentration (%)

Nup: Shoot nitrogen uptake (kg/ha)


NNI: Nitrogen nutrition index

SPAD: SPAD reading

ND: Not measured

Table 3. Comparison of spatial variation of rice yield between 33 grids with prescribed N and 33 grids with conventional uniform N

	Prescribed field section		Conventional field section	
	No fertilizer 2003	Prescribed N 2004	No fertilizer 2003	Uniform N 2004
	----- Kg/ha -----			
Minimum	3323	6413	3204	5546
Maximum	6171	9249	5541	8672
Mean	4460	7954	4207	7019
Standard deviation	954.0	647.9	656.0	1033
Coefficient of variation (%)	21.3	8.13	15.5	14.7

**Fig. 6.** Kriged maps of spatial variation of rice yield at the experimental field subjected to uniform and prescribed N topdressing at panicle initiation stage (NF: No fertilizer application in 2003, UN and VTR: uniform and variable rate of N topdressing in 2004, respectively).

nitrogen uptake. Some plant growth parameters such as DW, SN, and NNI at VRT field seem to be reduced at heading stage compared with UN (Table 2). Moreover, at harvest stage, variation of plant growth parameters at the field with VRT were significantly reduced compared with UN (CV for shoot nitrogen uptake at VRT was 17% while at UN is 22%). Shoot nitrogen uptake was also found to be the most variable among plant growth parameters of rice while SPAD reading was the least variable parameter at all stages (Table 2).

Comparison of spatial variability of rice yield and grain protein content between VRT and UN.

Spatial variation of rice yield

Spatial variation of rice yield at VRT and UN were presented in Table 3. Rice yield at VRT ranged from 6000 to 9000 kg/ha while 5500 to 8500 kg/ha was recorded for UN. An average of about 960 kg/ha of rice at VTR was higher than that of UN.

Spatial rice yield variability in 2004 was also significantly reduced at VTR in comparison with UN. Coefficients of variation were 8.13% and 14.6% for VRT and UN, respectively. Data of rice yield measured under no fertilization

condition in 2003 at the same field as in the present study of 2004 was also presented in Table 3 for comparison of the variation between 2003 and 2004. The spatial variability of rice yield was significantly reduced from 21.3 % in 2003 (no fertilization) only to 8.1% in 2004 (VRT) in VRT prescription, while no conceivable difference in the spatial variability was detected between years in UN. The kriged map of spatial variation distribution of rice yield at VTR and UN field were shown in Fig. 6. These maps also clearly indicate that spatial variation of rice yield was reduced due to the site-specifically prescribed nitrogen topdressing.

Spatial variation in grain protein content of milled rice

Variation of grain protein content at VRT was also significantly reduced compared with UN (Table 4). Grain protein content of milled rice at VRT ranged from 5.60% to 7.92% with CV of 7.05% while at UN it ranged from 4.41-7.66% with CV of 13.0%. And spatial variability of protein content was rather lower in 2004 (VRT N) than 2003 (no fertilizer application) at the VRT field, while the CVs increased significantly from 6.51% in 2003 to 13% in 2004 at UN field. kriged map of spatial variation distribution of protein content at VRT and UN were also compared (Fig. 7). These




Fig. 7. Kriged maps of spatial variation of rice protein content at the experimental field subjected to uniform and prescribed N topdressing at panicle initiation stage (NF: No fertilizer application in 2003, UN and VTR, uniform and variable rate of N topdressing in 2004, respectively).

Table 4. Comparison of spatial variation of grain protein content between 33 grids for the prescribed N and 33 grids with conventional uniform N.

	Prescribed field section		Conventional field section	
	No fertilizer 2003	Prescribed N 2004	No fertilizer 2003	Uniform N 2004
	----- % -----			
Minimum	4.68	5.60	4.73	4.41
Maximum	6.29	7.92	5.92	7.66
Mean	5.34	6.87	5.28	6.04
Standard deviation	0.44	0.52	0.35	0.91
Coefficient of variation (%)	8.23	7.05	6.51	13.0

maps also clearly indicate that spatial variation of rice protein content was reduced due to the site-specifically prescribed nitrogen topdressing.

DISCUSSION

Procedures to derive fertilizer recommendation for site-specific crop management

Methods for measurement of spatial crop variability were reported by Seney (1998), and Naiqian (2002) discussed several alternative managements of spatial crop variability. Variable rate treatment (VRT) is often used for managing the spatial variability, for example, VRT of fertilizer application, seed or pesticides etc. Estimation of nitrogen requirements is an important technique for VRT of fertilizer application for managing crop spatial variability (O'Neal *et al.*, 2000). In this experiment, the method for VRT of N top-dressing requirement at PIS was derived for rice, and examined its feasibility.

Regression equations for predicting rice yield and protein

content were well formulated using the information on rice yields, protein contents, plant nitrogen and biomass etc. from the previous year experiment in 2003 (Table 1.a and Table 1.b). In addition, partial least square (PLS) procedure was used to predict DW, NNI, and SN from canopy reflectance with acceptable precision and accuracy (Fig.3). With these procedures, nitrogen uptake (N_{up}) required from PIS to harvest for target rice yield or grain protein content could be well estimated and used for the calculation of variable nitrogen topdressing prescription. Site-specific N topdressing amount determined for targeting rice protein content ranged from 0 to 110 kg N/ha with average of 57 kg N/ha. The high spatial variation in prescribed N levels (Fig. 5) reflected the high spatial variation of natural soil N supplies that was obtained in the previous experiment of year 2003.

The method for site specific N fertilizer prescription for managing spatial rice yield variability is successful only if basic data on spatial variation in soil conditions and variation in plant growth status at PIS before N prescription of the same year are available. Firstly, the basic field information of the study sites includes spatial distribution of natural

soil N supplies and N recovery as well as spatial variation in yield or protein content in response to plant growth status at PIS. Although this information may be obtained from field research, its temporal variation among years should be considered. If the relative temporal variation in a given parameter is small compared to spatial variation, the spatial variation information obtained from one year or average over several years are useful information for site specific N fertilizer prescription, otherwise it is useless. That is why quite stable soil parameters such as CEC, clay content or organic matter were preferred to other high temporal variation parameters such as $\text{NO}_3\text{-N}$ and exchangeable cations (Dobermann, 1994; Cox *et al.*, 2003). Secondly, the prediction of crop variables at PIS by fast and nondestructive canopy reflectance method is required (Nguyen & Lee, 2004; Nguyen *et al.*, 2004; Hansen & Schjoerring, 2003). However, the precision and accuracy of the method depends on several factors such as weather condition, data analysis and skills of technicians.

Spatial variation of plant growth under VRT and UN

Spatial yield variability of rice was significantly caused by plant growth variations due to soil conditions (Dobermann, 1994; Casanova *et al.*, 1999) and management of plant growth variation was considered important for site-specific crop management (Naiqian, 2002). Variation of plant growth under VRT and UN were measured at several growth stages of rice crop. It is expected that most of plant growth parameters were equally varied at PIS because the whole field was managed uniformly until PIS. The differences in mean and CV of plant growth at PIS between two parts (VRT and UN) were probably due to different variations in soil properties. At heading stage, some plant parameters in VRT turned out to reduce the variation suggesting that prescription of nitrogen topdressing potentially reduced the variation in plant growth. The variation of plant growth reduced by VRT was clearer at harvest, most of plant parameters such as shoot N concentration and uptake being smaller in CV at VRT than at UN parts (Table 2).

Spatial yield variability and grain protein contents under VRT and UN

Obtained results of spatial variability of rice yield and protein content at VRT and UN (Table 3 and 4) confirmed that the method and procedure we developed for prescribing nitrogen topdressing at panicle initiation stage have successfully worked to manage spatial yield and protein variability in paddy field. Variation of rice yields and grain protein content of VRT were significantly reduced compared with UN.

Kriged maps of rice yield and grain protein contents (Fig. 6 and 7) also presented quite homogeneous distribution in all grids of VRT while they were highly variable among grids of UN.

One of the limitations of the study possibly was that the results did not show the N use efficiency or economic return from VRT in comparison with UN. The limitation was due to lack of a control treatment without N application at PIS at VRT and UN parts. We would like to recommend this remaining issue for further study. In conclusion, the research was carried out as just a case study for confirming the possibility of spatial variability management of rice crop. However, there is no doubt to recommend the result from this research for further examination to implement the site-specific crop management in large-scale rice field in Korea.

CONCLUSION

A possibility for management of spatial yield variability and grain protein contents using prescribed nitrogen topdressing at panicle initiation stage have been examined in a paddy field of Korea and obtained results of this research can be summarized as follows: The prescription of N fertilizer requirement for management of spatial variation in rice yield and protein contents at PIS using basic spatial information for previous study and in-situ prediction of variation in plant growth status was formulated successfully. Spatial variation of plant growth at VTR turned out to reduce slightly at HD after the prescription treatment of nitrogen topdressing and, especially, spatial variation of plant growths was significantly reduced at harvest stage. Design of experimental field for testing a possibility of management of spatial yield variability using prescription of nitrogen at PIS has successfully worked not only to reduce spatial yield and protein content variability but also increase rice yield in a paddy field. The developed method and procedure for estimation of nitrogen topdressing amount for site-specific fertilizer management was quite reliable. However, this was just a case study, further improvement is necessary before this procedure could be practically used for the site-specific rice management in large-scale rice field.

ACKNOWLEDGEMENT

This work was supported by a grant from Agricultural Research Promotion Center, Korea.

REFERENCES

- Bahman Eghball and Gary E. Varvel. 1997. Fractal analysis of temporal yield variability of crop sequences implication for

- site-specific management. *Agron. J.* 89: 851-855
- Beckett, P. H. T. and R. Webster. 1971. Soil variability: a review *Soils Fert.* 34, pp. 1-15.
- Booltink, H. W. G., G. J. Alphen, W. D. Batchelor, J. O. Paz, J. J. Stoorvogel, and R. Vargas. 2001. Tool for optimizing management of spatially variable fields. *Agricultural System* 70: 445-476.
- Bouma, J. 1997. Precision agriculture. introduction to the spatial and temporal variability of environmental quality. In: Lake, J.V., Bock, G.R., Goode, J.A (Eds.), *Precision Agriculture. Spatial and Temporal Variability of Environmental Quality.* Ciba Foundation Symposium, 210. Wiley, Wageningen, The Netherlands, pp 5-17.
- Cahn, M. D., J. W. Humel, and B. H. Brouer. 1994. Spatial analysis of soil fertility for site-specific crop management *Soil Sci Soc. Am J* 58:1240-1248.
- Casanova, D., G. F. Epema, and J. Goudriaan. 1998. Monitoring rice reflectance at field level for estimating biomass and LAI *Field Crops Res.* 83-92
- Casanova, D., J. Goudriaan, J. Bouma, and G. F. Epema. 1999. Yield gap analysis in relation to soil properties in direct-seeded flooded rice *Geoderma* 91(3-4): 191-216.
- Cox, M. S., P. D. Gerard, M. C. Wardlaw, and M. J. Abshire. 2003. Variability of selected soil properties and their relationships with soybean yield. *Soil Sci. Soc. Am. J.* 67. 1296-1302.
- Cui, R. X. and B. W. Lee. 2002. Spiklet number estimation model using nitrogen nutrition status and biomass at panicle initiation and heading stage of rice. *Korean J. Crop Sci.* 47 (5): 390-394.
- Cui, R. X., M. H. Kim, J. H. Kim, H. S. Nam, and B. W. Lee. 2002. Determination of critical nitrogen dilution curves for rice growth. *Korean J. Crop Sci.* 47 (2). 127-131.
- Delin, S. and B. Linden. 2002. Relations between net nitrogen mineralization and soil characteristics within an arable field. *Acta Agr. Scand. B-S P* 52, pp 78-85.
- Dobermann, A., A. F. Pampolino, and H. U. Neue. 1995. Spatial and temporal variability of transplanted rice at field scale. *Agron. J.* 87: 712-720.
- Dobermann, D. 1994. Factors causing field variation of direct-seeded flooded rice *Geoderma* 62(1-3) . 125-150.
- Environmental Systems Research Institute. Arcview GIS (ESRI), 1996.
- Hansen, P. M. and J. K. Schjoerring. 2003. Reflectance measurement of canopy biomass and nitrogen status in wheat crops using normalized difference vegetation indices and partial least squares regression. *Remote Sens. of Environ.* 86: 542-553.
- Jin J. and C. Jiang. 2002. Spatial variability of soil nutrient and site-specific management in the P.R. China *Computer and Electronics in Agriculture* 36 165-172
- Kahabka, J. E., H. M. Van Es, E. J. McClenahan, and W. J. Cox. 2004. Spatial analysis of maize response to nitrogen fertilizer in central New York. *Precision Agriculture* 5: 463-476.
- Machado, S., E. D. Bynum, T. L. Archer, R. J. Lascano, L. T. Wilson, J. Bordovsky, E. Segarra, K. Bronson, D. M. Nesmith, and W. Xu. 2000. Spatial and temporal variability of corn grain yield: site-specific relationships of biotic and abiotic factor. *Precision Agriculture* 2: 359-376
- McBratney, A. B. and M. J. Pringle. 1997. Spatial variability in soil implications for precision agriculture. In: Stafford, J. V. Ed. *Spatial Variability in Soil and Crop.* Precision Agriculture '97 I BIOS, Oxford, pp 1-31
- Miller, P. M., M. J. Singer, and D. R. Nielsen. 1995. Spatial variability of wheat yield and soil properties on complex hills. *Soil Sci.Soc.Am.J.* 52:1133-1141.
- Naiqian, Z., M. Wang, and N. Wang. 2002. Precision agriculture-a worldwide overview
- Nguyen T. Hung and B. W. Lee. 2004. Selection of most sensitive waveband reflectance for normalized difference vegetation index calculation to predict rice crop growth and yield. *Korean J. Crop. Sci.* 49(5): 394-406.
- Nguyen T. Hung, J. H. Kim, Nguyen Tuan Anh, and B.W. Lee. 2005. Application of hyperspectral canopy reflectance measurement and partial least square regression to prediction of crop growth and nitrogen status before heading stage of rice. *Precision Agriculture* 05 : 251-260.
- Nguyen Tuan Anh, J. C. Shin, and B. W. Lee. 2004. Analysis of spatial variation of yields and soil chemical properties in a paddy field. *Proc. Inter. Crop Sci Congress*, pp. 425 Australia.
- O'Neal, M. R., J. R. Frankenberger, and D. R. Ess. 2000. Spatial precipitation variability in the choice of nitrogen fertilization rates. *Proceedings of Fifth International Conference on Precision Agriculture (CD)*, July 16_/19, 2000. Bloomington, MN, USA
- Paz, J. O. 2000. Analysis of Spatial Yield Variability and Economic Prescription for Precision Agriculture: a Modeling Approach. Ph D. dissertation, Iowa State University, Ames, Iowa.
- Peng, S., F. V. Garcia, R. C. Laza, A. L. Sanico, R. M. Visperas, and K. G. Cassman. 1996. Increased N-use efficiency using a chlorophyll meter on high-yielding irrigated rice. *Fields Crops Res.* 47: 243-252.
- SAS Institute, 2001. SAS System for Windows Release 8.01. SAS Institute Inc. Cary, N.C.
- Seney, G. B., A. D. Ward, J. G. Lyon, N. R. Fausey, and S. E. Nokes. 1998. Manipulation of high spatial resolution aircraft remote sensing data for use in site-specific farming. *Trans. ASAE* 41 (2) : 489-495
- Verhagen, J. 1997. Site specific fertiliser application for potato production and effects on N leaching using dynamic simulation modeling *Agri Ecos and Environ* 66(2) · 165-175
- Verhagen, J., H. W. G. Bootink, and J. Bouma. 1995. Site specific management. balancing production and environmental requirements at farm level. *Agric. Syst.* 49 : 369-384.
- Yanai, J., C. K. Lee, M. Umeda, and T. Kosaki. 2000. Spatial of soil chemical properties in a paddy field *Soil Sci. Plant Nutr.* 4 : 473-482.