

## DIRECT PROOF OF EKELAND'S PRINCIPLE IN LOCALLY CONVEX HAUSDORFF TOPOLOGICAL VECTOR SPACES

JONG AN PARK

ABSTRACT. A.H.Hamel proved the Ekeland's principle in a locally convex Hausdorff topological vector spaces by constructing the norm and applying the Ekeland's principle in Banach spaces. In this paper we show that the Ekeland's principle in a locally convex Hausdorff topological vector spaces can be proved directly by applying the famous general principle of H.Brézis and F.E.Browder.

### 1. Introduction

H.Brézis and F.E.Browder[1] put forward the following general principle in nonlinear functional analysis which unifies the proofs of Ekeland's variational principles[3] and Caristi-Kirk fixed point theorem [2]and Bishop-Phelps lemma and Daneš' drop theorem. Also the invariance theorems for closed sets under flows in metric spaces were proved by the same principle in [1]. We define  $S(x) = \{y \in X | y \geq x\}$  in a ordered set  $X$ .

THEOREM 1.1. [1] *Let  $X$  be a Hausdorff topological space with an ordering structure. Let  $\psi : X \rightarrow \mathbb{R}$  be a function bounded below. Assume*

1.  $S(x)$  is sequentially closed for each  $x \in X$ ;
2.  $x \leq y$  and  $x \neq y$  imply  $\psi(y) < \psi(x)$ ;
3. any increasing sequence is relatively compact.

*Then for each  $a \in X$  there exists  $\bar{a} \in X$  such that  $a \leq \bar{a}$  and  $\bar{a}$  is maximal.*

---

Received December 22, 2004.

2000 Mathematics Subject Classification: Primary 49J40, 46A03.

Key words and phrases: lower semi-continuous, Minkowski functional, Ekeland's variational principle.

We define the Minkowski functional  $\mu_S$  of a subset  $S$  of the topological vector space  $X$  to be

$$\mu_S(x) := \inf\{t > 0 : x \in tS\}.$$

And we define  $\text{dom } \mu_S = \bigcup_{t>0} tS$  and  $\mu_S(x) = \infty, x \notin \text{dom } \mu_S$ . Let  $\{p_\lambda\}_{\lambda \in \Lambda}$  be a base of continuous seminorms generating the topology on a locally convex topological space  $X$ . Then we call  $X$  be a sequentially complete iff every  $p_\gamma$ -Cauchy sequence converges. Furthermore  $f : X \rightarrow (\infty, \infty]$  is called proper if  $\{x | f(x) < \infty\} \neq \emptyset$  and let  $\text{dom } f = \{x | f(x) < \infty\}$  and it is a sequentially lower semi-continuous function iff for every  $c \in \mathbb{R}$ ,  $\{x \in X | f(x) \leq c\}$  is sequentially closed.

LEMMA 1.1. *Let  $S \subset X$  be a sequentially closed, bounded and convex set of a Hausdorff locally convex topological space  $X$  containing 0. Then the followings hold;*

1.  $\mu_S : X \rightarrow [0, \infty]$  is an extended-valued proper and sequentially lower semi-continuous function
2. for any  $x, y \in \text{dom } \mu_S$  we have  $x + y \in \text{dom } \mu_S$  and

$$\mu_S(x + y) \leq \mu_S(x) + \mu_S(y)$$

3. for any  $x, y \in \text{dom } \mu_S, x - y \in \text{dom } \mu_S$  we have

$$\mu_S(x) - \mu_S(y) \leq \mu_S(x - y).$$

*Proof.* 1. Clearly since  $0 \in S, \mu_S(0) = 0$ ,  $\mu_S$  is proper. We must prove that  $C_c = \{x \in X | \mu_S(x) \leq c\}$  is sequentially closed for any  $c \in [0, \infty]$ . Indeed if  $c = \infty$ ,  $C_c = X$  is sequentially closed. if  $0 \leq c < \infty$ , and  $x_n \in C_c$  and  $x_n \rightarrow y$ , then  $\mu_S(x_n) \leq c$ . Hence for each  $n$  there exists  $\alpha_n, s_n \in S$  such that

$$0 \leq \alpha_n \leq c + \frac{1}{n}, x_n = \alpha_n s_n.$$

Suppose  $\alpha_{n_i} \rightarrow 0$  for some  $n_i$ , then  $x_{n_i} = \alpha_{n_i} s_{n_i} \rightarrow 0$  because  $S$  is bounded. That is,  $y = 0$  and  $\mu_S(y) = 0 \leq c, y = 0 \in C_c$ . Suppose  $0 < \delta \leq \alpha_n \leq c + \frac{1}{n}$  for all sufficiently large  $n$ , then  $\alpha_{n_i} \rightarrow \alpha$  for some  $n_i$  and  $0 < \delta \leq \alpha \leq c$ . Therefore

$$\frac{x_{n_i}}{\alpha_{n_i}} = s_{n_i} \rightarrow \frac{y}{\alpha}.$$

Since  $S$  is sequentially closed,  $\frac{y}{\alpha} \in S$  and  $y \in \alpha S, \mu_S(y) \leq \alpha \leq c$ . Hence  $y \in C_c$ .

2. Suppose  $x, y \in \text{dom } \mu_S$ , then for any  $s, t > 0$  such that  $\mu_S(x) < s, \mu_S(y) < t$  we have  $\frac{x}{s}, \frac{y}{t} \in S$  because  $S$  is convex and  $0 \in S$ . Let  $u = x + y$  and from the convexity of  $S$  we have

$$\frac{x + y}{u} = \left(\frac{s}{u}\right)\frac{x}{s} + \left(\frac{t}{u}\right)\frac{y}{t} \in S.$$

So  $x + y \in uS, \mu_S(x + y) \leq u = s + t$ . Therefore  $\mu_S(x + y) \leq \mu_S(x) + \mu_S(y)$ .

3. for any  $x, y \in \text{dom } \mu_S, x - y \in \text{dom } \mu_S$ , by 2

$$\mu_S(x) = \mu_S((x - y) + y) \leq \mu_S(x - y) + \mu_S(y)$$

That is

$$\mu_S(x) - \mu_S(y) \leq \mu_S(x - y).$$

□

We relate the Minkowski functional  $\mu_S$  with a family of continuous seminorms  $p_\lambda$ .

LEMMA 1.2. *Let  $S \subset X$  be a sequentially closed, bounded and convex set of a Hausdorff locally convex topological space  $X$  containing  $0$ . Let  $\{p_\lambda\}_{\lambda \in \Lambda}$  be a base of continuous seminorms generating the topology on  $X$ . Then there exists  $\{\alpha_\lambda\}_{\lambda \in \Lambda}$  a family of positive numbers such that*

$$\frac{1}{\alpha_\lambda} p_\lambda(x) \leq \mu_S(x), x \in \text{dom } \mu_S.$$

*Proof.* Since  $S$  is bounded in  $X$ , for any  $\lambda \in \Lambda$  there exists  $\alpha_\lambda > 0$  such that  $S \subset \alpha_\lambda U_\lambda$ , where  $U_\lambda := \{x \in X \mid p_\lambda(x) \leq 1\}$ . Then  $\mu_{U_\lambda} = p_\lambda$ . Therefore for any  $x \in S$

$$\frac{1}{\alpha_\lambda} p_\lambda(x) = \frac{1}{\alpha_\lambda} \mu_{U_\lambda}(x) \leq \mu_S(x).$$

Suppose  $x \in \text{dom } \mu_S$  and  $\mu_S(x) < t$  then  $x \in tS, x = ts, t > 0, s \in S$ . That is,

$$\frac{1}{\alpha_\lambda} p_\lambda(x) = \frac{1}{\alpha_\lambda} p_\lambda(ts) = \frac{t}{\alpha_\lambda} p_\lambda(s) \leq t \mu_S(s) \leq t$$

because  $\mu_S(s) \leq 1$  for any  $s \in S$ . Hence

$$\frac{1}{\alpha_\lambda} p_\lambda(x) \leq \mu_S(x), x \in \text{dom } \mu_S.$$

□

## 2. Main proof

Instead of the methods of proof in A.H.Hamel [4] we prove the following Ekeland's principle in locally convex Hausdorff spaces by using the above general principle directly.

**THEOREM 2.1.** *Let  $X$  be a Hausdorff locally convex topological space that is sequentially complete. Let  $f : X \rightarrow \mathbb{R}$  be a sequentially lower semi-continuous function, bounded below. Let  $\{p_\lambda\}_{\lambda \in \Lambda}$  be a base of continuous seminorms generating the topology on  $X$  and  $\{\gamma_\lambda\}_{\lambda \in \Lambda}$  a family of positive numbers. Then for every  $x_0 \in X$  there exists  $x^* \in X$  such that*

$$f(x^*) + \gamma_\lambda p_\lambda(x^* - x_0) \leq f(x_0)$$

for all  $\lambda \in \Lambda$ , and for all  $x \in X, x \neq x^*$  there exists  $\mu \in \Lambda$  such that

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*).$$

*Proof.* Let  $X$  be equipped with an ordering

$$x \leq y \text{ iff } f(y) - f(x) \leq -\gamma_\mu p_\mu(y - x) \quad \forall \mu \in \Lambda.$$

Indeed it is an ordering on  $X$ . That is,

- $x \leq x$
- $x \leq y$  and  $y \leq x$  imply  $x = y$
- $x \leq y$  and  $y \leq z$  imply  $x \leq z$

Clear  $\leq$  is reflexive. If  $x \leq y$  and  $y \leq x$ , then

$$f(y) - f(x) \leq -\gamma_\mu p_\mu(y - x) \quad \forall \mu \in \Lambda$$

and

$$f(x) - f(y) \leq -\gamma_\mu p_\mu(x - y) \quad \forall \mu \in \Lambda.$$

Therefore we have

$$0 \leq -\gamma_\mu (p_\mu(y - x) + p_\mu(x - y)) \quad \forall \mu \in \Lambda.$$

Hence  $p_\mu(x - y) \leq -p_\mu(y - x) \quad \forall \mu \in \Lambda$ . Since  $p_\mu \geq 0$ , we have  $p_\mu(x - y) = 0 \quad \forall \mu \in \Lambda$ . Since  $X$  is Hausdorff,  $x = y$ . We showed that  $\leq$  is antisymmetric. To prove that  $\leq$  is transitive, if  $x \leq y$  and  $y \leq z$ , then

$$f(y) - f(x) \leq -\gamma_\mu p_\mu(y - x) \quad \forall \mu \in \Lambda$$

and

$$f(z) - f(y) \leq -\gamma_\mu p_\mu(z - y) \quad \forall \mu \in \Lambda.$$

Then

$$f(z) - f(x) \leq -\gamma_\mu (p_\mu(y - x) + p_\mu(z - y)) \quad \forall \mu \in \Lambda.$$

Since  $p_\mu(z - x) \leq p_\mu(y - x) + p_\mu(z - y)$ ,

$$f(z) - f(x) \leq -\gamma_\mu(p_\mu(y - x) + p_\mu(z - y)) \leq -\gamma_\mu p_\mu(z - x) \quad \forall \mu \in \Lambda.$$

Hence  $z \leq x$ . In order to apply the above theorem by replacing  $f = \psi$  in [1],

1. Since  $f$  is sequentially lower semi-continuous,  $S(x)$  is sequentially closed for each  $x \in X$ ;
2. If  $x \leq y$  and  $x \neq y$ , then there exists  $\mu_0 \in \Lambda$  such that  $p_{\mu_0}(y - x) > 0$  because  $X$  is Hausdorff. Hence

$$f(y) - f(x) \leq -\gamma_{\mu_0} p_{\mu_0}(y - x) < 0, \quad f(y) < f(x);$$

3. for any increasing sequence  $x_n$

$$x_{n+1} \geq x_n, \quad f(x_{n+1}) - f(x_n) \leq -\gamma_\mu p_\mu(x_{n+1} - x_n) \leq 0 \quad \forall \mu \in \Lambda.$$

Therefore  $\{f(x_n)\}$  is decreasing and bounded below. So  $\{f(x_n)\}$  converges. Hence for any  $\mu \in \Lambda$ ,  $x_n$  is  $p_\mu$ -Cauchy. Since  $X$  is sequentially complete,  $\{x_n\}$  converges in  $X$ . Hence  $\{x_n\}$  is compact and relatively compact.

From the above principle for any  $x_0 \in X$  there exists  $x^* \in S(x_0)$  such that  $x^*$  is maximal. Since  $x^* \in S(x_0)$ ,

$$f(x^*) + \gamma_\lambda p_\lambda(x^* - x_0) \leq f(x_0)$$

for all  $\lambda \in \Lambda$ . Since  $x^*$  is maximal, for all  $x \neq x^* \in X$  there exists  $\mu \in \Lambda$  such that

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*).$$

□

**THEOREM 2.2.** [4] *Let  $X$  be a Hausdorff locally convex topological space that is sequentially complete. Let  $f : X \rightarrow (\infty, \infty]$  be an extended-valued proper and sequentially lower semi-continuous function, bounded below. Let  $\{p_\lambda\}_{\lambda \in \Lambda}$  be a base of continuous seminorms generating the topology on  $X$  and  $\{\gamma_\lambda\}_{\lambda \in \Lambda}$  a family of positive numbers. Then for every  $x_0 \in \text{dom } f$  there exists  $x^* \in X$  such that*

$$f(x^*) + \gamma_\lambda p_\lambda(x^* - x_0) \leq f(x_0)$$

for all  $\lambda \in \Lambda$ , and for all  $x \neq x^*$  there exists  $\mu \in \Lambda$  such that

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*).$$

*Proof.* For fixed  $x_0 \in \text{dom}f$  let us give the same ordering on the following sequentially closed subset  $C$  of  $X$

$$C = \{x \in X \mid f(x) + \gamma_\mu p_\mu(x - x_0) \leq f(x_0) \forall \gamma \in \Lambda\}.$$

That is,

$$x \leq y (\in C) \text{ iff } f(y) - f(x) \leq -\gamma_\lambda p_\gamma(y - x) \forall \gamma \in \Lambda.$$

Indeed it is an ordering on  $C$  that satisfies the three conditions of the above general principle. So  $C$  has a maximal  $x^*$ . We must prove for all  $x \neq x^*$  there exists  $\mu \in \Lambda$  such that

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*).$$

If  $x \in C (\neq x^*)$ , it is not  $x^* \leq x$ , so the conclusion holds. If  $x \notin C (\neq x^*)$  and  $x \in \text{dom}f$ ,

$$f(x) + \gamma_\mu p_\mu(x - x_0) > f(x_0)$$

for some  $\mu \in \Lambda$ . And since  $x^* \in C$ ,

$$f(x^*) + \gamma_\mu p_\mu(x^* - x_0) \leq f(x_0).$$

It follows that

$$f(x^*) + \gamma_\mu p_\mu(x^* - x_0) \leq f(x_0) < f(x) + \gamma_\mu p_\mu(x - x_0).$$

Since  $p_\mu(x - x_0) \leq p_\mu(x^* - x_0) + p_\mu(x - x^*)$ ,

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*)$$

for some  $\mu \in \Lambda$ . If  $x \notin \text{dom}f$ , clearly the inequality holds.  $\square$

**THEOREM 2.3.** [4] *Let  $X$  be a Hausdorff locally convex topological space that is sequentially complete. Let  $f : X \rightarrow (\infty, \infty]$  be an extended-valued proper and sequentially lower semi-continuous function, bounded below. Let  $S \subset X$  be a sequentially closed, bounded and convex set such that  $0 \in S$ . Then for every  $r > 0$ ,  $x_0 \in \text{dom}f$  there exists  $x^* \in X$  such that*

$$f(x^*) + r\mu_S(x^* - x_0) \leq f(x_0),$$

and for all  $x \in X, x \neq x^*$  we have

$$f(x^*) < f(x) + r\mu_S(x - x^*).$$

*Proof.* Let us fix  $x_0 \in \text{dom}f, r > 0$ . From the sequentially lower semi-continuity of  $f, \mu_S$  in Lemma 1.1(1), the following subset  $X' \subset X$  is sequentially closed.

$$X' = \{x \in X \mid f(x) + r\mu_S(x - x_0) \leq f(x_0)\}$$

Let  $\{p_\lambda\}_{\lambda \in \Lambda}$  be a base of continuous seminorms generating the topology on  $X$ . Then by Lemma 1.2 there exists  $\{\alpha_\lambda\}_{\lambda \in \Lambda}$  a family of positive numbers such that

$$\frac{1}{\alpha_\lambda} p_\lambda(x) \leq \mu_S(x), x \in \text{dom } \mu_S.$$

Let us  $\gamma_\lambda = \frac{r}{\alpha_\lambda}$  for all  $\lambda \in \Lambda$ . We give the following order structure on  $X'$ . That is,

$$x \leq y (\in X') \text{ iff } f(y) - f(x) \leq -\gamma_\lambda p_\lambda(y - x) \quad \forall \lambda \in \Lambda.$$

Indeed it is an ordering on  $C$  that satisfies the three conditions of the above general principle. So  $X'$  has a maximal  $x^*$ . We prove that for all  $x \in X (\neq x^*)$

$$f(x^*) < f(x) + \mu_S(x - x^*).$$

If  $x - x^* \notin \text{dom } \mu_S$ , this inequality holds. Hence we may assume that  $x - x^* \in \text{dom } \mu_S$ . Since  $x^* \in X'$ ,  $x^* - x_0 \in \text{dom } \mu_S$ . By Lemma 1.1(2)  $x - x_0 \in \text{dom } \mu_S$ . If  $x \in X' (\neq x^*)$ , then  $x^* \not\leq x$  and there exists  $\mu \in \Lambda$  such that

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*).$$

Since  $x - x^* \in \text{dom } \mu_S$ ,  $\gamma_\mu p_\mu(x - x^*) \leq r \mu_S(x - x^*)$  and

$$f(x^*) < f(x) + \gamma_\mu p_\mu(x - x^*) \leq f(x) + r \mu_S(x - x^*).$$

If  $x \notin X' (\neq x^*)$ , then

$$f(x) + r \mu_S(x - x_0) > f(x_0)$$

And since  $x^* \in X'$ ,

$$f(x^*) + r \mu_S(x^* - x_0) \leq f(x_0).$$

Since  $x - x_0 \in \text{dom } \mu_S$  and  $(x - x_0) - (x^* - x_0) = x - x^* \in \text{dom } \mu_S$ , by Lemma 1.1(3) it follows that

$$\mu_S(x - x_0) - \mu_S(x^* - x_0) \leq \mu_S(x - x^*).$$

From

$$f(x^*) < f(x) + r(\mu_S(x - x_0) - \mu_S(x^* - x_0)) \leq f(x) + \mu_S(x - x^*).$$

□

### References

- [1] H.Brézis and F.E.Browder, *A general principles on odered sets in nonlinear functional analysis*, Adv. in Math. 21(1976), 355-364.
- [2] J.Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc.215(1976), 241-251.
- [3] I.Ekeland, *Nonconvex minimization problems*, Bull.Amer.Math.Soc., 1(3)(1979), 443-474.
- [4] A.H.Hamel, *Phelps' lemma, Daneš drop theorem and Ekeland's principle in locally convex space* , Proc. Amer. Math. Soc. 131(2003), 3025-3038.

Department of Mathematics  
Kangwon National University  
Chunchon, 200-701, Korea  
*E-mail*: jongan@kangwon.ac.kr