

CONFORMAL CHANGE OF THE VECTOR S_ω IN 7-DIMENSIONAL g -UFT

CHUNG HYUN CHO

ABSTRACT. We investigate change of the vector S_ω induced by the conformal change in 7-dimensional g -unified field theory. These topics will be studied for the second class with the first category in 7-dimensional case.

1. Introduction

The conformal change in a generalized 4-dimensional Riemannian space conneted by an Einstein's connection was primarily studied by HLAVATÝ([6],1957). CHUNG([4],1968) also investigated the same topic in 4-dimensional $*g$ -unified field theory.

The Einstein's connection induced by the conformal change of the tensor $S_{\omega\mu}{}^\nu$ for the second class with the first category in 7-dimensional g -UFT were investigated by CHO([2],2001).

In the present paper, we investigate change of the vector S_ω induced by the conformal change in 7-dimensional g -unified field theory. These topics will be studied for the second class with the first category in 7-dimensional case.

2. Preliminaries

This chapter is a brief collection of basic concepts, notations, theorems, and results needed in our further considerations. They may be referred to CHUNG([3]), CHO([1],1992; [2],2001).

Received July 12, 2005.

2000 Mathematics Subject Classification: 53A45, 83E99.

Key words and phrases: Einstein connection, conformal change.

2.1. n -dimensional g -unified field theory. The n -dimensional g -unified field theory (n - g -UFT hereafter) was originally suggested by HLAVATÝ([6],1957) and systematically introduced by CHUNG([5],1963).

Let $X_n(n \geq 2)$ be an n -dimensional generalized Riemannian manifold, referred to a real coordinate system x^ν obeying coordinate transformations $x^\nu \rightarrow x^{\nu'}$, for which

$$(2.1) \quad \text{Det}\left(\left(\frac{\partial x}{\partial x'}\right)\right) \neq 0.$$

In the usual Einstein's n -dimensional unified field theory, the manifold X_n is endowed with a general real nonsymmetric tensor $g_{\lambda\mu}$ which may be split into its symmetric part $h_{\lambda\mu}$ and skew-symmetric part $k_{\lambda\mu}$:

$$(2.2) \quad g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu}$$

where

$$(2.3) \quad \text{Det}((g_{\lambda\mu})) \neq 0 \quad \text{Det}((h_{\lambda\mu})) \neq 0.$$

Therefore we may define a unique tensor $h^{\lambda\nu} = h^{\nu\lambda}$ by

$$(2.4) \quad h_{\lambda\mu} h^{\lambda\nu} = \delta_\mu^\nu.$$

In our n - g -UFT, the tensors $h_{\lambda\mu}$ and $h^{\lambda\nu}$ will serve for raising and/or lowering indices of the tensors in X_n in the usual manner.

The manifold X_n is connected by a general real connection $\Gamma_{\omega\mu}^\nu$ with the following transformation rule :

$$(2.5) \quad \Gamma_{\omega'\mu'}^{\nu'} = \frac{\partial x^{\nu'}}{\partial x^\alpha} \left(\frac{\partial x^\beta}{\partial x^{\omega'}} \cdot \frac{\partial x^\gamma}{\partial x^{\mu'}} \Gamma_{\beta\gamma}^\alpha + \frac{\partial^2 x^\alpha}{\partial x^{\omega'} \partial x^{\mu'}} \right)$$

and satisfies the system of Einstein's equations

$$(2.6) \quad D_\omega g_{\lambda\mu} = 2S_{\omega\mu}^\alpha g_{\lambda\alpha}$$

where D_ω denotes the covariant derivative with respect to $\Gamma_{\lambda\mu}^\nu$ and

$$(2.7) \quad S_{\lambda\mu}^\nu = \Gamma_{[\lambda\mu]}^\nu$$

is the *torsion tensor* of $\Gamma_{\lambda\mu}^\nu$. The connection $\Gamma_{\lambda\mu}^\nu$ satisfying (2.6) is called the *Einstein's connection*.

In our further considerations, the following scalars, tensors, abbreviations, and notations for $p = 0, 1, 2, \dots$ are frequently used :

$$\mathbf{g} = \text{Det}((g_{\lambda\mu})) \neq 0, \quad \mathbf{h} = \text{Det}((h_{\lambda\mu})) \neq 0, \quad (2.8a)$$

$$\mathbf{t} = \text{Det}((k_{\lambda\mu})),$$

$$g = \frac{\mathbf{g}}{\mathbf{h}}, \quad k = \frac{\mathbf{t}}{\mathbf{h}}, \quad (2.8b)$$

$$K_p = k_{[\alpha_1}^{\alpha^1} \dots k_{\alpha_p]}^{\alpha^p}, \quad (p = 0, 1, 2, \dots) \quad (2.8c)$$

$${}^{(0)}k_\lambda^\nu = \delta_\lambda^\nu, \quad {}^{(1)}k_\lambda^\nu = k_\lambda^\nu, \quad {}^{(p)}k_\lambda^\alpha = {}^{(p-1)}k_\lambda^\alpha k_\alpha^\nu, \quad (2.8d)$$

$$K_{\omega\mu\nu} = \nabla_\nu k_{\omega\mu} + \nabla_\omega k_{\nu\mu} + \nabla_\mu k_{\omega\nu}, \quad (2.8e)$$

$$\sigma = \begin{cases} 1 & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases} \quad (2.8f)$$

where ∇_ω is the symbolic vector of the covariant derivative with respect to the Christoffel symbols $\{\lambda_\mu^\nu\}$ defined by $h_{\lambda\mu}$. The scalars and vectors introduced in (2.8) satisfy

$$K_0 = 1; K_n = k \quad \text{if } n \text{ is even}; \quad K_p = 0 \quad \text{if } p \text{ is odd}, \quad (2.9a)$$

$$g = 1 + K_2 + \dots + K_{n-\sigma}, \quad (2.9b)$$

$${}^{(p)}k_{\lambda\mu} = (-1)^{p(p)} k_{\mu\lambda}, \quad {}^{(p)}k^{\lambda\mu} = (-1)^{p(p)} k^{\nu\lambda}. \quad (2.9c)$$

Furthermore, we also use the following useful abbreviations, denoting an arbitrary tensor $T_{\omega\mu\nu}$, skew-symmetric in the first two indices, by T :

$${}^{pqr}T = {}^{pqr}T_{\omega\mu\nu} = T_{\alpha\beta\gamma} {}^{(p)}k_\omega^{\alpha(q)} k_\mu^{\beta(r)} k_\nu^{\gamma}, \quad (2.10a)$$

$$T = T_{\omega\mu\nu} = {}^{000}T, \quad (2.10b)$$

$$2 {}^{pqr}T_{\omega[\lambda\mu]} = {}^{pqr}T_{\omega\lambda\mu} - {}^{pqr}T_{\omega\mu\lambda}, \quad (2.10c)$$

$$2 {}^{(pq)r}T_{\omega\lambda\mu} = {}^{pqr}T_{\omega\lambda\mu} + {}^{qpr}T_{\omega\lambda\mu}. \quad (2.10d)$$

We then have

$${}^{pqr}T_{\omega\lambda\mu} = -{}^{qpr}T_{\lambda\omega\mu}. \quad (2.11)$$

If the system (2.6) admits $\Gamma_{\lambda\mu}^\nu$, using the above abbreviations it was shown that the connection is of the form

$$\Gamma_{\omega\mu}^\nu = \{\nu_{\omega\mu}\} + S_{\omega\mu}^\nu + U_{\omega\mu}^\nu \quad (2.12)$$

where

$$U_{\nu\omega\mu} = 2 S_{\nu(\omega\mu)}^{001}. \quad (2.13)$$

The above two relations show that our problem of determining $\Gamma_{\omega\mu}^{\nu}$ in terms of $g_{\lambda\mu}$ is reduced to that of studying the tensor $S_{\omega\mu}^{\nu}$. On the other hand, it has also been shown that the tensor $S_{\omega\mu}^{\nu}$ satisfies

$$S = B - 3 S^{(110)} \quad (2.14)$$

where

$$2B_{\omega\mu\nu} = K_{\omega\mu\nu} + 3K_{\alpha[\mu\beta}k_{\omega]}^{\alpha}k_{\nu}^{\beta}. \quad (2.15)$$

DEFINITION 2.1. The vector S_{ω} defined by

$$S_{\omega} = S_{\omega\alpha}^{\alpha}. \quad (2.16)$$

2.2. Some results for the second class with the first category in 7- g -UFT. In this section, we introduce some results of 7- g -UFT without proof, which are needed in our subsequent considerations.

They may be referred to CHO([1],1992).

DEFINITION 2.2. In 7- g -UFT, the tensor $g_{\lambda\mu}(k_{\lambda\mu})$ is said to be the second class, if $K_2 \neq 0$, $K_4 = 0 = K_6$ with the first category.

THEOREM 2.3 (MAIN RECURRENCE RELATIONS). *For the second class in 7-UFT, the following recurrence relation hold*

$${}^{(p+3)}k_{\lambda}^{\nu} = -K_2^{(p+1)}k_{\lambda}^{\nu}, \quad (p = 0, 1, 2, \dots). \quad (2.17)$$

THEOREM 2.4 (FOR THE SECOND CLASS IN 7- g -UFT). *A necessary and sufficient condition for the existence and uniqueness of the solution of (2.5) is*

$$1 - (K_2)^2 \neq 0. \quad (2.18)$$

If the condition (2.18) is satisfied, the unique solution of (2.14) is given by

$$(1 - K_2^2)(S - B) = -2 B^{(10)1} + (K_2 - 1) B^{110} + 2 B^{(20)2} + 2 B^{112}. \quad (2.19)$$

3. Conformal change of the 7-dimensional vector S_ω for the second class

In this final chapter we investigate the change $S_\omega \rightarrow \bar{S}_\omega$ of the vector induced by the conformal change of the tensor $g_{\lambda\mu}$, using the recurrence relations and theorems introduced in the preceding chapter.

We say that X_n and \bar{X}_n are conformal if and only if

$$\bar{g}_{\lambda\mu}(x) = e^{\Omega} g_{\lambda\mu}(x) \tag{3.1}$$

where $\Omega = \Omega(x)$ is an at least twice differentiable function. This conformal change enforces a change of the vector S_ω . An explicit representation of the change of 5-dimensional vector S_ω for the second class will be exhibited in this chapter.

AGREEMENT 3.1. Throughout this section, we agree that, if T is a function of $g_{\lambda\mu}$, then we denote \bar{T} the same function of $\bar{g}_{\lambda\mu}$. In particular, if T is a tensor, so is \bar{T} . Furthermore, the indices of $T(\bar{T})$ will be raised and/or lowered by means of $h^{\lambda\nu}(\bar{h}^{\lambda\nu})$ and/or $h_{\lambda\nu}(\bar{h}_{\lambda\nu})$.

The results in the following theorems are needed in our further considerations. They may be referred to CHO([1],1992, [2],2001).

THEOREM 3.2. *In n - g -UFT, the conformal change (3.1) induces the following changes:*

$${}^{(p)}\bar{k}_{\lambda\mu} = e^{\Omega(p)} k_{\lambda\mu}, \quad {}^{(p)}\bar{k}_\lambda = {}^{(p)}k_\lambda^\nu, \quad {}^{(p)}\bar{k}^{\lambda\mu} = e^{-\Omega(p)} k^{\lambda\mu}, \tag{3.2a}$$

$$\bar{g} = g, \quad \bar{K}_p = K_p, \quad (p = 1, 2, \dots). \tag{3.2b}$$

THEOREM 3.3. *(For the second classes with the first category in 7- g -UFT). The change of the tensor $B_{\omega\mu\nu}$ induced by the conformal change (3.1) may be given by*

$$\begin{aligned} \bar{B}_{\omega\mu\nu} &= e^{\Omega}(B_{\omega\mu\nu} + k_{\nu[\omega}\Omega_{\mu]} - k_{\omega\mu}\Omega_\nu \\ &\quad - h_{\nu[\omega}k_{\mu]}^\delta\Omega_\delta + 2^{(2)}k_{\nu[\omega}k_{\mu]}^\delta\Omega_\delta + k_{\omega\mu}{}^{(2)}k_\nu^\delta\Omega_\delta). \end{aligned} \tag{3.3}$$

Now, we are ready to derive representations of the changes $S_\omega \rightarrow \bar{S}_\omega$ in 7- g -UFT for the second class with the first category induced by the conformal change (3.1).

THEOREM 3.4. *The conformal change (3.1) induces the following change:*

$$\begin{aligned} \overline{{}^{(10)1} B}_{\omega\mu\nu} &= e^{\Omega} 2 \left[\overline{{}^{(10)1} B}_{\omega\mu\nu} + (-2) k_{\nu[\omega} k_{\mu]}^{\delta} \right. \\ &\quad \left. + 2 k_{\nu[\omega} k_{\mu]}^{\delta} \Omega_{\delta} + k_{\omega\mu} (2) k_{\nu}^{\delta} \Omega_{\delta} - (3) k_{\nu[\omega} \Omega_{\mu]} \right]. \end{aligned} \quad (3.4)$$

THEOREM 3.5. *The conformal change (3.1) may be represented by*

$$\begin{aligned} \overline{{}^{ppq} B}_{\omega\mu\nu} &= e^{\Omega} \left[\overline{{}^{ppq} B}_{\omega\mu\nu} + (-1)^p \{ 2^{(p+q+2)} k_{\nu[\omega} (p+1) k_{\mu]}^{\delta} \right. \\ &\quad \left. + (2p+1) k_{\omega\mu} (2+q) k_{\nu}^{\delta} - (2p+1) k_{\omega\mu} (q) k_{\nu}^{\delta} \right. \\ &\quad \left. + (p+q+1) k_{\nu[\omega} (p) k_{\mu]}^{\delta} - (p+q) k_{\nu[\omega} (p+1) k_{\mu]}^{\delta} \} \Omega_{\delta} \right]. \\ &\quad \left(\begin{array}{l} p = 0, 1, 2, 3, 4, \dots \\ q = 0, 1, 2, 3, 4, \dots \end{array} \right) \end{aligned} \quad (3.5)$$

THEOREM 3.6. *The change $S_{\omega\mu}{}^{\nu} \rightarrow \overline{S}_{\omega\mu}{}^{\nu}$ induced by conformal change (3.1) may be represented by*

$$\begin{aligned} \overline{S}_{\omega\mu}{}^{\nu} &= S_{\omega\mu}{}^{\nu} + 1 - h^{\nu}{}_{[\omega} k_{\mu]}^{\delta} \Omega_{\delta} \\ &\quad + (K_2 - 1) k_{\omega\mu} \Omega^{\nu} + (1 - K_2) k_{\omega\mu} (2) k^{\nu\delta} \Omega_{\delta} \\ &\quad + \frac{1}{K_2^2 - 1} [(-1 + K_2) k^{\nu}{}_{[\omega} \Omega_{\mu]}] \\ &\quad + (-1 + 2K_2 + K_2^2) (2) k^{\nu}{}_{[\omega} k_{\mu]}^{\delta} \Omega_{\delta} \\ &\quad + (K_2 + K_2^2 - 2K_2^3) k^{\nu}{}_{[\omega} (2) k_{\mu]}^{\delta} \Omega_{\delta}]. \end{aligned} \quad (3.6)$$

THEOREM 3.7. *The change $S_{\omega} \rightarrow \overline{S}_{\omega}$ induced by conformal change (3.1) may be represented by*

$$\begin{aligned} \overline{S}_{\omega} &= S_{\omega} + 1 + \frac{1}{C} [(3 - 3K_2 - 7K_2^2) \\ &\quad + 5K_2^3 - 2K_2^4) k_{\omega}^{\delta} \Omega_{\delta} \\ &\quad + (1 - 2K_2 - K_2^2) (2) k_{\alpha}^{\alpha} k_{\omega}^{\delta} \Omega_{\delta}] \end{aligned} \quad (3.7)$$

where $C = 2(K_2^2 - 1)$.

Proof. In virtue of Definition 2.1 and Agreement (3.1), we have

$$\bar{S}_\omega = \bar{S}_{\omega\alpha}{}^\alpha \quad (3.8)$$

The relation(3.7) follows by substituting (3.2), (3.3), (2.10), Definition (2.2) into Theorem 3.6.

References

- [1] C.H. Cho, Conformal change of the connection in 3- and 5-dimensional $*g^{\lambda\nu}$ -unified Field Theory, *BIBS, Inha Univ.* **13** (1992), 11–19.
- [2] C.H. Cho, Conformal change of the tensor $S_{\omega\mu}{}^\nu$ in 7-dimensional g -UFT, *Bull.Korean Math.Soc.***38** (2001),No.1, 197–203.
- [3] K.T. Chung, Three- and Five-dimensional considerations of the geometry of Einsteins's $*g$ -unified field theory, *International Journal of Theoretical Physics* **27(9)** (1988), 1105–1136.
- [4] K.T. Chung, Conformal change in Einstein's $*g^{\lambda\nu}$ -unified field theory, *Nuove Cimento* (**X**)**58B** (1968), 201–212.
- [5] K.T. Chung, Einstein's connection in terms of $*g^{\lambda\nu}$, *Nuove Cimento* (**X**) **27** (1963), 1297–1324.
- [6] V. HLAVATÝ, Geometry of Einstein's unified field theory, *P. Noordhoff Ltd* (1957).

Department of Mathematics
Inha University
Incheon, 402–751, Korea
E-mail: chcho@inha.ac.kr