

외상성 뇌손상 후 이차적으로 발생한 성인 주의력결핍
과잉행동장애에서 OROS Methylphenidate의 치료효과 :
증례 및 고찰*

임 명 호**† · 이 우 철**

OROS Methylphenidate Treatment of Secondary Adult ADHD after
Traumatic Brain Injury*

Myung Ho Lim, M.D., **† Woo Chul Lee, M.D., **

ABSTRACT

The incidence of the Attention Deficit Hyperactivity Disorder secondary to the traumatic brain injury, such as traffic accidents, is increasing ; the variety of the treatment modality is also increasing. This case was studied to see if OROS Methylphenidate(Concerta), which is one of the most commonly used medication in Attention Deficit Hyperactivity Disorder patients, not only improves the patient's attention, but also their impulsivity, hyperactivity and aggression. According to the case result, the medication showed an improvement of the impulsivity, aggression, and attention in the secondary Attention Deficit Hyperactivity Disorder patients after the traumatic brain injury.

KEY WORDS : OROS methylphenidate(Concerta) · Traumatic brain injury(TBI) · Adult attention deficit hyperactive disorder(ADHD).

서 론

가 가

가

가

2004

Department of Psychiatry, College of Medicine, Dankook
University, Cheonan, Korea

†교신저자 : , 330 - 715 16
) (041) 550 - 3945,) (041) 561 - 3007
E - mail) paperose@kornet.net

가

Fann 2) 939

1 49% , 증 례
34%가, 18%가
6 가 27
4.0
(95% 2.4~6.8) 가
2.8(95% 2.4~6.8) 가
1 가 가
가
2003 , 9 8 가
가
, ,
Glasgow coma scale 7
가 , 7 45
. 3) . 11 21
5
“
, , , , ”
. 3) , 가 “ 가
”
가 가 . 2003 12
가 , 4)
aman- ris-
tadine methylphenidate peridone 4mg, orfil 1200mg . 2004 1
, 5) 6)7)
가 가
OROS Methylpheni-
date
OROS Methylphenidate

가

Wender - Utah ⁹⁾ 8

7

5 가 가 (Adult ADHD Self - Report Scale - v.11 ; ASRS)¹⁰⁾ 33 , MINI PLUS 가 11 , Buss & Dukee ¹¹⁾ 46 . Continuous Performance Test Computerized Attention Deficit System¹²⁾ 52 , 40 , risperidone 4mg, 71 , 65 sodium valproate 1000mg, clonazepam 0.5mg . Compute- rized Attention Deficit System 2004 40~60 7 가 , 60 가 , 가 (WAIS) 80, 78, 87 . Beck ¹³⁾ Hamilton 가 , ¹⁴⁾ Beck ¹⁵⁾ Spielberg , ¹⁶⁾ Maudsley ¹⁷⁾ , ¹⁸⁾ ¹⁹⁾ 가 , ²⁰⁾ 가

OROS Methylphenidate 54mg . 2 가 가 가 22 , MINI PLUS 가 8 . Buss & Dukee 38 8 . Computerized Attention Deficit System 41 , 44 , 52, 43

토 론

1 washout 가 Brown ²¹⁾ 31 . DSM - ⁸⁾ 2 3

가

가

OROS Methylphenidate DSM -

5)

, NMDA

amantadine

가 , Barthel index Brief Psychiatric Rating Scale 6)

가

Methylphenidate 5

22) 7) 18

, DSM - Methylphenidate 20mg 1 superLab

가 pro 2.0

가

27

OROS Methylphenidate

2~4%

23)

가

24)25)

DSM -

가 , 27) OROS Methylpheni-

Wender - Utah date 27

가 가 , MINI PLUS

가 , Computerized Attention Deficit

System

가

Methylphenidate

Gering 26) 99

1

Mooney 28)

Whyte 29)

0.2 24

0.04 , 0.3mg/kg/dose

가

Methylphenidate

가

가

30)

3)

가

Faraone 31)

Methylphenidate

6

effect

size 0.9

0.7 effect size

1.05mg/kg

1.3 effect size

54mg(1.0mg/kg)

OROS Methylphenidate Methylphenidate

12

Methylphenidate

32)

33)

OROS Methylphenidate

2

가

2

가

Methylphenidate

가

가

34)

요 약

가

가

가

OROS Methylphenidate

중심 단어 :

참고문헌

1. 박원명, 김 원, 전태연. 정신과에서의 외상성 뇌 손상. 진단과 치료 2004;24:740-744.
2. Fann JR, Burington B, Leonetti A, Jaffe K, Katon WJ, Thompson RS. Psychiatric illness following traumatic brain injury in an adult health maintenance organization population. Arch Gen Psychiatry 2004;61:53-61.
3. Jorge RE. Neuropsychiatric consequences of traumatic brain injury: a review of recent finding. Current Opinion Psychiatry 2005;18:289-299.
4. Zafonte RD, Lexell J, Cullen N. Possible applications for dopaminergic agents following traumatic brain injury: part 2. J Head Trauma Rehabil 2001;16:112-116.
5. 정한용, 이소영, 김양래. 외상성 뇌손상 환자에서 Amantadine의 치료적 효과: 2중례 및 고찰. 생물정신의학 2001;8:156-161.
6. 강경주, 윤서라, 조길호, 한승상, 노성만. 뇌손상 환자에서 Methylphenidate의 인지기능에 대한 효과. 대한재활의학회지 1998;22:230-235.
7. 김연희, 나승용, 신용일, 서정환. 뇌손상 후 인지기능 장애에 대한 메틸페니데이트의 효과: 이중맹검 위약대조 연구. 대한재활의학회지 2000;24:611-617.
8. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th ed. Washington, DC: American Psychiatric Association;1994.
9. Ward MF, Wender PH, Reimherr FW. The Wender Utah Rating Scale: an aid in the retrospective diagnosis of childhood attention deficit hyperactivity disorder. Am J Psychiatry 1993;150:885-890.
10. Adler LA, Kessler RC, Spencer T. Adult ADHD Self-report Scale v1.1 (ASRS-v1.1) symptom checklist. New York, NY: World Health Organization;2003. Available from URL: <http://www.med.nyu.edu/Psych/training/adhd.html>
11. 고영인. 대학생의 분노표현 양식과 우울 및 공격성과의 관계(박사학위). 부산: 부산대학교;1994.
12. 신민섭, 조성준, 전선영, 홍강의. 전산화된 주의력장애 진단시스템의 개발 및 표준화연구. 소아청소년정신의학 2000;11:91-99.
13. 이영호, 송중용. BDI, SDS, MMPI-D 척도의 신뢰도 및 타당도에 대한 연구. 한국심리학회지 임상 1991; 15:98-113.
14. 김 연. 우울증의 귀인모형에 관한 연구(박사학위). 서울: 고려대학교;1985.

15. 서수균. 자기집중적 주의: 실제 자기개념 및 자기안 내자 선호성을 고려한 자기 불일치와 우울 및 불안의 관계(석사학위). 서울: 서울대학교;1996.
16. 김정택. 특성불안과 사회성과의 관계: Spielberger의 STAI를 중심으로(석사학위). 서울: 고려대학교;1978.
17. 조대경. 강박행동의 객관적 측정에 관한 연구: Maudsley 강박행동목록의 표준화. 학생연구 1995;20:78-90.
18. 강영수. 외상후 스트레스장애 진단을 위한 미시시 피 척도와 MMPI-PTSD 척도의 비교(박사학위). 전주: 전북대학교;1997.
19. 김영아, 오경자. 양극성 장애의 고위험군으로서의 경 조성 집단의 심리적인 특성. 한국심리학회지 임상 1996;15:103-115.
20. 이훈진, 원호택. 편집성 척도(Paranoia Scale)의 신뢰도, 타당도 연구. 한국임상심리학회지 임상 1995;14: 83-94.
21. Brown G, Chadwick O, Shaffer D, Rutter M, Traub M. A prospective study of children with head injuries, III: psychiatric sequelae. Psychol Med 1981;11:63-78.
22. Wender PH. Attention deficit hyperactivity disorder in adults: a wide view of a widespread condition. Psychiatric Annals 1998;27:8:556-562.
23. Barkley RA, Fischer M, Smallish L, Fletcher K. The persistence of attention-deficit/hyperactivity disorder into young adulthood as a function of reporting source and definition of disorder. J Abnorm Psychol 2002;111:279-289.
24. Adler LA, Chua HC. Management of ADHD in Adults. J Clin Psychiatry 2002;63:29-35.
25. Murphy KR, Adler LA. Assessing attention-deficit/hyperactivity disorder in adults: focus on rating scales. J Clin Psychiatry 2004;65 (Suppl 3) :12-17.
26. Gerring JP, Brady KD, Chen A, Vasa R, Grados M, Bandeen-Roche KJ, et al. Premorbid prevalence of ADHD and development of secondary ADHD after closed head injury. J Am Acad Child Adolesc Psychiatry 1998;6:647-654.
27. Greenhill LL, Pliszka S, Dulcan MK, Bernet W, Arnold V, Beitchman J, et al. Summary of the practice parameter for the use of stimulant medications in the treatment of children, adolescents, and adults. J Am Acad Child Adolesc Psychiatry 2001;40:1352-1355.
28. Mooney GF, Haas LJ. Effect of methylphenidate on brain injury-related anger. Arch Phys Med Rehabil 1993; 74:153-160.
29. Whyte J, Hart T, Vaccaro M, Grieb-Neff P, Risser A, Polansky M, et al. Effects of methylphenidate on attention deficits after traumatic brain injury: a multidimensional, randomized, controlled trial. Am J Phys Med Rehabil 2004;83:401-420.
30. Speech TJ, Rao SM, Osmon DC, Sperry LT. A double-blind controlled study of methylphenidate treatment in closed head injury. Brain Inj 1993;7:333-338.
31. Faraone SV, Spencer T, Alcardi M, Pagano C, Biederman J. Metaanalysis of the efficacy of methylphenidate for treating adult attentiondeficit/hyperactivity disorder. J Clin Psychopharmacol 2004;24:24-29.
32. Pelham WE, Gnagy EM, Burrows-Maclean L, Williams A, Fabiano GA, Morrisey SM, et al. Once-a-day Concerta methylphenidate versus three-times-daily methylphenidate in laboratory and natural settings. Pediatrics 2001; 107:E105.
33. Wilens TE, Hammerness PG, Biederman J, Kwon A, Spencer TJ, Clark S, et al. Blood pressure changes associated with medication treatment of adults with attention-deficit/hyperactivity disorder. J Clin Psychiatry 2005;66:253-259.
34. The MTA Cooperative Group. Multimodal Treatment Study of Children with ADHD. A 14-month randomized clinical trial of treatment strategies for Attention-deficit/hyperactivity disorder. Arch Gen Psychiatry 1999;56: 1073-1086.