

New Hydroxamic Acid Derivative Prevents Sq-OOH Induced Wrinkle

Ho Sik Rho,^{*} Heung Soo Baek, Byoung-Seok Lee, Ji Hyun Kim, Duck Hee Kim, and Ih Seop Chang

R & D Center, AmorePacific Corporation, Yongin, Gyeonggi-do 446-729, Korea. *E-mail: thiocarbon@freechal.com

Received August 12, 2005

Key Words : Hydroxamic acid, Lipid peroxidation, Sq-OOH, Epidermal aging

Lipid peroxidation has been suggested to play a key role in damaging biological processes resulting from excessive exposure to UV light.¹ Aging of skin has been associated with increased peroxidation of skin lipid.² In particular, squalene, which is a main component of skin surface polyunsaturated lipids, is easily peroxidized.³ Squalene-mono-hydroperoxide (Sq-OOH), the initial product of peroxidized squalene, is produced at the human skin by natural exposure to sunlight during day time activities.

Recently, a few researchers reported on the damages of hairless mice skin by the topical application of Sq-OOH.⁴ Repeated application of Sq-OOH to the hairless mice induced its the skin wrinkles with clear and fine appearance.^{4c} Histologically were seen, more keratinocyte hypertrophy with Sq-OOH treated skin than with chronic UVB irradiated group in epidermis. In dermis, characteristics, found in UVB irradiated skin, such as the loss of polarity, fractured collagen bundle and elastosis, were not observed severely. The alteration of epidermal layer might be the main factor for the formation of wrinkle caused by Sq-OOH. From the previous works,⁴ Sq-OOH is considered as an important source of damage in the early stage of aging process to result in epidermal wrinkle. Thus, it is very important to minimize such damage that would be induced mainly by Sq-OOH against healthy skin condition. Therefore, we tried to develop new anti-aging materials for protecting skin damage by Sq-OOH. Hydroxamic acids, the naturally occurring products exhibit low toxicities in general, and are of interest for many therapeutic applications.⁵ Recently, hydroxamic acids and derivatives were reported largely as effective antibacterial and antifungal agents.⁶ Their activities originate from their ability to coordinate on metal ions.⁷ That is, hydroxamic acid group is regarded as an one electron donor. Hydroxamic acid moiety has also drawn much attention as a reactive oxygen species (ROS) scavenger.⁸

In this study, a new skin protecting compound, (4-(*N*-hydroxycarbonyl)phenyl)-*N*-adamantanylecarboxamide (**1**)





Figure 1. Structure of hydroxamic acid derivative.

is synthesized to show inhibitory activities of lipid peroxidation both *in vitro* and *in vivo*. The targeted compound **1** is a hydroxamic acid derivative which possesses an amide linker between hydroxamic acid and adamantane moiety. Adamantane group was adopted as a hydrophobic moiety to optimize antioxidant activity of hydroxamic acid in animal test. Compound **1** is believed to be adequate in skin permeation because of its balance in hydrophilic and hydrophobic character.

Experimental Section

Synthesis. Mono-methyl terephthalate **2** was refluxed in thionyl chloride to afford an acid chloride. This compound was reacted immediately with 1-adamantamine HCl in pyridine to produce the corresponding amide derivatives **3**. The ester group was hydrolyzed under standard condition (NaOH, ethanol) to produce corresponding acid **4**. The acid was reacted with ethylchloroformate and *N*-methyl morpholine in THF to convert the carboxylic acid to an anhydride.⁹ The anhydride was reacted immediately with hydroxylamine HCl to produce the corresponding hydroxamic acid derivative **1**.

TLC, SiO₂, EtOAc/hexanes 2 : 1, R_f = 0.41 ¹H-NMR (300 MHz, DMSO-d₆) δ 11.20 (s, 1H), 9.12 (s, 1H), 7.80 (m, 4H), 7.61 (s, 1H), 2.10 (s, 9H), 1.78 (s, 6H). IR ν_{max} (KBr) 3298,



Scheme 1. Reaction conditions; (a) SOCl₂, reflux; (b) 1-adamantamine HCl, pyridine; (c) NaOH, EtOH; (d) ethylchloroformate, *N*-methyl morpholine, THF; (e) hydroxylamine HCl, TEA, DMF

2911, 1634, 1542 cm^{-1} . Ms-FAB (m/e) 315 ($M^+ + 1$).

DPPH assay. DPPH reagent was prepared at a DPPH concentration of 80 $\mu\text{g}/\text{mL}$ in MeOH. A test sample (50 μL) was dissolved in DMSO and mixed with 100 mM Tris-HCl buffer (pH 7.4, 50 μL), distilled water, and 400 μL of DPPH ethanolic solution (50 μL). The mixture was shaken well and allowed to stand for 20 min in the dark. The absorbance was measured at 515 nm using an Elx800 microtiter plate reader (Bio-Tek Instruments, Vermont, USA).

MTT growth assay. HaCaT keratinocytes were maintained in DMEM (Gibco, Grand Island, NY, USA) supplemented 10% fetal bovine serum, previously inactivated at 56 $^{\circ}\text{C}$ for 20 min. The cytotoxic effects of test materials were monitored by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as dose dependent manner.

Lipid peroxidation. HaCaT keratinocytes were grown in DMEM medium containing 10% fetal bovine serum and 1% antibiotic and antimycotic solution. For experiments, cells were maintained in DMEM supplemented with 1% fetal bovine serum (FBS) and test materials for 18 h. After HaCaT keratinocytes were incubated with test materials for 18 h, the cells were exposed to 4 mM *t*-BOOH for 4 h. Following incubation, the cell were washed twice in phosphate-buffered saline (PBS), and lysed by repetitive freeze/thawing in distilled water. To establish the levels of lipid peroxidation, malondialdehyde (MDA) and 4-hydroxy-2(*E*)-nonenal (4-HNE) levels were quantified using a commercial colorimetric lipid peroxidation assay kit (Calbiochem, San Diego, CA). This method analyzes MDA and 4-HNE by their reaction with a chromogen (*N*-methyl-2-phenylindole) at 45 $^{\circ}\text{C}$ to produce a stable chromogen. The reaction products were measured by spectrophotometry at 586 nm. The procedure was performed in accordance with the manufacturer's specifications and data were expressed in $\mu\text{mol}/\text{mg}$ protein.

Animal. Female SHR-1 hairless mice (Charles River, Japan), age ranges of 7-8 weeks, were used. They were housed 5 per cage (10 mice per groups) in a temperature (23 ± 3 $^{\circ}\text{C}$) and the relative humidity (40-60%) controlled room. Lighting was adjusted automatically to give a cycle of 12 hours light and 12 hours dark. Throughout the study, the animals were allowed to access freely to laboratory diets (Purina Co., Korea) and tap water. The water analysis for bacteriologic and chemical contaminant was regularly conducted by local water supply authority. No contaminants were present in diet or water to the levels of interfering with the objective of the study. One week of acclimatization period, was allowed before any treatment. Both animal care and protocol for this study were in accordance with IACUC (Institutional Animal Care and Use Committee) and OECD guideline.

Sq-OOH preparation and treatment. The procedure for preparation of Sq-OOH has been described previously.^{4a,4b} Briefly, Sq-OOH was produced from squalene by UVA irradiation (50 $\text{J}/\text{m}^2/\text{S}$). UV-irradiated squalene containing Sq-OOH was extracted with methanol. The methanol solution was displaced with ethanol solution by evaporation

under a nitrogen gas stream. Sq-OOH was confirmed for the hydroperoxide assay by HPLC coupled with chemiluminescence's detection. In group 2 and 3, mice were treated with 0.1 mL of Sq-OOH. Treatment was applied to the dorsum (neck to tail area, constituting an area of approximately 20 cm^2) using a plastic pipette, 2 times/7 day/weeks for 4 weeks. Skin samples were taken at 4 week. To group 3, 0.1 mL of ethanol solution containing 1% compound **1** was applied to the dorsum at 90 min prior to Sq-OOH treatment.

Evaluation of skin wrinkle. Treated animals were estimated grossly and photographed by digital camera (C740uz, Olympus[®], Japan) for evaluation of skin wrinkling. Skin replica image was acquired for statistical analysis. Skin negative replicas were made of the dorsum using a silicon-based gum material (Silflo, Flexico Development Ltd., UK). Animals were anesthetized with pentobarbital, and silflo was applied according to the manufacture's instructions. The appearance of wrinkles was assessed using an image analysis method. In brief, replicas were illuminated under an illumination lamp at an angle of 20 degree to maximize the formation of shadows behind the wrinkles. The image was observed under a stereoscopic microscope and photographed using a CCD camera installed with skin-visiometer SV600 software (C+K, Germany). Results of raw data were R1-R5 by analysis of the horizontal line. Each parameter acquired by horizontal analysis of replica image mean as follows;

Histology of skin. Animals were sacrificed at the end of the study. Dorsal skin was removed, and adherent subcutaneous fat was dissected away. Skin specimens obtained from dorsal skin were fixed in 10% buffered formalin, embedded in paraffin and sectioned 3-4 μm for light microscopy. Sections were stained with H & E, and Masson-Trichrome stain for collagen.

Results and Discussion

To evaluate antioxidant effect of hydroxamic acid derivative **1**, we compared its activity with known antioxidant such as trolox,¹⁰ EGCG¹¹ and kojic acid.¹² Diphenylpicrylhydrazyl (DPPH) is widely used for assessing the ability of antioxidant agent to transfer labile H-atoms to radicals, a common mechanism of radical scavenging. Trolox and EGCG showed a potent scavenging activity (Table 1). Compound **1** showed mild scavenging activity ($\text{IC}_{50} = 43.7$ mM). However, compound **4** (4-(*N*-adamantanylcabamoyl)benzoic acid) which is the precursor of hydroxamic acid derivative showed no inhibitory activity. These results suggest that the presence of hydroxamic moiety was critical for radical scavenging activity. Kojic acid showed no activity.


Lipid peroxidation is an indicator of possible free radical damage to cells. Cytotoxicity and inhibitory potency of compound **1** in lipid peroxidation were evaluated in HaCat cells. Cell viability was assessed by the MTT reduction assay. HaCat cells were resistant to up to 10 μM concentration of all test materials (Fig. 2).

After confirming cell viability, we evaluated inhibitory activity of compound **1** in lipid peroxidation. Their activities


Table 1. Radical-scavenging Effects

Compound	DPPH IC ₅₀ (μ M)
Trolox	20.7
EGCG	1.52
Kojic acid	–
Compound 1	43.7
Compound 4	–

–; Not effective.

**Figure 2.** Cytotoxic effects of test compounds in HaCat cell line. All compounds were tested at 10 μ M concentration.


were examined in terms of ability to reduce the oxidative factors such as malondialdehyde (MDA) and 4-hydroxy-2(*E*)-nonenal (4-HNE), generated by TBHP (*tert*-butyl-hydroperoxide) in HaCaT cell line. Treatment of 4 mM of TBHP increased lipid peroxide level up to about three times as compared with untreated sample. When 10 μ M of compounds were treated, trolox, EGCG and compound 1 (4-(*N*-hydroxycarbonyl)phenyl)-*N*-adamantanylcaboxamide) were active (Fig. 3). Trolox decreased the level of lipid peroxidation by about 40% in contrast with TBHP-

**Figure 3.** Inhibitory effects on lipid peroxidation induced by TBHP in HaCat cell line. All compounds were tested at 10 μ M concentration.

treated control. Reduced lipid peroxidation level was obtained about 50% after being treated with compound 1, which was close to the effect with EGCG. In this assay, compound 4 and kojic acid also showed no inhibitory activity at the same concentration.

In addition, the effect of compound 1 on the wrinkle formation by Sq-OOH was investigated in order to evaluate its availability for suppressing the lipid peroxidation in animal model. The features of the dorsal skin and replica images of hairless mice in groups 1 [vehicle, Sq-OOH(-)], group 2 [vehicle, Sq-OOH(+)] and group 3 [compound 1, Sq-OOH(+)] were shown in Figure 4. At the end of 4-week topical application period, all the mice in Sq-OOH treated group exhibited wrinkles with clear and fine appearance (A2 and B2). However, reduced wrinkle was observed in the mice group 3 (A3 and B3). Based on these results, it is suggested that the application of compound 1 prevents the formation of wrinkle induced by Sq-OOH.

At the end of experimental period, the skin sections were cut and stained with H&E, and Masson-Trichrome stain for collagen. A lot of alteration in epidermal layer was seen in

**Figure 4.** Photographs of skin surface (A) and replica image (B) of hairless mouse. 1: Untreated skin, 2: Sq-OOH treated skin, 3: Sq-OOH and compound 1 treated skin.




Figure 5. Histological findings of hairless mouse. 1: Untreated skin, 2: Sq-OOH treated skin, 3: Sq-OOH and compound 1 treated skin. A, Hematoxyline & Eosin stain, B, Masson & Trichrome stain, Original magnification X 160

Sq-OOH treated animal such as hypertrophy, hyperkeratosis and increasing epidermal thickness (A2 and B2). Epidermal thickness was particularly increased up to three times as compared with the untreated skin. However, a change of dermis layer was not significant. The formation of wrinkle induced by Sq-OOH may not be due to not alteration of dermal layer but to alteration of epidermal layer. The application of compound 1 suppressed alteration in epidermis layer to the Sq-OOH not application (A3 and B3). These histological data are correlated fairly well with features of dorsal skin and replica images.

In our best knowledge, this is the first report showing that an antioxidant agent prevents Sq-OOH induced wrinkle formation *in vivo*. Hydroxamic acid has both radical scavenging activity and iron chelating activity. The availability of new hydroxamic acid derivative **1** as a radical scavenger should contribute to the suppression of damage induced by Sq-OOH.

References

- (a) Meffert, H.; Diezel, W.; Sonnichsen, H. *Experientia* **1976**, *32*, 1397. (b) Chamberlain, J.; Moss, S. H. *Photochem. Photobiol.* **1987**, *45*, 625.
- (a) Uchino, T.; Tokunaga, H.; Onodera, H.; Ando, M. *Biol. Pharm. Bull.* **2002**, *25*, 605. (b) Ohsawa, K.; Watanabe, T.; Matsukawa, R.; Yoshimura, Y.; Imacda, K. *J. Toxicol. Sci.* **1984**, *9*, 151.
- (a) Kwon, T. W.; Olcott, H. S. *J. Food Sci.* **1966**, *31*, 552. (b) Yoo, H. C. H.; Shibamoto, T. *Lipids* **1992**, *27*, 50.
- (a) Chiba, K.; Sone, T.; Kawakami, K.; Onoue, M. *Exp. Dermatol.* **1999**, *8*, 471. (b) Chiba, K.; Kawakami, K.; Sone, T.; Onoue, M. *Skin Pharmacology and Applied Skin Physiology* **2003**, *16*, 242. (c) Lee, B. S.; Son, W. C.; Hyun, J. S.; Jung, J. M.; Rho, H. S.; Kim, D. H.; Shim, Y. C.; Kim, B. H. *Laboratory Animal Research* **2005**, *21*, 5.
- Vanjari, H.; Pande, R. *Journal of Pharmaceutical and Biochemical Analysis* **2003**, *33*, 783.
- Jackman, J. E.; Fierke, C. A.; Tumej, L. N.; Pirrung, M.; Uchiyama, T.; Tahir, S. H.; Hinds Gaul, O.; Raetz, C. R. H. *Journal of Biological Chemistry* **2000**, *275*, 11002.
- Farkas, E.; Enyedy, E. A.; Micera, G.; Garribba, E. *Polyhedron* **2000**, *19*, 1727.
- Taira, J.; Miyagi, C.; Aniya, Y. *Biochemical Pharmacology* **2002**, *63*, 1019.
- Reddy, A. S.; Kumar, M. S.; Reddy, G. R. *Tetrahedron Lett.* **2000**, *41*, 6285.
- (a) Peus, D.; Meves, A.; Pott, M.; Beyerle, A.; Pittelkow, M. R. *Free Radic. Biol. Med.* **2001**, *30*, 425. (b) Delicado, E. N.; Ferrer, A. S.; Carmona, F. G. *Biochimica et Biophysica Acta* **1997**, *1335*, 127.
- (a) Saffari, Y.; Hossein Sadrzadeh, S. M. *Life Sciences* **2004**, *74*, 1513. (b) Bors, W.; Michel, C. *Free Radic. Biol. Med.* **1999**, *27*, 1413.
- Mitani, H.; Koshiishi, I.; Sumita, T.; Imanari, T. *European Journal of Pharmacology* **2001**, *411*, 169.