

Bite Force, Occlusal Contact Area and Occlusal Pressure of Patients with Temporomandibular Joint Internal Derangement

Ki-Seo Kim, D.D.S., M.S.D., Jong-Hoon Choi, D.D.S., M.S.D., Ph.D.,
Seong-Taek Kim, D.D.S., M.S.D., Chong-Youl Kim, D.D.S., M.S.D., Ph.D.,
Hyung-Joon Ahn, D.D.S., M.S.D., Ph.D.

Department of Oral Medicine, College of Dentistry, Yonsei University

Temporomandibular joint (TMJ) internal derangement, especially disc displacement with reduction (DDwR) is the most common TMJ arthropathy and has been thought to do some effects on masticatory performance. Measuring of maximal bite force has been widely used as objective and quantitative method of evaluating masticatory performance, but previous studies showed various results due to various characteristics of subjects and different measuring devices and techniques. In a few studies about the correlation of bite force and temporomandibular disorders (TMD), some authors reported that bite force and masticatory performance would be reduced in patients with TMD because of pain. But the correlation of changes in structure of articular disc and masticatory performance has not been well investigated yet. In this study, to investigate the influences of non-painful disc change on the masticatory performance, we measured the value of maximal bite force, occlusal contact area and occlusal pressure of 39 patients with non-painful DDwR of the TMJ using pressure sensitive film, and compared it with that of 59 controls. The results are summarized as follows:

1. The maximal bite force ($P < 0.01$) and the occlusal contact area ($P < 0.05$) of the DDwR patients were greater than the controls.
2. There was no significant difference in occlusal pressure between the DDwR patients and the controls ($P > 0.05$).
3. The maximal bite force of the male group was greater than that of the female group ($P < 0.05$). However, the occlusal contact area and the occlusal pressure between the male and the female group didn't show significant difference ($P > 0.05$).

From the results above, we can suggest that DDwR could be a factor of changing bite force, but more controlled, large scaled and EMG related further study is needed.

Key words : Bite force, Occlusal contact area, Occlusal pressure, Pressure sensitive film, Disc displacement with reduction

Corresponding Author: Prof. Hyung-Joon Ahn

*Department of Oral Medicine, College of Dentistry,
Yonsei University, 134 Shinchon-Dong, Seodaemun-Ku,
Seoul 120-752, Korea*

E-mail: hjahn@yumc.yonsei.ac.kr

received: 2006-07-20

accepted: 2006-09-08

I. INTRODUCTION

Mastication, or chewing, is one of the main functions of the stomatognathic system¹⁾. Masticatory performance can be influenced by many changes on the stomatognathic system such as orthognathic surgery²⁾, muscle problems³⁾,

disorders of the temporomandibular joint⁴⁾, loss of occlusal support⁵⁾, different types of dentures⁶⁾, age⁷⁾ and sex⁸⁾.

Temporomandibular disorders(TMD) are divided into temporomandibular joint(TMJ) disorders and masticatory muscle disorders. TMJ disorders include internal derangement(or articular disc displacement) and it is the most common TMJ arthropathy and is characterized by several stages of clinical dysfunction that involve the condyle-disc relation. It is subdivided into disc displacement with reduction(DDwR) and disc displacement without reduction(DDw/oR)⁹⁾.

Masticatory function has been evaluated using both subjective and objective methods. Bite force is generally evaluated as objective method. Some investigators have reported the comparison of bite force between denture wearers and normal dentate subjects in the dental clinics^{10,11)}. But, until now, the effects of temporomandibular disorders(TMD) on masticatory performance have not frequently been investigated. A few studies have assessed bite force of patients with TMD^{12,13)}.

Although Helkimo¹²⁾ found an increasing bite force correlated with decreasing TMD symptoms, this observation could not be confirmed in a more recent study¹³⁾. Patients with DDw/oR had decreased bite force endurance in static chewing, caused by increasing pain in the joints¹⁴⁾. Electromyographic studies of patients with DDw/oR¹⁵⁾ revealed a significant reduction in bite force when patients were compared with controls. In addition to their studies of bite force and occlusal contact area, the same authors also evaluated the masticatory function of patients with DDw/oR using photometric devices¹⁶⁾.

All parameters were reduced in patients compared with controls and reductions in occlusal contact area and/or impaired chewing were discussed as sources of reduced masticatory function.

Many studies describe reduced chewing ability caused by pain among patients with TMD, especially DDw/oR^{12,14)}, but influence of structural

change of disc-condyle relation such as non-painful DDwR to masticatory function has not been well described.

A number of appliances for bite force measurement have been reported including strain gauge transducer¹⁷⁻²⁵⁾, biting fork²⁶⁻²⁸⁾ and bite force dynamometers^{29,30)}. But these methods were time consuming and unstable because of various level of vertical occlusal height.

Recently, an occlusal diagnostic system called the Dental Prescale System(Fuji Film Co., Tokyo, Japan.) was developed. As compared with the appliances described above, the latter system is more flexible, permits natural occlusion and prevents mandibular displacement during clenching³¹⁾. The reproducibility of this system has been confirmed for use in complete dentitions^{32,33)} and complete denture patients³¹⁾ as well as in implant supported fixed cantilever prosthesis³⁴⁾ and hemi-maxillectomy patients³⁵⁾.

The aim of this study is to investigate the influences of non-painful disc change on the masticatory performance by means of comparing the maximal bite force, the occlusal contact area and the biting pressure.

II. MATERIALS AND METHODS

1. Subjects

39 patients(18 males, 21 females) diagnosed with DDwR of the TMJ served as the experimental group. Their ages ranged from 21 to 38 years, with an average of 25.9 years. They were fully informed of the nature of the study and agreed to participate. Reciprocal clicking was a clinical sign of DDwR. The clicking can occur at various stages during opening and closing. Patients who have pain on mandibular movement were ruled out. Bony change on the condylar head area of the TMJ was evaluated by radiographic examination to rule out osteoarthritis and osteoarthrosis.

59 individuals were selected as controls(30 males, 29 females). The controls had no past history or

Table 1. Classification of subject group

Subjects		
	Control group	Experimental group
n	59	39
Male/Female	30/29	18/21
Age	25.9 ± 3.1(21~38)	25.6 ± 3.5(19~34)

present symptoms related to the TMD, joint sounds or pain, muscle pain, deviation on opening or limited mouth opening. Their ages ranged from 19 to 34 years, with an average of 25.6 years. In both the patients with DDwR and the controls, the upper and lower dentition was complete. They had neither malocclusion nor dento-facial deformity. Third molars were either absent or unerupted, uncontacted (Table 1).

2. Methods

1) Measurement of maximal bite force, occlusal contact area, occlusal pressure

Maximal bite force, occlusal contact area, occlusal pressure was measured by a Dental Prescale System (Fuji Film Co.), consisting of pressure sensitive sheets (Dental Prescale®, Fuji Film Co., Fig. 1) and an analyzing computer



Fig. 1. Dental Prescale® film (50H type-R, large size)

(Occluzer®, GC company Co., Fig. 2).

The Dental Prescale® consists of two papersheets and numerous microcapsules containing a red dye between them. When the teeth are brought into occlusion, these microcapsules rupture and discharge the dye, staining one of the papers red. The density of the color is in proportion to the degree of pressure applied. The occlusal contact area was measured, and the occlusal pressure and the bite force calculated for each subject. All measurements were made with the subject seated with the head upright, looking forward, and in an unsupported natural head position. The subject was instructed to bite as forcefully as possible. Tooth contact area and the density of the color for the occlusal pressure recorded on the Dental Prescale® were measured by an Occluzer® computer (FPD707) and the bite force was also calculated using this apparatus. In this study, the value of the occlusal contact area and the value of the bite force were explained, respectively, as the total contact areas and the total force calculated on contact area between the upper and lower teeth in occlusion for each subject. The value of the occlusal pressure was represented by the mean value of pressures measured on the contact area between the upper and lower teeth in occlusion for each subject.

Before recording, each subject was seated at the



Fig. 2. Occluzer®(GC company Co.)

upright position, in parallel with the Frankfort plane and instructed to bite so as to exert and produce maximal clenching. Dental Prescale® sheet(50H, type R) was placed between the upper and lower dentitions and the subjects were instructed to bite at the intercuspal position for 2~3 seconds. On occlusion, the midline of the Dental Prescale® was matched with that of the subject's dental arch. After testing, the sheets were stored in a lightproof box for at least 3 hours to stabilize the intensity of color.

2) Statistical analysis

Differences in maximal bite force, occlusal contact area, occlusal pressure between patients with DDwR and the controls, and between male and female group were evaluated with paired *t*-test.

III. RESULTS

1. Maximal bite force

The maximal bite force of experimental group(patients with DDwR) were greater than that of control group in both of male(989.0 ± 309.7 versus 818.6 ± 227.50 N) and female group(813.8 ± 186.3 versus 700.2 ± 178.2 N). Standard deviation were relative small in female group in both of control and experimental group. The mean and standard deviation(S.D.) of maximal bite force of each subject groups are listed below(Table 2).

The value of bite force measured in patients with DDwR was significantly greater than that in the controls ($P < 0.01$).

Table 2. Maximal bite force(Mean±S.D.)

Group	Maximal bite force (N)		
	Control group	Experimental group	<i>P</i> value
Male	818.6 ± 227.50	989.0 ± 309.7	0.0336*
Female	700.2 ± 178.2	813.8 ± 186.3	0.0339*
Total	760.4 ± 211.6	894.2 ± 262.7	0.0064**

* : $P < 0.05$ ** : $P < 0.01$

2. Occlusal contact area

The occlusal contact area of experimental group(26.8 ± 9.6 mm²) (patients with DDwR) were greater than the controls (23.1 ± 7.7 mm²). The mean and standard deviation(S.D.) of occlusal contact area of each subject groups are listed below(Table 3).

The value of occlusal contact area measured in patients with DDwR was greater than that in the controls($P < 0.05$).

3. Occlusal pressure

There was no significant difference in occlusal pressure between the DDwR patients(34.0 ± 4.7 Mpa) and the controls(33.9 ± 5.8 Mpa). The mean and standard deviation(S.D.) of occlusal pressure of each subject groups are listed below(Table 4).

4. Comparison of male and female group

In both of control and experiment group, only the value of the bite force measured in males was significantly greater than that in females($P < 0.05$).

The maximal bite force of the male group(818.6 ± 227.50 N in control, 989.0 ± 309.7 N in experimental group) was greater than that of the female group(700.2 ± 178.2 N in control, 813.8 ± 186.3 N in experimental group) in both of control and experimental group($P < 0.05$). However, the occlusal contact area and the occlusal pressure between the male and the female group didn't show significant difference($P > 0.05$). The mean and

Table 3. Occlusal contact area(Mean±S.D.)

Group	Occlusal contact area(mm^2)		
	Control group	Experiment group	P value
Male	23.5 ± 7.5	29.3 ± 11.6	0.0713
Female	22.8 ± 6.0	24.6 ± 7.1	0.3950
Total	23.1 ± 7.7	26.8 ± 9.6	0.0408*

* : $P < 0.05$ ** : $P < 0.01$

Table 4. Occlusal pressure(Mean±S.D.)

Group	Occlusal pressure(Mpa)		
	Control group	Experimental group	P value
Male	35.4 ± 5.1	34.5 ± 4.7	0.5700
Female	32.4 ± 6.2	33.5 ± 4.8	0.5066
Total	33.9 ± 5.8	34.0 ± 4.7	0.9615

* : $P < 0.05$ ** : $P < 0.01$

Table 5. Comparison between male and female control group

Group	Control group		
	Male	Female	P value
Maximal bite force(N)	818.6 ± 227.50	700.2 ± 178.2	0.0303*
Occlusal contact area(mm^2)	23.5 ± 7.5	22.8 ± 6.0	0.7084
Occlusal pressure (Mpa)	35.4 ± 5.1	32.4 ± 6.2	0.0504

* : $P < 0.05$ ** : $P < 0.01$

Table 6. Comparison between male and female experiment group

Group	Experimental group(DDwR)		
	Male	Female	P value
Maximal bite force (N)	989.0 ± 309.7	813.8 ± 186.3	0.0455*
Occlusal contact area (mm^2)	29.3 ± 11.6	24.6 ± 7.1	0.1513
Occlusal pressure (Mpa)	34.5 ± 4.7	33.5 ± 4.8	0.5023

* : $P < 0.05$ ** : $P < 0.01$

standard deviation(S.D.) of each evaluated values of control and experiment group are listed below (Table 5, 6).

IV. DISCUSSION

Most of previous studies provided variable results on maximal bite force, and it maybe due to a lack in the control of such variables to affect bite force measurements as age^{17,26,28,36)}, sex^{17,19,21,22,36)}, physical characteristics of subjects including height and weight^{19,26,21,22)}, the number of teeth present^{37,36)}, diverse population and different measuring devices and techniques^{29,30,38)}.

In the present study, age, sex, the number of tooth present and measuring device were considered except physical characteristics including body height and weight. Several authors found positive correlation between bite force and growth variables such as age and body height and weight. It is assumed that persons with larger body build, size and/or weight may exhibit a greater bite force. Kiliaridis *et al.*²⁷⁾ noticed a correlation between body height and the mean amplitude of the maximal bite force. Ringqvist³⁰⁾ found significant correlation between body height and maximal molar bite force in female, whereas Linderholm and Wennstörm²⁹⁾ and Braun *et al.*^{21,22)} did not. In this study, the standard deviation of maximal bite force ranged from 178.2 to 309.2, and it is consistent to that of the previous studies^{36,16,39)}. Therefore, a group of similar body height and weight could be an important factor in future studies.

In the previous studies, maximal bite force was measured using a metal material such as transducer, biting fork or bite force dynamometer of various sizes and thicknesses. The thickness(4~16 mm) is one of the reasons for unstable positions. Moreover, subjects might be reluctant to bite maximally because of fear of tooth damage or pain. In this study, maximal bite force was measured using the Dental Prescale System, which consists of pressure sensitive sheet(Dental Prescale®, Fuji Film Co., Tokyo, Japan) and a computer for

analysis(Occluzer®, GC company Co., Japan). Suzuki *et al.*³¹⁾ reported that Dental Prescale System has some advantages as follows: (i) the thin material induces only a small change in the occlusal vertical dimension(98 μm thickness), and makes it available to measure at a position near the intercuspal position, (ii) it is not necessary to prepare special measurement equipment, (iii) many patients may be examined for a short time, (iv) recording storage, even for an extended period, is simplified and (v) it is easy to explain the treatment to patients by using dental images. In addition, the clinical usefulness of this system was documented in terms of quantitative evaluation for occlusal load distribution^{35,31)}.

In the present study, the mean value of maximal bite force was greater than that obtained by previous studies. Bakke *et al.*⁴⁰⁾ examined a large sample of subjects(63 females and 59 males), and found a mean bite force of 522 N in males and 441 N in females. Dean *et al.*⁴¹⁾ found the mean values at the molar region to be 490 N in males and 402 N in females. Braun *et al.*^{21,22)} reported a mean value of 814 N for males and 615 N for females. This variability might be due to the differences of measuring point and appliances used.

There are few studies which have examined occlusal pressure, contact area and bite force in patients with TMD or anterior disc displacement of TMJ. Takenoshita *et al.*⁴²⁾ observed no difference in occlusal contact area between the patients with TMJ dysfunction and the controls without TMJ dysfunction. Our results show that the value of the occlusal contact area measured in patients with DDwR of the TMJ was slightly greater than that in the controls($P < 0.05$), and are thus not consistent with the results of Sato *et al.*¹⁶⁾, who reported smaller occlusal contact area in patients with anterior disc displacement of the TMJ. It remains obscure why the value of occlusal contact area measured in patients with DDwR was greater than that in the controls. But we can guess increased masticatory muscle activity, as in bruxism patients, could make increased tooth wear, thus occlusal

contact area could be increased. Further EMG related studies could be valuable. Our results also show that the bite force in patients with DDwR of the TMJ was greater than that in the controls, although there was no difference in the occlusal pressure. From these results we can think that occlusal contact area should have more important role in masticatory performance than occlusal pressure. In previous studies, the bite force and occlusal contact area in patients with DDwR of the TMJ was smaller than that in the controls and it was thought because the patients' symptoms – such as TMJ pain or masticatory muscle pain, fear to TMJ noise etc. – could disturb a stronger bite force. Sinn DP *et al.*⁴³⁾ examined the bite force in patients with TMJ dysfunction. They also found that the maximal bite force in patients was smaller than that in the controls without TMJ dysfunction. However, most of previous studies included various types of TMJ dysfunction, including myofascial pain, internal derangement of the TMJ(DD with or without reduction), osteoarthritis etc. In our study, we examined only in patients with non-painful DDwR of the TMJ, therefore, influence of disc displacement could be reflected more critically. In other words, increased bite force and occlusal contact area could be a cause of disc displacement. Both of the control and the DDwR group, maximal bite force of male group was greater than that of female group. This result is consistent with the results of previous studies^{8,40,36)}.

In conclusion, we can suggest that DDwR could be a factor of changing bite force regardless of pain, but more controlled, large scaled and EMG related further study is needed.

V. CONCLUSIONS

To investigate the influences of non-painful disc change on the masticatory performance, we measured the value of maximal bite force, occlusal contact area and occlusal pressure of 39 patients with non-painful DDwR of the TMJ using pressure sensitive film, and compared it with that of 59

controls. the results are summarized as follows:

1. The maximal bite force ($P < 0.01$) and the occlusal contact area ($P < 0.05$) of the DDwR patients were greater than those of the controls.
2. There was no significant difference in the occlusal pressure between the DDwR patients and the controls ($P > 0.05$).
3. The maximal bite force of the male group was greater than that of the female group ($P < 0.05$). However, there was no significant difference in the occlusal contact area and the occlusal pressure between the male and the female group ($P > 0.05$).

From the results above, we can suggest that DDwR could be a factor of changing bite force, but more controlled, large scaled and EMG related further study is needed.

REFERENCES

1. Peroz I, Tai S. Masticatory performance in patients with anterior disk displacement without reduction in comparison with reduction in comparison with symptom-free volunteers. Eur J Oral Sci 2002; 110:341-344.
2. Zarrinkelk HM, Throckmorton GS, Ellis E, Sinn DP. A longitudinal study of changes in masticatory performance of patients undergoing orthognathic surgery. J Oral Maxillofac Surg 1995;53:777-782.
3. Liu ZJ, Yamagata K, Kasahara Y, Ito G. Electromyographic examination of jaw muscles in relation to symptoms and occlusion of patients with temporomandibular joint disorders. J Oral Rehabil 1999;26:33-47.
4. Harper RP, Brown CM, Triplett MM, Villasenor A, Gatchel RJ. Masticatory function in patients with juvenile rheumatoid arthritis. Pediatr Dent 2000;22:200-206.
5. van der Bilt A, Olthoff LW, Bosman F, Oosterhaven SP. Chewing performance before and after rehabilitation of postcanine teeth in man. J Dent Res 1994;73:1677-1683.
6. Fontijn-Tekamp FA, Slagter AP, Van Der Bilt A, Van 'T Hof MA, Witter DJ, Kalk W, Jansen JA. Biting and chewing in overdentures, full dentures,

- and natural dentitions. *J Dent Res* 2000;79:1519–1524.
- 7. Hirai T, Ishijima T, Koshino H, Anzai T: Age-related change of masticatory function in complete denture wearers: evaluation by a sieving method with peanuts and a food intake questionnaire method. *Int J Prosthodont* 1994;7:454–460.
 - 8. Julien KC, Buschang PH, Throckmorton GS, Dechow PC: Normal masticatory performance in young adults and children. *Arch Oral Biol* 1996;41:69–75.
 - 9. Okeson J: Orofacial pain : guidelines for classification, assessment, and management. 3rd ed. 1996, pp. 127–141, Quintessence Pub Co Inc. Chicago.
 - 10. Atkinson HF, Ralph WJ: Tooth loss and biting force in man. *J Dent Res* 1973;52:225–228.
 - 11. Haraldson T, Karlsson U, Carlsson GE: Bite force and oral function in complete denture wearers. *J Oral Rehabil* 1979;6:41–48.
 - 12. Helkimo E, Carlsson GE, Carmeli Y: Bite force in patients with functional disturbances of the masticatory system. *J Oral Rehabil* 1975;2:397–406.
 - 13. Rudy TE, Greco CM, Yap GA, Zaki HS, Leader JK, Boston JR: The association between research diagnostic criteria for temporomandibular disorder findings and biting force and endurance in patients with temporomandibular disorders. *Pain Med* 2001;2:35–45.
 - 14. Stegenga B, Broekhuijsen ML, De Bont LG, van Willigen JD: Bite-force endurance in patients with temporomandibular joint osteoarthritis and internal derangement. *J Oral Rehabilfac Surg* 1992;19:639–647.
 - 15. Sato S, Ohta M, Goto S, Kawamura H, Motegi K: Electromyography during chewing movement in patients with anterior disc displacement of the temporomandibular joint. *Int J Oral Maxillofac Surg* 1998;27:274–277.
 - 16. Sato S, Ohta M, Sawatari M, Kawamura H, Motegi K: Occlusal contact area, occlusal pressure, bite force, and masticatory efficiency in patients with anterior disc displacement of the temporomandibular joint. *J Oral Rehabil* 1999;26:906–911.
 - 17. Garner LD, Kotwal NS: Correlation study of incisive biting forces with age, sex and anterior occlusion. *J Dent Res* 1973;52:698–702.
 - 18. Proffit WR, Fields HW, Nixon WL: Occlusal forces in normal and long-face adults. *J Dent Res* 1983;62:566–570.
 - 19. Fields HW, Proffit WR, Case JC, Vig KWL: Variables affecting measurements of vertical force. *J Dent Res* 1986;65:135–138.
 - 20. Waltimo A, Nyström M, Könönen M: Bite force and dentofacial morphology in men with severe dental attrition. *Scand J Dent Res* 1994;102:92–96.
 - 21. Braun S, Bantleon HP, Hnat WP, Freudenthaler JW, Marcotte MR, Johnson BE: A study of bite force, part 1. Relationship to various physical characteristics. *Angle Orthod* 1995;65:367–372.
 - 22. Braun S, Bantleon HP, Hnat WP, Freudenthaler JW, Marcotte MR, Johnson BE: A study of bite force, part 2. Relationship to various cephalometric measurements. *Angle Orthod* 1995;65:373–377.
 - 23. Kikuchi M, Korioth TWP, Hannam AG: The association among occlusal contacts, clenching effort, and bite force distribution in man. *J Dent Res* 1997;76:13–16.
 - 24. Raadsheer MC, van Eijden TM, van Ginkel FC, Prahl-Andersen B: Contribution of jaw muscle size and craniofacial morphology to human bite force magnitude. *J Dent Res* 1999;78:31–42.
 - 25. Throckmorton GS, Ellis E, Buschang PH: Morphologic and biomechanical correlates with maximum bite forces in orthognathic surgery patients. *J Oral Maxillofac Surg* 2000;58:515–524.
 - 26. Helkimo E, Carlsson GE, Helkimo M: Bite force and state dentition. *Acta Odontol Scand* 1977;35:297–303.
 - 27. Kiliaridis S, Kjellberg H, Wenneberg B, Engström C: The relationship between maximal bite force, bite force endurance and facial morphology during growth. A cross sectional study. *Acta Odontol Scand* 1993;51:323–331.
 - 28. Fogle LL, Glaros AG: Contribution of facial morphology, age and gender to EMG activity under biting and resting condition. A canonical correlation analysis. *J Dent Res* 1995;74:1496–1500.
 - 29. Linderholm H, Wennstöm A: Isometric bite force and its relation to general muscle force and body build. *Acta Odontol Scand* 1970;28:679–689.
 - 30. Ringqvist M: Isometric bite force and its relation to dimensions of the facial skeleton. *Acta Odontol Scand* 1973;31:35–42.
 - 31. Suzuki T, Kumagai H, Watanabe T, Uchida T, Nagao M: Evaluation of complete denture occlusal contacts using pressure-sensitive sheets. *Int J Prosthod* 1997;10:386–391.
 - 32. Watanabe M, Hattori Y, Satoh C: Bite force distribution on the dental arch in normal dentitions. In *Brain and Oral Functions, oral motor function and dysfunction*. Morimoto T, Matsuya T, Takada K, eds., 1995, pp. 399–403, Elsevier Science, Oxford.

33. Kumagai H, Suzuki T, Hamada T, Sondang P, Fujitani M, Nikawa H: Occlusal force distribution on the dental arch during various levels of clenching. *J Oral Rehabil* 1999;26:932-935.
34. Suzuki T, Kumagai H, Yoshitomi N, McGlumphy EA: Occlusal contacts of edentulous patients with mandibular hybrid dentures opposing maxillary complete dentures. *Int J Maxillofac Impl* 1999;14: 504-509.
35. Matsui Y, Ohno K, Michi K, Suzuki Y, Yamagata K: A computerized method for evaluating balance of occlusal load. *J Oral Rehabil* 1996;23:530-535.
36. Miyaura K, Matsuka Y, Morita M, Yamashita A, Watanabe T: Comparison of biting forces in different age and sex groups: a study of biting efficiency with mobile and non-mobile teeth. *J Oral Rehabil* 1999;26:223-227.
37. Hidaka O, Iwasaki M, Saito M, Morimoto T: Influence of clenching intensity on bite force balance, occlusal contact area, and average bite pressure. *J Dent Res* 1999;78:1336-1344.
38. Tortopidis D, Lyons MF, Baxendale RH, Gilmour WH: The variability of bite force measurement between sessions, in different positions within the dental arch. *J Oral Rehab* 1998;25:681-686.
39. Sondang P, Kumagai H, Tanaka E, Ozaki H, Nikawa H, Tanne K, Hamada T: Correlation between maximal bite force and craniofacial morphology of young adults in Indonesia. *J Oral Rehabil* 2003;30:1109-1117.
40. Bakke M, Holm B, Jensen L, Michler L, Moller E: Unilateral, isometric bite force in eight 68-year-old women and men related to occlusal factors. *Scand J Dent Res* 1990;98:149-158.
41. Dean JS, Throckmorton GS, Ellis EE, Sinn DP: A preliminary study of maximal voluntary bite force and jaw muscle efficiency in pre-orthognathic surgery patients. *J Oral Maxillofac Surg* 1992;50: 1284-1288.
42. Takenoshita Y, Ikebe T, Yamamoto M, Oka M: Occlusal contact area and temporomandibular joint symptoms. *Oral Surg Oral Med Oral Pathol* 1991;72: 388-394.
43. Sinn DP, De Assis EA, Throckmorton GS: Mandibular excursions and maximal bite forces in patients with temporomandibular joint disorders. *J Oral Maxillofac Surg* 1996;54:671-679.

국문요약

측두하악관절 내장증 환자의 교합력, 교합 접촉 면적 및 교합압

연세대학교 치과대학 구강내과학교실

김기서 · 최종훈 · 김성택 · 김종열 · 안형준

측두하악관절 내장증, 그 중에서도 정복성 관절원판 변위(Disc displacement with reduction; DDwR)는 가장 흔하게 발생하는 측두하악관절 질환으로, 저작 능력에 다양한 영향을 미칠 것으로 생각되어져 왔다. 저작 능력을 평가하는 객관적이고 가시적인 수단으로 최대 교합력의 측정이 널리 통용되어져 왔는데, 과거 많은 연구자들이 측두하악장애(TMD)와 교합력의 관계에 대한 연구를 시도하였으나, 피험자의 개인적 특성이나 측정 방법의 차이 등으로 인하여 통일성 있는 결과를 얻지 못하고 있으며, 측두하악장애로 인한 통증이 교합력 및 저작능력을 감소시킨다고 보고된 바 있으나 측두하악관절의 구조적 변화와 저작능력의 관계에 대한 연구는 아직 충분히 이루어져 있지 않다. 본 연구에서는 측두하악관절의 구조적 변화와 저작 능력의 관계를 파악해 보고자 개구시 무통성의 관절음을 보이는 환자 39명 및 대조군 59명을 대상으로 최대 교합력, 교합 접촉 면적 및 교합압을 감압필름법(pressure sensitivie film)을 이용하여 분석, 평가한 후 다음과 같은 결과를 얻었다.

1. 정복성 관절원판 변위 환자군이 대조군보다 높은 최대 교합력($P < 0.01$) 및 넓은 교합 접촉 면적($P < 0.05$)을 나타내었다.
2. 단위 면적당 평균 교합압은 대조군과 정복성 관절원판 변위 환자군 간에 있어서 유의성있는 차이를 보이지 않았다($P > 0.05$).
3. 남녀간의 차이에 있어서, 정복성 관절원판 환자군과 대조군 모두에서 남성이 여성보다 높은 교합압을 나타내었으나($P < 0.05$), 평균 교합압과 교합 접촉 면적은 유의성있는 차이를 보이지 않았다($P > 0.05$).

이상의 결과로부터 얻을 수 있는 결론은 측두하악관절의 정복성 관절원판 변위가 교합 기능을 변화시키는 한 요소로서 작용할 수 있다는 점이다. 향후의 연구에 있어서 근전도 등을 이용한 저작근 활성도 측정을 병행함이 유용할 것으로 사료되며, 교합력에 영향을 미치는 다른 요소가 통제된 보다 대규모 환자집단에 대한 연구가 필요할 것으로 사료된다.

주제어 : 교합력, 교합 접촉 면적, 교합압, 감압필름, 정복성 관절원판 변위
