

## Effect of Soil Factors on Vegetation Values of Salt Marsh Plant Communities: Multiple Regression Model

Ihm, Byung-Sun\*, Jeom-Sook Lee<sup>1</sup>, Jong-Wook Kim and Joon-Ho Kim<sup>2</sup>

*Department of Biology, Mokpo National University, Muan-Gun 534-830, Korea*

<sup>1</sup>*Department of Biology, Kunsan National University, Kunsan 573-701, Korea*

<sup>2</sup>*School of Biological Sciences, Seoul National University, Seoul 151-742, Korea*

**ABSTRACT:** The objective of the current study was to characterize and apply multiple regression model relating to vegetation values of the plant species over salt marshes. For each salt marsh community, vegetation and soil variables were investigated in the western coast and the southern coast in South Korea. Osmotic potential of soil and  $\text{Cl}^-$  content of soil as independent variable had positive and negative influences on vegetation values. Multiple regression model showed that vegetation values of 14 coastal plant communities were determined by pH of soil, osmotic potential of soil and sand content. The multiple regression equation may be applied to the explanation of distribution and abundance of plant communities with existing ordination plots.

**Key words:** Coastal plant community, Multiple regression model, Osmotic potential, Sand content, Vegetation value

### INTRODUCTION

Salt marshes embody boundary characteristics of terrestrial and marine communities. They are plastic coastal features, shaped by the interaction of water, sediments and vegetation. For stability, they require protection from high-energy waves and therefore usually develop in sheltered salt marshes. In some parts of the world they show extremely diverse vegetation, while in other regions they are dominated by a few plant species, usually grasses (Ihm et al. 2001a, Isacch et al. 2006). A better understanding of the environmental factors that determine plant community structure is a major goal of plant ecologists studying coastal wetlands. Sharp spatial boundaries of plant species on coastal wetland have been observed along environmental gradients (Armstrong et al. 1985). In controlling the distribution and abundance of wetland plants within and across coastal wetlands types, factors related to hydrology, salinity and soil texture are believed to play a major role (Ustin et al. 1982, Brewer and Grace 1990, Glenn et al. 1991, Ihm and Lee 1998).

In South Korea, the coastal communities are among the most poorly understood vegetation units, although research has been conducted on the physiognomic, floristic, and ecophysiological characteristics of halophytic species (Kim 1971, 1975, Kim et al. 1982, Kim and Song 1983, Oh and Ihm 1983, Ihm and Lee 1998, Jung and Kim 1998, Min and Kim 1999a, 1999b, 2000). The objective of the current study was to characterize and apply multiple regres-

sion model relating to the vegetation values of the plant communities over salt marshes.

### MATERIAL AND METHODS

Two coastal areas in South Korea were investigated: 1) the western coast at Kimje, Iksan, Kochang, Yongkwang, Hampyung, Muan, Shinan, and Yongam; 2) the southern coast at Haenam, Changhung, Bosung, Sunchon, and Masan. Coastal vegetation in these regions was monitored from September 1997 to August 2001. Salt marshes were described according to their hydrology, salinity, soil texture, etc. (Ihm and Lee, 1998, Ihm et al. 2001a, b).

For each community, we measured data of twelve soil variables on vegetation value (VV) (= cover (%) in quadrat  $\times$  plant height (cm),  $\% \cdot \text{cm}$  or  $\text{cm}^3$ ), which indicates occupying or dominant volume and competitive ability as the dependent variable for multiple regression analysis of vegetation (Lee and Park 2005). In order to examine the soil environment near the roots, samples were taken at a depth of 10 cm from the surface. Air-dried samples of soils were used for physico-chemical analysis. Water potential and osmotic potential of soils were determined by the method of Ihm (1989). Soil texture was determined by the Köhn's apparatus method. Organic matter content of soil was determined by ashing the samples at 550°C for 4hr. Electrical conductivity and pH of soil were measured by S-C-T meter (YSI Model 33) and Orion Ion-analyser (Model 407A), using a water to fresh soil ratio of about 5:1(w/w).

\* Corresponding author; Phone: +82-61-450-2343, e-mail: ihmbs@mokpo.ac.kr

Total Kjeldahl nitrogen content of soil was determined by micro-Kjeldahl method (Allen et al. 1986).  $\text{Cl}^-$  content of soil was measured by argentometric method (Kalthoff and Stenger 1947) and  $\text{Na}^+$  content of soil was measured by flame photometer (Coleman 51).

These soil variables were entered into a backward stepwise multiple regression with vegetation values as the dependent variable (SAS 2000). We constructed linear models to evaluate significant correlations between these soil variables and vegetation values. The significant variables at the 0.1000 level were selected in the model.

## RESULTS AND DISCUSSION

Vegetation value ranged from 900 to 9,000 in 14 plant communities of salt marshes. pH of soil ranged from 5.6 to 7.2. Water potential of soil ranged from  $-5.13$  to  $-0.95$  MPa. Soil moisture content ranged from 15.3 to 29.0%. Osmotic potential of soil ranged from  $-3.83$  to  $-0.75$  MPa. Electrical conductivity of soil ranged from 0.63 to 1.92 mS/cm.  $\text{Cl}^-$  content of soil ranged from 1.8 to 22.2 mg/g.  $\text{Na}^+$  content of soil ranged from 3.9 to 21.3 mg/g. Total Kjeldahl nitrogen of soil ranged from 0.15 to 0.65 mg/g. Organic matter content of soil ranged from 2.6 to 4.9%. Sand content of soil ranged from 8 to 82%. Silt content of soil ranged from 11 to 62%. Clay content of soil ranged from 8 to 33%.

In simple regression models analyzed to account for the vegetation value variability,  $r$  in models using osmotic potential of soil and  $\text{Cl}^-$  content of soil as independent variable are 0.541 and 0.618

(both  $P < 0.05$ ). Osmotic potential and  $\text{Cl}^-$  content had positive and negative influences on vegetation values (Table 1, Fig. 1 and 2). The relationships between vegetation value and pH, water potential, moisture content, electrical conductivity,  $\text{Na}^+$  content, total Kjeldahl nitrogen content, organic matter content and contents of sand, silt and clay were not statistically significant ( $P > 0.05$ , Table 1).

In multiple regression models analyzed to account for the vegetation value variability,  $r$  in model using soil variables as indepen-




Fig. 1. Relationship between vegetation value and osmotic potential of soil for 14 plant communities of salt marshes. The regression line is shown in Table 1.

Table 1. Simple regression analyses relating to vegetation values as dependent variables in 14 plant communities of salt marshes

Independent variable	Regression coefficient	Intercept	$r$	N	$P$
pH	1,543.417	-6,280.338	0.237	14	NS
Water potential (-MPa)	-970.810	6,362.831	0.473	14	NS
Moisture content (%)	-66.929	5,453.795	0.114	14	NS
Osmotic potential (-MPa)	-1,444.888	6,714.503	0.541	14	<0.05
Electrical conductivity (mS/cm)	-2,835.548	6,750.530	0.396	14	NS
$\text{Cl}^-$ content (mg/g)	-245.955	6,212.613	0.618	14	<0.05
$\text{Na}^+$ content (mg/g)	-170.773	5,704.345	0.395	14	NS
Total Kjeldahl nitrogen (mg/g)	5,087.890	2,140.353	0.295	14	NS
Organic matter (%)	526.397	1,980.336	0.148	14	NS
Sand (%)	13.065	3,298.171	0.122	14	NS
Silt (%)	-18.231	4,521.373	0.118	14	NS
Clay (%)	-35.366	4,611.237	0.114	14	NS

NS,  $P > 0.05$ .

dent variable is 0.831. The variability in vegetation value could be explained by three variables of soil pH, osmotic potential of soil and sand content of soil (Table 2 and Eq. 1):  $P > F$  of soil pH, osmotic potential and sand content were 0.024, 0.0011 and 0.0107, respectively. The 3 soil variables were significant at the 0.05 level. Soil pH and osmotic potential had positive influences on vegetation values, but sand content of soil had negative influences.

The 14 coastal plant communities were plotted in the space defined by the vegetation value observed and the vegetation value predicted by the multiple regression model (Fig. 3). *Zoysia sinica*, *Suaeda japonica* and *Limonium tetragonum* communities were located in salt marshes with low osmotic potential and sand content of soil and high pH of soil. *Phacelurus latifolius*, *Scirpus fluviatilis* and *Phragmites communis* communities were located in salt marshes with high osmotic potential and sand content of soil and low pH of soil. With exiting ordination plots such as Canonical Correspondence Analysis (CCA), Detrended Correspondence Analysis (DCA), Principal Component Analysis (PCA) etc., the equation of Multiple Regression Analysis (MRA) can be used for explaining plant abun-




Fig. 2. Relationship between vegetation value and  $Cl^-$  content of soil for 14 plant communities of salt marshes. The regression line is shown in Table 1.

Table 2. Type II SS,  $F$  value and  $P > F$  of soil pH, osmotic potential and sand content in backward stepwise multiple regression

Variable	Type II SS	F value	$P > F$
Intercept	1,979,403	0.75	0.407
pH	18,842,045	7.12	0.024
Osmotic potential	53,601,236	20.25	0.0011
Sand content	25,913,658	9.79	0.0107




Fig. 3. Relationship between the vegetation value observed and the vegetation value predicted by the multiple regression model. The diagonal line represents that predicted values are equal to observed ones. SAHE: *Salicornia herbacea*, SUJA: *Suaeda japonica*, SUAS: *Suaeda asparagoides*, ATGM: *Atriplex gmelini*, ARSC: *Artemisia scoparia*, LITE: *Limonium tetragonum*, ASTR: *Aster tripolium*, JOSI: *Zoysia sinica*, SUMA: *Suaeda maritima*, SCFL: *Scirpus fluviatilis*, PHCO: *Phragmites communis*, CASC: *Carex scabrifolia*, TRMA: *Triglochin maritimum*, PHLA: *Phacelurus latifolius*.

$$\text{Vegetation value} = -6659.826 + 3248.345 (\text{pH}) - 3345.134 (\text{Osmotic potential}) - 93.426 (\text{Sand content}) \quad (r = 0.831, P < 0.01) \quad (1)$$

dance and distribution (Gauch 1982, Digby and Kempton 1987, Jongman et al. 1995).

Salt marsh vegetation is generally characterized by simplicity of structure and uniform species composition (Asri and Ghorbanli 1997). Each of our communities had one or sometimes two dominants with or without associated species. *Suaeda herbacea*, *S. japonica* and *S. maritima* are known as a pioneer species on reclaimed land and salt marsh with low soil osmotic potential ( $\sim 5.0$  MPa), low sand content of soil and high light intensity (Kim 1971, Chapman 1974, Beefink 1977, Kim and Song 1983, Ihm and Lee 1998, Ihm et al. 2001a). Also, several salt marsh species were located in estuaries or in salt marshes supplied by fresh water springs. They were dominated by *Carex scabrifolia*, *Phragmites communis*, *Scirpus fluviatilis* and *Triglochin maritimum* communities. They were located mostly in salt marshes with high soil osmotic potential and sand content of soil (Oh and Ihm 1983, Kim and Song 1983, Ihm et al. 2001a, b).

Because vegetation values were consistent with typical distribution patterns of plant communities, we suggest that pH of soil, os-

otic potential of soil and sand content play the primary role in determining plant distribution. Ihm and Lee (1998) reported that edaphic factors related to water potential of soil, salinity and soil texture also affected the distribution and abundance of wetland plants within and across coastal wetland types. Similar results were found in the same or other areas (Min and Kim 1999a, b, Kim and Ihm 1988, Ihm et al. 2001a).

### ACKNOWLEDGEMENT

This paper was supported in part by the 2003 research funds of Mokpo National University, Republic of Korea.

### LITERATURE CITED

- Allen SE, Grimshaw HM, Rowland AP. 1986. Chemical analysis. In: Methods in Plant Ecology, (Moore PB, Chapman SB, eds). Blackwell Scientific Publications, Oxford, pp 285-344.
- Armstrong W, Wright EJ, Lythe S, Gaynard TJ. 1985. Plant zonation and the effects of the spring-neap tidal cycle on soil aeration in humber salt marsh. *J Ecol* 73: 323-339.
- Asri Y, Ghorbanli M. 1997. The halophilous vegetation of the Orumieh lake salt marshes, NW Iran. *Plant Ecol* 132: 155-170.
- Beefink WG. 1977. The coastal salt marshes of western and northern Europe: An ecological and phytosociological approach. In: *Ecosystem of the World I*. Elsevier Sci Co., New York, pp 109-156.
- Brewer JS, Grace JB. 1990. Plant community structure in an oligohaline tidal marsh. *Vegetatio* 90: 92-107.
- Chapman VJ. 1974. *Salt Marshes and Salt Deserts of the World*. 2nd ed. Cramer, Lehre.
- Digby PGN, Kempton RA. 1987. *Multivariate Analysis of Ecological Communities*. Chapman and Hall, New York.
- Gauch HG. 1982. *Multivariate Analysis in Community Ecology*. Cambridge University Press, Cambridge.
- Glenn EP, O'Leary JW, Watson MC, Thompson TL, Kuehl RO. 1991. *Salicornia bigelovii* Torr.: An oilseed halophyte for seawater irrigation. *Science* 251: 1065-1067.
- Ihm B-S. 1989. Distribution of coastal plant communities in response to soil water potential and plant osmotic adjustment. (Ph. D. thesis). Seoul National Univ., Seoul.
- Ihm B-S, Lee J-S. 1998. Soil factors affecting the plant communities of wetland on southwestern coast of Korea. *Korean J Ecol* 21: 321-328.
- Ihm B-S, Lee J-S, Kim J-W. 2001a. Coastal vegetation on the western, southern, and eastern coasts of South Korea. *J Plant Biol* 44: 163-167.
- Ihm H-B, Ihm B-S, Lee J-S, Kim J-W, Kim H-S. 2001b. Carbon dioxide budget in *Phragmites communis* stands. *Korean J Ecol* 24: 335-339.
- Isacch JP, Costa CSB, Rodriguez-Gallego L, Conde D, Escapa M, Gagliardini DA, Iribarne OO. 2006. Distribution of saltmarsh plant communities associated with environmental factors along a latitudinal gradient on the south-west Atlantic coast. *J Biogeogr* 33: 888-900.
- Jongman RHG, ter Braak CJF, van Tongeren OFR. 1995. *Data Analysis in Community and Landscape Ecology*. Cambridge University Press, Cambridge.
- Jung Y-K, Kim J-W. 1998. Coastal sand dune vegetation in Kyungpook province. *Korean J Ecol* 21: 257-262.
- Kalthoff LM, Stenger VA. 1947. *Volumetric Analysis*, 2nd ed, Vol. 2. Interscience Publishers, New York, pp 4242-4245.
- Kim CS. 1971. An ecological study on the process of plant community formation in tidal land. *J Plant Biol (Korean J Bot)* 14:163-169.
- Kim CS. 1975. A study on standing crops in *Phragmites communis* communities and their environmental factors. *J Plant Biol (Korean J Bot)* 18: 129-134.
- Kim CS, Ihm BS. 1988. Studies on the vegetation of the salt marsh in the southwestern coast of Korea. *Korean J Ecol* 11: 175-192.
- Kim CS, Song TG. 1983. Ecological studies on the halophyte communities at western and southern coasts in Korea (IV) -The halophyte communities at the different salt marsh habitats -. *Korean J Ecol* 6: 167-176.
- Kim J-H, Kim HS, Lee IK, Kim JW, Mun HT, Suh KH, Kim W, Kwon DH, Yoo SA, Suh YB, Kim YS. 1982. Studies on the estuarine ecosystem of the Nagdong River. *Proc Coll Natur Sci SNU* 7: 121-163.
- Lee, K-S, Park S-D. 2005. Relationship between the aboveground vegetation structure and fine roots of the topsoil in the burnt forest areas, Korea. *Korean J Ecol* 28: 149-156.
- Min BM, Kim J-H. 1999a. Plant distribution in the relation to soil properties of reclaimed lands on the west coast of Korea. *J Plant Biol* 42: 279-286.
- Min BM, Kim J-H. 1999b. Plant community structure in reclaimed lands on the west coast of Korea. *J Plant Biol* 42: 287-293.
- Min BM, Kim J-H. 2000. Plant succession and interaction between soil and plants after land reclamation on the west coast of Korea. *J Plant Biol* 43: 41-47.
- Oh K-H, Ihm B-S. 1983. Seasonal changes in the productivity and soil nutrients of *Phragmites communis* community in the salt marsh of the Sumjin-River estuary. *Korean J Ecol* 6: 90-97.
- SAS. 2000. *SAS/STAT User's guide*, Version 8.02. SAS Institute., Inc., Cary, NC.
- Ustin SL, Percy RW, Bayer DE. 1982. Plant water relations in a San Francisco Bay salt marsh. *Bot Gaz* 143: 368-373.

(Received June 9, 2006; Accepted August 21, 2006)