

Synthesis and Biological Evaluation of Arylsulfonylpiperazine Derivatives as 5-HT₆ Receptor Ligands

Sun Ah Jeon, Hyunah Choo, Woo-Kyu Park,[†] Hyewhon Rhim, Soo Y. Ko,[‡]
Yong Seo Cho, Hun Yeong Koh,[§] and Ae Nim Pae*

Life Sciences Division, Korea Institute of Science and Technology, P.O. Box 131, Cheongryang, Seoul 130-650, Korea

*E-mail: anpae@kist.re.kr

[†]Pharmaceutical Screening Research Team, Korea Research Institute of Chemical Technology, Daejeon 305-343, Korea

[‡]Department of Chemistry, Ewha Womans University, Seoul 120-750, Korea

[§]Department of Chemistry, Inha University, Incheon 402-751, Korea

Received November 2, 2006

The 5-HT₆ antagonists are mainly related to the treatment of cognitive dysfunction or impairment associated with Alzheimer's disease and schizophrenia. There have been lots of efforts to develop 5-HT₆ antagonists. As in our efforts, arylsulfonylpiperazine derivatives **1-3** were designed, synthesized and biologically evaluated against the human recombinant 5-HT₆ serotonin receptor. Total 36 compounds were synthesized and the most active compound among the synthesized compounds is compound **2h** with an IC₅₀ value of 1.5 μM. The compound **2h** is novel as 5-HT₆ receptor ligand and could act as lead for the novel 5-HT₆ receptor ligands.

Key Words : Serotonin, 5-HT₆ Receptor, Antagonists

Introduction

Serotonin (5-hydroxytryptamine, 5-HT) is a major neurotransmitter which interacts with 5-HT receptors to produce a lot of its effects. There are 15 different human serotonin receptors that have been cloned and divided into 7 subclasses (5-HT₁₋₇).¹ The 5-HT₆ receptor is one of the latest subtypes to have been identified and belongs to the G-protein coupled receptor (GPCR) superfamily, which is positively coupled to adenylate cyclase.² 5-HT₅ receptors are present in brain regions such as the cerebral cortex, nucleus accumbens, caudate-putamen and hippocampus with high densities and thalamus and substantia nigra with low densities, which are associated with learning and memory.¹ Actually, compromised serotonergic function may have an important contribution to cognitive decline related to aging, Alzheimer's disease (AD) and schizophrenia.³ Thus serotonergic system became a potential target for the treatment of memory dysfunction.⁴ It is also suggested that the 5-HT₆ receptor has a major role in obesity, based on the knockout mice study that 5-HT₆ receptor-knockout mice are resistant to weight gain when exposed to a high-fat diet.⁵

Up to date, there have been various 5-HT₆ receptor antagonists developed, among which three compounds have

entered clinical trials (Fig. 1).⁶ SB-742457 developed by GlaxoSmithKline is in the phase II clinical trial for the treatment of cognitive dysfunction associated with Alzheimer's disease. The Lilly compound LY-483518 is also in the phase II clinical trial for the treatment of cognitive impairment associated with schizophrenia.⁶ SB-271046 is in the phase I clinical trial, developed by GlaxoSmithKline and also for the treatment of Alzheimer's disease and schizophrenia.

Most of 5-HT₆ receptor antagonists have a sulfone moiety and/or a piperazine moiety (Figure 1).⁶ As a part of our ongoing project to develop 5-HT₆ receptor antagonists, we designed, synthesized and biologically evaluated arylsulfonylpiperazine derivatives **1-3** as novel 5-HT₆ receptor ligands (Fig. 2).

Results and Discussion

Design. The arylsulfonylpiperazine derivatives **1-3** were designed to possess a sulfone moiety and a piperazine moiety as well as a hydrophobic aromatic group through the structural analysis of the 5-HT₆ antagonists reported (Fig. 2). As shown in Figure 1, all the three 5-HT₆ antagonists in clinical trial have the three moieties in common. Particularly, the designed compounds have a characteristic arylsulfonyl-





Figure 1. 5-HT₆ antagonists under clinical trials.



- 1 R¹ = phenyl, R² = CF₃
 2 R¹ = ethyl, R² = CF₃
 3 R¹ = ethyl, R² = CN
 R³ = H, F, Cl, CH₃, OMe or dimethyl

Figure 2. Structures of arylsulfonylpiperazine derivatives.


piperazine which is linked to the hydrophobic group such as a trifluoromethylphenoxy and a cyanophenoxy group.

Chemistry. The arylsulfonylpiperazine derivatives 1-3 were synthesized from β -chloroketones 4 and 5 in three steps, respectively, as racemic mixtures (Scheme 1). The β -chloroketones 4 and 5 were treated with NaBH₄ in a solution of aq. THF at room temperature to give secondary alcohols 6 and 7 in 85% and 89% yields, respectively. The secondary alcohols 6 and 7 underwent Mitsunobu reactions with *p*-trifluoromethylphenol 8 (R² = CF₃) and *p*-cyanophenol 8 (R² = CN) by treating with DIAD and PPh₃ in THF at room temperature to afford compounds 9, 10 and 11 in 49%, 66% and 28% yields, respectively. The compounds 9, 10 and 11 were coupled with various arylsulfonylpiperazines 12 which have twelve different R³ substituents such as H, *o*-Cl, *m*-Cl, *p*-Cl, *o*-F, *m*-F, *p*-F, *o*-Me, *m*-Me, *p*-Me, *m*-OMe and 3,4-Me₂. Thus, the compounds 9, 10 and 11 underwent S_N2 type reactions with twelve arylsulfonylpiperazines 12 by treatment with K₂CO₃ in acetonitrile under reflux conditions to give the target compounds 1-3 in 24-96% yields. Total 36 of the title compounds 1-3 were synthesized.

Biological Evaluation. All the synthesized compounds were biologically evaluated against the human recombinant 5-HT₆ serotonin receptor stably expressed by HEK293 cell line through [³H]-lysergic acid diethylamide (LSD) binding assay.⁷ The results are summarized in Table 1. The compounds 1 with a phenyl group as R¹ are not active against the human recombinant 5-HT₆ serotonin receptor up to 10 μ M, except for the compound 1e (Table 1) with an *o*-fluoro-

Table 1. IC₅₀ values of the arylsulfonylpiperazine derivatives 1 against 5-HT₆ receptor

compd	R ¹	R ²	R ³	%inhibition at 10 μ M	IC ₅₀ (μ M)
1a	phenyl	CF ₃	H	24	>10
1b	phenyl	CF ₃	<i>o</i> -Cl	32	>10
1c	phenyl	CF ₃	<i>m</i> -Cl	30	>10
1d	phenyl	CF ₃	<i>p</i> -Cl	23	>10
1e	phenyl	CF ₃	<i>o</i> -F	52	6.0
1f	phenyl	CF ₃	<i>m</i> -F	32	>10
1g	phenyl	CF ₃	<i>p</i> -F	41	>10
1h	phenyl	CF ₃	<i>o</i> -Me	41	>10
1i	phenyl	CF ₃	<i>m</i> -Me	34	>10
1j	phenyl	CF ₃	<i>p</i> -Me	33	>10
1k	phenyl	CF ₃	<i>m</i> -OMe	30	>10
1l	phenyl	CF ₃	3,4-Me ₂	20	>10
2a	ethyl	CF ₃	H	58	4.2
2b	ethyl	CF ₃	<i>o</i> -Cl	32	>10
2c	ethyl	CF ₃	<i>m</i> -Cl	59	4.3
2d	ethyl	CF ₃	<i>p</i> -Cl	54	7.8
2e	ethyl	CF ₃	<i>o</i> -F	43	>10
2f	ethyl	CF ₃	<i>m</i> -F	55	4.7
2g	ethyl	CF ₃	<i>p</i> -F	44	>10
2h	ethyl	CF ₃	<i>o</i> -Me	60	1.5
2i	ethyl	CF ₃	<i>m</i> -Me	50	7.1
2j	ethyl	CF ₃	<i>p</i> -Me	58	6.2
2k	ethyl	CF ₃	<i>m</i> -OMe	50	9.6
2l	ethyl	CF ₃	3,4-Me ₂	47	>10
3a	ethyl	CN	H	29	>10
3b	ethyl	CN	<i>o</i> -Cl	37	>10
3c	ethyl	CN	<i>m</i> -Cl	43	>10
3d	ethyl	CN	<i>p</i> -Cl	37	>10
3e	ethyl	CN	<i>o</i> -F	60	3.6
3f	ethyl	CN	<i>m</i> -F	67	2.4
3g	ethyl	CN	<i>p</i> -F	48	>10
3h	ethyl	CN	<i>o</i> -Me	55	3.7
3i	ethyl	CN	<i>m</i> -Me	68	2.4
3j	ethyl	CN	<i>p</i> -Me	52	7.0
3k	ethyl	CN	<i>m</i> -OMe	58	6.7
3l	ethyl	CN	3,4-Me ₂	57	5.8
methiothepin					0.0016



Scheme 1. Reagents and conditions: (a) NaBH₄, aq. THF, rt, 85-89% yields; (b) DIAD, PPh₃, THF, rt 28-66% yields; (c) K₂CO₃, CH₃CN, reflux, 24-96% yield.

phenylsulfonylpiperazine group which has an IC_{50} value of $6.0 \mu M$. The compounds **2** and **3** with an ethyl group as R^1 are relatively more active than the compounds **1**. Among the compounds **2** and **3**, all the compounds **2h-2j** and **3h-3j** with a methyl group as R^3 showed inhibitory activity against the human recombinant 5-HT₆ serotonin receptor with IC_{50} values of 1.5 to $7.1 \mu M$, while only some of the compounds with a fluoro or chloro group showed inhibitory activity with IC_{50} values of 2.4 to $4.7 \mu M$. The compounds **2k** and **3k** with a *m*-methoxy group also show marginal inhibitory activity with IC_{50} values of 9.6 and $6.7 \mu M$, respectively. Among the compounds **1l**, **2l** and **3l** with a 3,4-dimethyl group at R^3 , only the compound **3l** show inhibitory activity with an IC_{50} value of $5.8 \mu M$. The most active compound among the series of the synthesized compounds is compound **2h** with an IC_{50} value of $1.5 \mu M$.

Discussion. According to the clinical trials, the 5-HT₆ antagonists are mainly related to the treatment of cognitive dysfunction or impairment associated with Alzheimer's disease and schizophrenia.⁶ Until now, there is no marketed drug as 5-HT₆ antagonist. In this study, arylsulfonylpiperazine derivatives were designed, synthesized and biologically evaluated against the human recombinant 5-HT₆ serotonin receptor. The synthesized compounds **1-3** were designed to have three characteristic moieties such as R^1 , R^2 and R^3 . R^1 is a phenyl group or an ethyl group which constitutes the main skeleton. The hydrophobic phenoxy group with R^2 substituents was expected to be one of the important factors for the inhibitory activity against 5-HT₆ receptor. R^3 substituents gave variety to the arylsulfonylpiperazine moiety.

According to the biological results shown in Table 1, R^1 group prefers a simple alkyl (ethyl) group to the aromatic (phenyl) group. The compounds with a phenyl group as R^1 show no inhibitory activity against 5-HT₆ receptor except the compound **1e**, while most of the compounds with an ethyl group as R^1 are active. There are two substituents as R^2 : a trifluoromethyl group and a cyano group. The two electron-withdrawing groups were tested as substituents at the R^2 position. There shows a little activity difference in the position and kind of the substituents at the R^3 position. The compound **2a** with hydrogen as R^3 shows inhibitory activity with an IC_{50} value of $4.2 \mu M$, while the compound **3a** has no activity up to $10 \mu M$. Among the compounds **2b-2d** and **3b-3d** with a chloro substituent as R^3 , only the compound **2c** with a *m*-chloro substituent shows inhibitory activity with an IC_{50} value of $4.3 \mu M$. In the case of fluoro substituents (compounds **2e-2g** and **3e-3g**), the compounds **2f**, **3e** and **3f** show activity with IC_{50} values of 4.7, 3.6 and $2.4 \mu M$, respectively. On the other hand, all the compounds with a methyl substituent show inhibitory activity with IC_{50} values of 1.5 to $7.1 \mu M$. A methyl substituent may be preferred to electronwithdrawing substituents such as fluoro and chloro for the 5-HT₆ antagonist effect. The compounds **2k** and **3k** with an electron donating group, a *m*-methoxy group, are only marginally active against 5-HT₆ receptor with IC_{50} values of 9.6 and $6.7 \mu M$, respectively. The compounds **1l** and **2l** with a 3,4-dimethyl group are not active but the

compound **3l** is active against 5-HT₆ receptor with an IC_{50} value of $5.8 \mu M$. The compound **2h** with an *o*-methyl group shows the best inhibitory activity against 5-HT₆ receptor among the synthesized compounds **1-3**. The activity of the compound **2h** is not so good compared to that of methiothepin, but the structure of the compound **2h** is novel as a 5-HT₆ ligand. Therefore, the compound **2h** could act as lead for the novel 5-HT₆ receptor ligands.

Based on the results of the limited structure activity relationship study on new 5-HT₆ receptor ligands, more extensive structure-activity relationship studies are under the due. A small molecule library with more various substituents for R^1 , R^2 and R^3 positions will be designed, synthesized and biologically evaluated against 5-HT₆ receptor.

Experimental

All the commercially available reagents were obtained from Aldrich and Fluka, and generally used without further purification. ¹H NMR and ¹³C NMR spectra were obtained on Bruker Advance 400 (or 300) spectrometer. Nuclear magnetic resonance spectra were acquired at 400 (or 300) MHz for ¹H, and 100 MHz for ¹³C NMR. Infrared spectra were obtained on a Perkin Elmer 16FPC FT-IR spectrometer using KBr pellet. CHCl₃ or neat. GC/MSD was obtained on a Hewlett Packard 5890. HRMS spectra were obtained on a JMS-700 mass spectrometer (Jeol). Analytical thin layer chromatographies (TLC) were carried out on precoated silica gel plates (Merck Kieselgel 60F254, layer thickness 0.25 mm). Flash column chromatographies were conducted with silica gel grade 230-400 mesh (Merck Kieselgel 60 Art 9385).

3-Chloro-1-phenylpropan-1-ol (6). To the solution of 3-chloro-1-phenyl-1-propanone **4** (4.0 g, 24 mmol) in 100 mL of THF was added NaBH₄ (1.8 g, 48 mmol) portionwise at 0 °C. The reaction mixture was stirred at room temperature for 40 min and quenched with water. The resulting mixture was extracted with EtOAc (100 mL × 2) and the organic layer was washed with water and brine, dried with MgSO₄, filtered and concentrated to afford the product **6** (3.6 g, 21 mmol) in 85 % yield as oil: ¹H NMR (400 MHz, CDCl₃) δ 7.40-7.37 (m, 4H), 7.35-7.32 (m, 1H), 4.98 (dd, $J = 8.5, 4.7$ Hz, 1H), 3.79-3.74 (m, 1H), 3.62-3.58 (m, 1H), 2.30-2.24 (m, 1H), 2.15-2.11 (m, 1H).

1-Chloropentan-3-ol (7). The procedure was same as the preparation of the compound **6**. 1-Chloropentan-3-one **5** was converted to the product **7** in 30 mmol-scale in 89% yield: ¹H NMR (400 MHz, CDCl₃) δ 3.72-3.67 (m, 3H), 1.87-1.85 (m, 2H), 1.54-1.48 (m, 2H), 0.96 (t, $J = 7.4$ Hz, 3H).

1-(3-Chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene (9). A solution of 3-chloro-1-phenyl-1-propanol **6** (1.0 g, 5.9 mmol), 4-(trifluoromethyl)phenol (1.0 g, 5.9 mmol) and triphenylphosphine (1.5 g, 5.9 mmol) in 15 mL of THF was treated with DIAD (1.2 mL, 5.9 mmol) at 0 °C under N₂ atmosphere. After 2 h, the reaction mixture was concentrated and purified by column chromatography with a 60 : 1 mixture of hexanes and EtOAc to give the product **9** (0.90 g,

2.9 mmol) in 49% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.46 (d, $J = 8.6$ Hz, 2H), 7.41-7.32 (m, 5H), 6.93 (d, $J = 8.7$ Hz, 2H), 5.45 (dd, $J = 11.3, 6.2$ Hz, 1H), 3.83-3.77 (m, 1H), 3.65-3.58 (m, 1H), 2.51-2.47 (m, 1H), 2.28-2.24 (m, 1H).

1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene (10). The procedure was same as the preparation of the compound **9**. 1-chloropentan-3-ol **7** was converted to the product **10** in 5.3 mmol-scale in 66% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.54 (d, $J = 8.8$ Hz, 2H), 6.90 (d, $J = 8.7$ Hz, 2H), 4.55 (m, 1H), 3.68-3.65 (m, 2H), 2.19-2.17 (m, 1H), 2.07-2.06 (m, 1H), 1.74-1.69 (m, 2H), 0.97 (t, $J = 7.4$ Hz, 3H).

4-(1-Chloropentan-3-yloxy)bezointrile (11). The procedure was same as the preparation of the compound **9**. 1-chloropentan-3-ol **7** was converted to the product **11** in 2.3 mmol-scale in 28% yield: $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.60-7.56 (m, 2H), 7.00-6.96 (m, 2H), 4.58-4.55 (m, 1H), 3.67-3.63 (m, 2H), 2.20-2.17 (m, 1H), 2.09-2.06 (m, 1H), 1.77-1.70 (m, 2H), 0.97 (t, $J = 7.5$ Hz, 3H).

1-(3-Phenyl-3-(4-(trifluoromethyl)phenoxy)propyl)-4-(phenylsulfonyl)piperazine (1a). A mixture of 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** (48 mg, 0.15 mmol), 1-(phenyl sulfonyl)piperazine (38 mg, 0.17 mmol) and K_2CO_3 (25 mg, 0.18 mmol) in 2 mL of CH_3CN was refluxed for 48 h at 110 °C. The reaction mixture was cooled to room temperature and extracted with CH_2Cl_2 . The organic layer was dried over MgSO_4 , filtered, concentrated and purified by column chromatography with a 5 : 1 mixture of hexanes and EtOAc to give the product **1a** (31 mg, 0.06 mmol) in 40% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.69-7.67 (m, 2H), 7.54-7.52 (m, 1H), 7.49-7.47 (m, 2H), 7.32-7.30 (m, 2H), 7.23-7.16 (m, 5H), 6.77 (d, $J = 8.6$ Hz, 2H), 5.10 (dd, $J = 8.0, 5.0$ Hz, 1H), 2.94 (br s, 4H), 2.44-2.39 (m, 6H), 2.07-2.04 (m, 1H), 1.89-1.85 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 528.8 [(M + Na)] $^+$.

1-(2-Chlorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1b). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1b** in 0.13 mmol-scale in 74% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.04 (d, $J = 7.9$ Hz, 2H), 7.54-7.47 (m, 2H), 7.42-7.40 (m, 3H), 7.38-7.27 (m, 4H), 7.26-7.24 (m, 1H), 6.88 (d, $J = 8.6$ Hz, 2H), 5.24 (dd, $J = 7.8, 5.1$ Hz, 1H), 3.32-3.29 (m, 4H), 2.50-2.49 (m, 6H), 2.20-2.15 (m, 1H), 2.01-1.95 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 562.9 [(M + Na)] $^+$.

1-(3-Chlorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1c). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1c** in 0.05 mmol-scale in 55% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.75 (s, 1H), 7.64 (d, $J = 7.7$ Hz, 1H), 7.59 (d, $J = 8.0$ Hz, 1H), 7.51-7.47 (m, 1H), 7.40 (d, $J = 8.5$ Hz, 2H), 7.34-7.23 (m, 5H), 6.86 (d, $J = 8.5$ Hz, 2H), 5.21-5.18 (m, 1H), 3.05 (br s, 4H), 2.53-2.20 (m, 6H), 2.20-2.11 (m, 1H), 1.98-1.93 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 562.9 [(M + Na)] $^+$.

1-(4-Chlorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1d). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1d** in 0.12 mmol-scale in 75% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.71-7.69 (m, 2H), 7.68-7.52 (m, 2H), 7.41-7.37 (m, 2H), 7.32-7.25 (m, 5H), 6.85 (d, $J = 8.7$ Hz, 2H), 4.15-4.07 (m, 1H), 3.02 (br s, 4H), 2.50 (br s, 6H), 2.15-2.10 (m, 1H), 2.00-1.96 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 562.9 [(M + Na)] $^+$.

1-(2-Fluorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1e). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1e** in 0.13 mmol-scale in 85% yield: $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.84-7.82 (m, 1H), 7.60-7.58 (m, 1H), 7.40 (d, $J = 8.5$ Hz, 2H), 7.30-7.20 (m, 7H), 6.87 (d, $J = 8.5$ Hz, 2H), 5.23 (dd, $J = 7.8, 5.1$ Hz, 1H), 3.22 (br s, 4H), 2.52-2.50 (m, 6H), 2.22-2.15 (m, 1H), 1.99-1.95 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 546.9 [(M + Na)] $^+$.

1-(3-Fluorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1f). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1f** in 0.12 mmol-scale in 75% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.54-7.51 (m, 2H), 7.47-7.45 (m, 1H), 7.38 (d, $J = 8.6$ Hz, 2H), 7.31-7.22 (m, 6H), 6.84 (d, $J = 8.7$ Hz, 2H), 5.18 (dd, $J = 7.7, 5.0$ Hz, 1H), 3.03 (br s, 4H), 2.51-2.46 (m, 6H), 2.15-2.11 (m, 1H), 1.98-1.95 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 546.8 [(M + Na)] $^+$.

1-(4-Fluorophenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1g). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1g** in 0.04 mmol-scale in 24% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.79-7.76 (m, 2H), 7.41-7.32 (m, 2H), 7.31-7.20 (m, 7H), 6.85 (d, $J = 8.7$ Hz, 2H), 5.21-5.18 (m, 1H), 3.05 (br s, 4H), 2.54 (br s, 6H), 2.19-2.14 (m, 1H), 2.01-1.99 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 546.9 [(M + Na)] $^+$.

1-(3-Phenyl-3-(4-(trifluoromethyl)phenoxy)propyl)-4-(*o*-tolylsulfonyl)piperazine (1h). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1h** in 0.11 mmol-scale in 59% yield: $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.60 (d, $J = 8.3$ Hz, 1H), 7.50-7.43 (m, 3H), 7.40-7.24 (m, 7H), 6.88 (d, $J = 8.6$ Hz, 2H), 5.25 (dd, $J = 7.7, 5.1$ Hz, 1H), 3.20 (br s, 4H), 2.64 (s, 3H), 2.52 (br s, 6H), 2.22-2.13 (m, 1H), 2.02-1.95 (m, 1H). $\text{M/S} - \text{ESI } m/z$ 542.9 [(M + Na)] $^+$.

1-(3-Phenyl-3-(4-(trifluoromethyl)phenoxy)propyl)-4-(*m*-tolylsulfonyl)piperazine (1i). The procedure was same as the preparation of the compound **1a**. 1-(3-chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1i** in 0.10 mmol-scale in 66% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.55 (d, $J = 4.9$ Hz, 2H), 7.43-7.38 (m, 4H), 7.31-7.25 (m, 6H), 6.85 (d, $J = 8.7$ Hz, 2H),

5.18 (dd, $J = 7.9, 5.0$ Hz, 1H), 3.02 (br s, 4H), 2.53-2.47 (m, 6H), 2.44 (s, 3H), 2.15-2.12 (m, 1H), 1.97-1.95 (m, 1H). M/S – ESI m/z 542.9 [(M + Na)]⁺.

1-(3-Phenyl-3-(4-(trifluoromethyl)phenoxy)propyl)-4-tosylpiperazine (1j). The procedure was same as the preparation of the compound **1a**. 1-(3-Chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1j** in 0.08 mmol-scale in 49% yield: ¹H NMR (300 MHz, CDCl₃) δ 7.56 (d, $J = 8.2$ Hz, 2H), 7.31-7.14 (m, 10H), 6.77 (d, $J = 8.7$ Hz, 2H), 5.13 (br s, 1H), 2.99 (br s, 4H), 2.50 (br s, 6H), 2.37 (s, 3H), 7.97 (br s, 1H), 1.92 (br s, 1H). M/S – ESI m/z 542.9 [(M + Na)]⁺.

1-(3-Methoxyphenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1k). The procedure was same as the preparation of the compound **1a**. 1-(3-Chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1k** in 0.08 mmol-scale in 47% yield: ¹H NMR (400 MHz, CDCl₃) δ 7.45-7.39 (m, 3H), 7.34-7.26 (m, 8H), 7.25-7.25 (m, 1H), 6.85 (d, $J = 8.7$ Hz, 2H), 5.19 (dd, $J = 7.9, 5.1$ Hz, 1H), 3.86 (s, 3H), 3.03 (br s, 4H), 2.51-2.47 (m, 6H), 2.17-2.12 (m, 1H), 1.96-1.93 (m, 1H). M/S – ESI m/z 558.9 [(M + Na)]⁺.

1-(3,4-Dimethylphenylsulfonyl)-4-(3-phenyl-3-(4-(trifluoromethyl)phenoxy)propyl) piperazine (1l). The procedure was same as the preparation of the compound **1a**. 1-(3-Chloro-1-phenylpropoxy)-4-(trifluoromethyl)benzene **9** was converted to the product **1l** in 0.08 mmol-scale in 51% yield: ¹H NMR (400 MHz, CDCl₃) δ 7.50-7.46 (m, 2H), 7.39 (d, $J = 8.8$ Hz, 2H), 7.33-7.23 (m, 7H), 6.85 (d, $J = 8.7$ Hz, 2H), 5.18 (dd, $J = 7.9, 5.0$ Hz, 1H), 3.00 (br s, 4H), 2.53-2.16 (m, 6H), 2.29 (s, 6H), 2.17-2.11 (m, 1H), 1.96-1.91 (m, 1H). M/S – ESI m/z 556.9 [(M + Na)]⁺.

1-(Phenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2a). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2a** in 0.07 mmol-scale in 36% yield: ¹H NMR (400 MHz, CDCl₃) δ 7.75 (d, $J = 7.8$ Hz, 2H), 7.63-7.60 (m, 1H), 7.56-7.53 (m, 2H), 7.44 (d, $J = 8.1$ Hz, 2H), 6.89 (d, $J = 8.5$ Hz, 2H), 4.28 (br s, 1H), 3.01 (br s, 4H), 2.44-2.31 (m, 6H), 1.77 (br s, 2H), 1.67-1.62 (m, 2H), 0.91 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 480.8 [(M + Na)]⁺.

1-(2-Chlorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2b). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2b** in 0.11 mmol-scale in 58% yield: ¹H NMR (400 MHz, MeOD) δ 8.04-7.01 (m, 1H), 7.63-7.61 (m, 2H), 7.53-7.05 (m, 3H), 7.04 (d, $J = 8.6$ Hz, 2H), 4.46-4.45 (m, 1H), 3.27-3.25 (m, 4H), 2.51-2.44 (m, 6H), 1.86-1.82 (m, 2H), 1.71-1.68 (m, 2H), 0.95 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 514.7 [(M + Na)]⁺.

1-(3-Chlorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2c). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2c** in 0.05 mmol-scale in 29% yield: ¹H NMR (400

MHz, MeOD) δ 7.98-7.76 (m, 2H), 7.58-7.53 (m, 2H), 7.48-7.43 (m, 3H), 7.00-6.98 (m, 2H), 4.42-4.39 (m, 2H), 3.29-3.19 (m, 4H), 2.45-2.38 (m, 6H), 1.79-1.77 (m, 2H), 1.64-1.62 (m, 2H), 0.90 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 514.7 [(M + Na)]⁺.

1-(4-Chlorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2d). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2d** in 0.10 mmol-scale in 52% yield: ¹H NMR (400 MHz, MeOD) δ 7.77-7.75 (m, 2H), 7.67-7.64 (m, 2H), 7.49 (d, $J = 8.6$ Hz, 2H), 7.02 (d, $J = 8.6$ Hz, 2H), 4.46-4.40 (m, 1H), 3.00 (br s, 4H), 2.58-2.55 (m, 2H), 2.51-2.47 (m, 4H), 1.84-1.80 (m, 2H), 1.70-1.65 (m, 2H), 0.94 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 514.7 [(M + Na)]⁺.

1-(2-Fluorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2e). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2e** in 0.10 mmol-scale in 52% yield: ¹H NMR (400 MHz, MeOD) δ 7.84-7.80 (m, 1H), 7.73-7.68 (m, 1H), 7.50-7.48 (m, 2H), 7.40-7.33 (m, 2H), 7.03-7.01 (m, 2H), 4.46-4.40 (m, 1H), 3.13-3.12 (m, 4H), 2.55-2.50 (m, 2H), 2.46-2.44 (m, 4H), 1.87-1.77 (m, 2H), 1.71-1.64 (m, 2H), 0.93 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 498.8 [(M + Na)]⁺.

1-(3-Fluorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2f). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2f** in 0.06 mmol-scale in 29% yield: ¹H NMR (400 MHz, MeOD) δ 7.68-7.64 (m, 1H), 7.61-7.59 (m, 1H), 7.54-7.45 (m, 4H), 7.03-7.00 (m, 2H), 4.44-4.41 (m, 1H), 3.00 (br s, 4H), 2.57-2.53 (m, 2H), 2.51-2.44 (m, 4H), 1.84-1.78 (m, 2H), 1.71-1.64 (m, 2H), 0.94 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 497 [(M + Na)]⁺.

1-(4-Fluorophenylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2g). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2g** in 0.09 mmol-scale in 46% yield: ¹H NMR (300 MHz, CDCl₃) δ 7.72-7.68 (m, 2H), 7.40 (d, $J = 8.4$ Hz, 2H), 7.19-7.13 (m, 2H), 6.84 (d, $J = 8.7$ Hz, 2H), 4.23 (br s, 1H), 2.93 (br s, 4H), 2.40 (br s, 4H), 1.98 (br s, 2H), 1.71-7.52 (m, 4H), 0.85 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 498.8 [(M + Na)]⁺.

1-(*o*-Tolylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2h). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2h** in 0.13 mmol-scale in 71% yield: ¹H NMR (400 MHz, CDCl₃) δ 7.90-7.88 (m, 2H), 7.88-7.45 (m, 1H), 7.65-7.31 (m, 5H), 6.94 (d, $J = 8.7$ Hz, 5H), 4.34-4.32 (m, 1H), 3.18-3.15 (m, 4H), 2.64 (s, 3H), 2.49-2.42 (m, 6H), 1.81-1.79 (m, 2H), 1.70-1.66 (m, 2H), 0.94 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 494.8 [(M + Na)]⁺.

1-(*m*-Tolylsulfonyl)-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2i). The procedure was same as the

preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2i** in 0.09 mmol-scale in 48% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.48-7.47 (m, 2H), 7.40-7.34 (m, 4H), 6.84 (d, $J = 8.7$ Hz, 2H), 4.22 (t, $J = 5.7$ Hz, 1H), 2.96 (br s, 4H), 2.47-2.38 (m, 4H), 1.73 (br s, 2H), 1.63-1.55 (m, 2H), 1.25-1.18 (m, 2H), 0.58 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 493 [(M + Na)] $^+$.

1-Tosyl-4-(3-(4-(trifluoromethyl)phenoxy)pentyl)piperazine (2j). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2j** in 0.12 mmol-scale in 65% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.56 (d, $J = 8.2$ Hz, 2H), 7.38 (d, $J = 8.6$ Hz, 2H), 7.27 (d, $J = 7.9$ Hz, 2H), 6.84 (d, $J = 8.6$ Hz, 2H), 4.23-4.19 (m, 1H), 2.94 (br s, 4H), 2.14-2.40 (m, 4H), 1.73 (br s, 2H), 1.60-1.55 (m, 2H), 1.25-1.18 (m, 2H), 0.84 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 494.8 [(M + Na)] $^+$.

1-(3-Methoxyphenylsulfonyl)-4-(3-(4-trifluoromethyl)phenoxy)pentyl)piperazine (2k). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2k** in 0.11 mmol-scale in 59% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.54-7.48 (m, 2H), 7.34-7.31 (m, 1H), 7.26-7.24 (m, 2H), 7.02-7.00 (m, 2H), 4.45-4.39 (m, 1H), 3.88 (s, 3H), 3.06-2.99 (m, 4H), 2.55-2.43 (m, 6H), 1.84-1.76 (m, 2H), 1.71-7.64 (m, 2H), 0.93 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 507 [(M + Na)] $^+$.

1-(3,4-Dimethylphenylsulfonyl)-4-(3-(4-trifluoromethyl)phenoxy)pentyl)piperazine (2l). The procedure was same as the preparation of the compound **1a**. 1-(1-Chloropentan-3-yloxy)-4-(trifluoromethyl)benzene **10** was converted to the product **2l** in 0.06 mmol-scale in 31% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.54-7.46 (m, 3H), 7.31-7.26 (m, 2H), 6.91 (d, $J = 8.6$ Hz, 2H), 4.28 (br s, 1H), 2.99 (br s, 4H), 2.66 (br s, 6H), 2.33 (s, 6H), 1.77-1.64 (m, 2H), 1.58 (br s, 2H), 0.97-0.89 (m, 3H). M/S – ESI m/z 507 [(M + Na)] $^+$.

4-(1-(4-(Phenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3a). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3a** in 0.15 mmol-scale in 66% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.78-7.76 (m, 2H), 7.72-7.66 (m, 1H), 7.65-7.64 (m, 2H), 7.54-7.51 (m, 2H), 7.02-6.99 (m, 2H), 4.50-4.40 (m, 1H), 2.95 (br s, 4H), 2.51-2.47 (m, 2H), 2.46-2.42 (m, 4H), 1.80 (m, 2H), 1.69-1.67 (m, 2H), 0.92 (t, $J = 6.1$ Hz, 3H). M/S – ESI m/z 436 [(M + Na)] $^+$.

4-(1-(4-(2-Chlorophenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3b). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3b** in 0.16 mmol-scale in 74% yield: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.01-8.02 (m, 1H), 7.55-7.51 (m, 4H), 7.43-7.41 (m, 1H), 6.94-6.91 (m, 2H), 4.37 (t, $J = 5.7$ Hz, 1H), 3.37-3.31 (m, 4H), 2.47 (br s, 4H), 1.88-1.84 (m, 2H), 1.26 (t, $J = 7.1$ Hz, 2H), 0.93 (t, $J = 7.5$ Hz, 3H). M/S – ESI m/z 470 [(M + Na)] $^+$.

4-(1-(4-(3-Chlorophenylsulfonyl)piperzine-1yl)pentan-

3-yloxy)benzotrile (3c). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3c** in 0.13 mmol-scale in 61% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.69-7.63 (m, 4H), 7.62-7.56 (m, 1H), 7.49-7.47 (m, 2H), 6.95-6.93 (m, 2H), 4.40-4.37 (m, 1H), 2.90 (br s, 4H), 2.47-2.43 (m, 2H), 2.40-2.35 (m, 4H), 1.75-1.72 (m, 2H), 1.62-1.58 (m, 2H), 0.86-0.83 (m, 3H). M/S – ESI m/z 470 [(M + Na)] $^+$.

4-(1-(4-(4-Chlorophenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3d). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3d** in 0.15 mmol-scale in 67% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.76-7.73 (m, 2H), 7.66-7.63 (m, 2H), 7.55-7.52 (m, 2H), 7.02-6.98 (m, 2H), 4.47-4.42 (m, 1H), 2.95 (br s, 4H), 2.52-2.49 (m, 2H), 2.46-2.41 (m, 4H), 1.83-1.76 (m, 2H), 1.68-1.63 (m, 2H), 0.91 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 470 [(M + Na)] $^+$.

4-(1-(4-(2-Fluorophenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3e). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3e** in 0.21 mmol-scale in 96% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.83-7.82 (m, 1H), 7.75-7.65 (m, 1H), 7.56 (dd, $J = 6.9, 2.1$ Hz, 2H), 7.42-7.35 (m, 2H), 7.03 (dd, $J = 6.9, 2.1$ Hz, 2H), 4.51-4.41 (m, 1H), 3.11 (br s, 4H), 2.52-2.51 (m, 2H), 2.47-2.44 (m, 4H), 1.85-1.81 (m, 2H), 1.70-1.67 (m, 2H), 0.93 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 455.1 [(M + Na)] $^+$.

4-(1-(4-(3-Fluorophenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3f). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3f** in 0.17 mmol-scale in 77% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.69-7.68 (m, 1H), 7.61-7.59 (m, 1H), 7.56-7.48 (m, 4H), 7.02 (dd, $J = 6.9, 2.0$ Hz, 2H), 4.50-4.39 (m, 1H), 2.99 (br s, 4H), 2.52-2.43 (m, 6H), 1.82-1.80 (m, 2H), 1.69-1.66 (m, 2H), 0.92 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 454 [(M + Na)] $^+$.

4-(1-(4-(4-Fluorophenylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3g). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3g** in 0.21 mmol-scale in 94% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.85-7.82 (m, 2H), 7.56 (dd, $J = 6.9, 2.0$ Hz, 2H), 7.38 (t, $J = 8.7$ Hz, 2H), 7.02 (dd, $J = 6.9, 2.0$ Hz, 2H), 4.50-4.40 (m, 1H), 2.96 (br s, 4H), 2.52-2.42 (m, 6H), 1.81-1.79 (m, 2H), 1.69-1.67 (m, 2H), 0.93 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 454 [(M + Na)] $^+$.

4-(1-(4-(*o*-Tolylsulfonyl)piperzine-1yl)pentan-3-yloxy)benzotrile (3h). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3h** in 0.17 mmol-scale in 76% yield: $^1\text{H NMR}$ (400 MHz, MeOD) δ 7.87-7.86 (m, 1H), 7.59-7.53 (m, 3H), 7.43-7.40 (m, 2H), 7.04 (dd, $J = 6.9, 1.9$ Hz, 2H), 4.50-4.40 (m, 1H), 3.12-3.10 (m, 4H), 2.61 (s, 3H), 2.50-2.43 (m, 6H), 1.85-1.83 (m, 2H).

1.70-1.67 (m, 2H), 0.94 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 451.0 [(M + Na)]⁺.

4-(1-(4-(*m*-Tolylsulfonyl)piperazine-1yl)pentan-3-yloxy)benzotrile (3i). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3i** in 0.15 mmol-scale in 68% yield: ¹H NMR (300 MHz, MeOD) δ 7.60-7.59 (m, 1H), 7.57-7.51 (m, 5H), 7.03-7.00 (m, 2H), 4.45-4.43 (m, 1H), 2.95-2.94 (m, 4H), 2.51-2.42 (m, 9H), 1.81-1.79 (m, 2H), 1.70-1.66 (m, 2H), 0.93 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 450 [(M + Na)]⁺.

4-(1-(4-(Tosylpiperazine-1yl)pentan-3-yloxy)benzotrile (3j). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3j** in 0.18 mmol-scale in 83% yield: ¹H NMR (400 MHz, MeOD) δ 7.65-7.63 (m, 2H), 7.53-7.50 (m, 2H), 7.46 (d, $J = 8.1$ Hz, 2H), 7.00 (dd, $J = 6.9, 2.0$ Hz, 2H), 4.46-4.44 (m, 1H), 2.92 (br s, 4H), 2.51-2.41 (m, 9H), 1.81-1.80 (m, 2H), 1.69-1.65 (m, 2H), 0.92 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 450 [(M + Na)]⁺.

4-(1-(4-(3-Methoxysulfonylpiperazine-1yl)pentan-3-yloxy)benzotrile (3k). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3k** in 0.21 mmol-scale in 94% yield: ¹H NMR (400 MHz, MeOD) δ 7.54 (dd, $J = 6.9, 2.1$ Hz, 3H), 7.34-7.31 (m, 1H), 7.28-7.25 (m, 2H), 7.01 (dd, $J = 6.9, 2.1$ Hz, 2H), 4.50-4.40 (m, 1H), 3.89 (s, 3H), 2.97 (br s, 4H), 2.52-2.51 (m, 2H), 2.48-2.42 (m, 4H), 1.82-1.69 (m, 2H), 1.67-1.66 (m, 2H), 0.92 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 466 [(M + Na)]⁺.

4-(1-(4-(3,4-Dimethylsulfonylpiperazine-1yl)pentan-3-yloxy)benzotrile (3l). The procedure was same as the preparation of the compound **1a**. 4-(1-Chloropentan-3-yloxy)benzotrile **11** was converted to the product **3l** in 0.16 mmol-scale in 74% yield: ¹H NMR (400 MHz, MeOD) δ 7.53-7.51 (m, 3H), 7.48-7.46 (m, 1H), 7.40 (d, $J = 8.0$ Hz, 1H), 7.06 (dd, $J = 6.9, 2.1$ Hz, 2H), 4.50-4.40 (m, 1H), 2.93 (br s, 4H), 2.59-2.50 (m, 2H), 2.50-2.38 (m, 9H), 2.01 (s, 3H), 1.80-1.78 (m, 2H), 1.69-1.65 (m, 2H), 0.92 (t, $J = 7.4$ Hz, 3H). M/S – ESI m/z 464 [(M + Na)]⁺.

[³H]-LSD Binding assay to serotonin 5-HT₆ receptor. Membranes from stable HEK-293 cell line expressing the human recombinant 5-HT₆ serotonin receptor (PerkinElmer Life and Analytical Sciences, Boston, USA) were used. For the binding assay, aliquots of receptor membranes, 1.6 nM [³H]LSD and appropriate concentrations of test compounds were added to 0.25 mL of 50 mM Tris-HCl (pH 7.4) buffer containing 10 mM MgCl₂ and 0.5 mM EDTA. Incubations were carried out for 60 min at 37 °C, and these were terminated by rapid filtration using an Innotech cell harvester (Innotech Biosystems, Switzerland) through Whatman GF/C glass fiber filter presoaked in 0.3% polyethylenimine. The filter was covered with MeltiLex, sealed in a sample bag followed by drying in the microwave oven, and counted by MicroBeta Plus (Wallac, Finland). Nonspecific binding was determined in the presence of 0.5 μ M methiothepin. Competition binding studies were carried out with 7-8 varied concentrations of the test compounds run in duplicate tubes, and isotherms from three assays were calculated by computerized nonlinear regression analysis (GraphPad Prism Program, San Diego, USA) to yield inhibition values (IC₅₀).

Acknowledgments. This work was supported by Korea Ministry of Science and Technology.

References and Notes

- Hoyer, D.; Clarke, D. E.; Fozard, J. R.; Hartig, P. R.; Martin, G. R.; Mylecharane, E. J.; Saxena, P. R.; Humphrey, P. P. *Pharmacol. Rev.* **1994**, *46*, 157.
- Ruat, M.; Traiffort, E.; Arrang, J. M. *Biochem. Biophys. Res. Commun.* **1993**, *193*, 268.
- (a) Meneses, A. *Learn Memory* **2003**, *10*, 363. (b) Meneses, A. *Neurosci. Biobehav. Rev.* **1999**, *8*, 1111.
- Roth, B. L.; Hanizavareh, S. M.; Blum, A. E. *Psychopharmacology* **2004**, *174*, 17.
- Vickers, S. P.; Dourish, C. T. *Curr. Opin. Investig. Drugs* **2004**, *5*, 377.
- Holenz, J.; Pauwels, P. J.; Diaz, J. L.; Merce, R.; Codony, X.; Buschmann, H. *Drug Discov. Today* **2006**, *11*, 283.
- (a) Drummond, A. H.; Bucher, F.; Levitan, I. B. *Brain Research* **1980**, *184*, 163. (b) Drummond, A. H.; Bucher, F.; Levitan, I. B. *J. Biol. Chem.* **1980**, *255*, 6679.