



Rapid Enumeration of *Salmonella* spp. in Contaminated Pork Meat Using Competitive PCR

Aerie Moon and Weon Sang Choi*

Department of Biotechnology, Dongguk University, Gyeongju Korea

Competitive PCR을 이용한 돼지고기 오염 살모넬라의 신속 계수

문애리 · 최원상*

동국대학교 생명공학과

(Received September 23, 2007/Accepted December 5, 2007)

ABSTRACT – In this study, the competitive polymerase chain reaction (cPCR) was used to develop a direct enumeration method of *Salmonella* spp. in pork meat. After comparing three DNA extraction methods, the modified guanidine thiocyanate-phenol-chloroform method was chosen for *Salmonella* DNA extraction in artificially inoculated pork meat. The previously reported 284-bp *invA* gene (Rahn et al. Mol. Cell. Probes 1992) was tested for specificity, and 57 *Salmonella* strains and 24 non-*Salmonella* strains were evaluated. All *Salmonella* strains tested were *invA* positive, and all non-*Salmonella* strains produced no false positive amplification products. The detection limit achieved was as low as 1,460 colony-forming units (cfu) per 0.1 g of pork meat. For cPCR, the *invA* gene, which features a 82 bp-deletion, was cloned in the pGEM-4Z vector. A known amount of competitor DNA, which has the same primer binding sites, was co-amplified with *Salmonella* chromosomal DNA from the artificially inoculated pork meat. The cell-number determined by cPCR was approximately equal to the cfu from the most probable number (MPN) method. Finally, the whole procedure took only 5 hr.

Key words: Detection, Enumeration, *Salmonella* spp., PCR

Introduction

Salmonella spp. are members of the *Enterobacteriaceae*, and are divided into seven subspecies based on their biochemical characteristics. Serotyping has identified more than 2,000 serovars¹. *Salmonella* spp. have been isolated from a variety of foods. The ingestion of contaminated food can cause salmonellosis. Thus, a rapid detection and enumeration method will play an important role for epidemiological surveillance, monitoring of food-borne outbreaks, and the identification of sources of contamination in food processing plants.

Detection and enumeration of *Salmonella* continues to be an important issue in clinical and applied microbiology, and new media and diagnostic tests are continuously being developed^{2,3}. Biochemical substrate utilization is the basis

of species identification for *Salmonella*. The majority of *Salmonella* spp. are recognized as non-lactose fermenters (Lac⁻) and hydrogen-sulfide producers (H₂S⁺). However, because the majority of the H₂S⁺ and/or Lac⁻ colonies turn out not to be *Salmonella* spp., but insteads are related species such as *Proteus* or *Citrobacter*, conventional approaches require confirmatory testing of all H₂S⁺ and/or Lac⁻ colonies, which extends the time for identification³. Traditional culture methods for the detection of *Salmonella* spp. require about 5 days to obtain a negative result, and it takes at least 2 days for clinical laboratories to give a positive report^{4,6}.

Several polymerase chain reaction (PCR) assays, not dependent on substrate utilization or the expression of antigens and thereby circumventing the phenotypic variations, have been developed to detect *Salmonella* spp. Regions that have been selected for PCR detection include *agfA*, *fimA*, *hin*, *H-li*, *iagAB*, *IS200*, *invA*, *iroB*, *mkfA*, *ompC*, *oriC*, *spvR*, *viaB*, *fimY*, *rfb*, *pefA*, *sefA*, *iagA*, *mntH*, *gyrA*, and *prot6e*^{3,7-25}. However, limited data are available on the performance of PCR assays for the detection of *Salmonella*

*Correspondence to: Weon Sang Choi, Department of Biotechnology, Dongguk University, Gyeongju Korea
Tel: 82-54-770-2227, Fax: 82-54-770-2386
E-mail: weonsang@dongguk.ac.kr

in foods.

Recently, we reported that competitive PCR (cPCR) can be applied to directly estimate the numbers of *Listeria monocytogenes*²⁶⁾, *Salmonella enteritidis*²⁷⁾, and *Yersinia enterocolitica*²⁸⁾ in milk, and *Listeria monocytogenes* in pork meat²⁹⁾. In this study we evaluated the specificity of primers based on the invasion (*invA*) gene that is located on pathogenicity island 1 of *Salmonella* spp., which encodes a protein of the type III secretion system³⁰⁾ for *Salmonella* detection, and then utilized it for the enumeration of *Salmonella* in pork meat using cPCR.

Materials and Methods

Bacterial strains

The *Salmonella* strains (Table 1) were obtained from the Seoul Metropolitan City Institute of Health and Environment (Seoul, Korea) and Korean National Institute of Health (Seoul, Korea) and were grown in Luria-Bertani (LB) medium containing 1% tryptone (Becton Dickinson, Sparks,

MD, USA), 0.5% yeast extract (Becton Dickinson), and 1% sodium chloride (Sigma, St. Louis, MO, USA). The non-*Salmonella* strains used in this study are listed in Table 2. *Listeria* strains were obtained from the National Veterinary Research and Quarantine Service (Anyang, Korea) and were grown in TSBY containing Tryptic soy broth and 0.6% yeast extract (Becton Dickinson, Sparks, MD, USA). The *Escherichia coli* (*E. coli*) strains, except *E. coli* O157 H:7 ATCC43888 and *E. coli* K12 KCTC1041, were from the National Veterinary Research and Quarantine Service (Anyang, Korea). *E. coli* O157 H:7 ATCC 43888 and *Yersinia enterocolitica* ATCC 27729 were obtained from the Korean National Institute of Health (Seoul, Korea). *E. coli* K12 KCTC 1041 (ATCC 23736), *Shigella flexneri* ATCC 9199 (KCTC 2008), and *Shigella sonnei* ATCC 9290 (KCTC 2009) were obtained from the Korea Research Institute of Bioscience and Biotechnology (Daejeon, Korea).

Reagents

Restriction enzymes and a Wizard Genomic DNA

Table 1. *Salmonella* spp. used in this study

	Serovar	Serogroup		Serovar	Serogroup
1	<i>S. typhi</i> ATCC ^a 19430	D1	30	<i>S. djugu</i>	C1
2	<i>S. paratyphi</i> ATCC11511	A	31	<i>S. meleagridis</i>	E1
3	<i>S. enteritidis</i> ATCC4931	D1	32	<i>S. hillingdon</i>	D2
4	<i>S. schottmulleri</i> ATCC10719	B	33	<i>S. istanbul</i>	C3
5	<i>S. choleraesuis</i> ATCC13312	C1	34	<i>S. lindenburg</i>	C2
6	<i>S. typhimurium</i> ATCC14028	B	35	<i>S. derby</i>	B
7	<i>S. gallinarum</i> ATCC9184	D1	36	<i>S. mbandaka</i>	C1
8	<i>S. london</i> ATCC8389	E1	37	<i>S. haardt</i>	C3
9	<i>S. enteritidis</i> ATCC13076	D1	38	<i>S. schwarzengrund</i>	B
10	<i>S. schwarzengrund</i> IVK B01177	B	39	<i>S. stanley</i>	B
11	<i>S. eingedi</i> IVK B01183	C1	40	<i>S. virchow</i>	C1
12	<i>S. uppsala</i>	B	41	<i>S. montevideo</i>	C1
13	<i>S. ohio</i>	C1	42	<i>S. weltevreden</i>	E1
14	<i>S. budapest</i>	B	43	<i>S. bareilly</i>	C1
15	<i>S. wien</i>	B	44	<i>S. rissen</i>	C1
16	<i>S. virginia</i>	C3	45	<i>S. senftenberg</i>	E4
17	<i>S. muenchen</i>	C2	46	<i>S. bardo</i>	C3
18	<i>S. cerro</i>	K	47	<i>S. anatum</i>	E1
19	<i>S. nigeria</i>	C1	48	<i>S. richmond</i>	C1
20	<i>S. blegdam</i>	D1	49	<i>S. uganda</i>	E1
21	<i>S. yeerongpilly</i>	E1	50	<i>S. ahmadi</i>	E4
22	<i>S. give</i>	E1	51	<i>S. dublin</i>	D1
23	<i>S. hadar</i>	C2	52	<i>S. hindmarsh</i>	C3
24	<i>S. essen</i>	B	53	<i>S. pakistan</i>	C3
25	<i>S. ruanda</i>		54	<i>S. eingedi</i>	C1
26	<i>S. nieukerk</i>	C4	55	<i>S. braenderup</i>	C1
27	<i>S. benin</i>	D2	56	<i>S. infantis</i>	C1
28	<i>S. sereman</i>	D1	57	<i>S. heidelberg</i>	B
29	<i>S. kentucky</i>	C3			

^a ATCC: American Type Culture Collection, Manassas, VA

Table 2. Non-*Salmonella* strains used in this study

Species	Strains	Serotypes ^a	Isolation ^b
<i>Listeria monocytogenes</i>	ATCC ^c 19113	3	human
<i>Listeria monocytogenes</i>	ATCC19114	4a	
<i>Listeria monocytogenes</i>	ATCC19115	4b	human
<i>Listeria monocytogenes</i>	ATCC19117	4d	sheep
<i>Listeria monocytogenes</i>	ATCC19118	4e	chicken
<i>Listeria monocytogenes</i>	HPB410	1/2a	
<i>Listeria monocytogenes</i>	ATCC35152		guinea pig
<i>Listeria ivanivii</i>	ATCC19119		sheep
<i>Listeria innocua</i>	ATCC33090	6a	cow brain
<i>Listeria welshimeri</i>	ATCC35897	6b	decaying plant
<i>Listeria seeligeri</i>	ATCC35967		soil
<i>Listeria grayi</i>	ATCC19120		chinchilla feces
<i>Listeria murrayi</i>	ATCC25401		corn stalks and leaves
<i>Escherichia coli</i> O26			
<i>Escherichia coli</i> O55			
<i>Escherichia coli</i> O111			
<i>Escherichia coli</i> O114			
<i>Escherichia coli</i> O119			
<i>Escherichia coli</i> O157:H7	ATCC43888		
<i>Escherichia coli</i> O157:H7	ATCC43894		
<i>Escherichia coli</i> K12	ATCC23736		
<i>Shigella flexeri</i>	ATCC9199		
<i>Shigella sonnei</i>	ATCC9120		
<i>Yersinia enterocolitica</i>	ATCC27729		

^a Source: <http://www.atcc.org/> except *L. monocytogenes* HPB410.

^b Source: <http://www.atcc.org/>.

purification system were purchased from Promega (USA). The *Taq* DNA polymerase and reagents for the PCR and G-

spinTM genomic DNA extraction kit were obtained from Intron Biotechnology (Korea). Ligase was obtained from Roche (Germany). Pork meat (for cutlet) was purchased at a supermarket (Samsung TESCO Home Plus, Gyeongju, Korea), and was tested before use as negative for the *Salmonella* with PCR. All chemicals, unless otherwise noted, were purchased from Sigma (USA).

Primers

The DNA primers used in this study are listed in Table 3. All primers were purchased from GenoTech (Daejeon, Korea). The *invA* gene (GenBank accession number AL513382) was chosen because this sequence was previously tested for detection of *Salmonella*^{13,22,31-37}.

Construction of the plasmid encoding *Salmonella invA* gene containing an 82- base deletion

All DNA manipulations were performed according to standard procedures³⁸. The *EcoRV* site in the *invA* gene was created by overlap extension. The DG187/DG188 primer pair was used to amplify the DNA segment that contained the *EcoRV* site, together with upstream *EcoRI* sequences for subcloning. The DG189/DG190 primer pair was used to amplify the DNA segment that contained the *EcoRV* site together with downstream *HindIII* sequences for subcloning. The amplified DNAs were purified with a PCRquick-spinTM PCR Product Purification kit (Intron) and were used for amplification with the DG187/DG190 pair. The amplified DNA segment was cleaved with *EcoRI/HindIII*, and was cloned into the pGEM-4Z plasmid to get the pGEM-4Z

Table 3. Primers used in this study

Primer Name	Oligonucleotide sequence (5'-3')	Use
DG75	GAC CGC AAG GTT GAA ACT CA	Detection of <i>Listeria</i> 16S rDNA
DG76	CAG CCT ACA ATC CGA ACT GA	
DG129	TGG GAA ACT GCC TGA TGG AG	Detection of <i>Salmonella</i> 16S rDNA
DG130	ACC TTC CTC CCC GCT GAA AG	
DG131	GCT GGT CTG AGA GGA TGA CC	Detection of <i>E. coli</i> , <i>Yersinia</i> , and <i>Shigella</i> 16S rDNA
DG132	AAG GGC ACA ACC TCC AAG TC	
139	GTG AAA TTA TCG CCA CGT TCG GGC AA	Detection of <i>invA</i>
141	TCA TCG CAC CGT CAA AGG AAC C	
DG158	CGG ATC TCA TTA ATC AAC AAT A	Detection of <i>invA</i>
DG159	CTC TTT CGT CTG GCA TTA TC	
DG187	AAA TAC CGG CAG CGG GAA TTC CGG CAG AGT	Creation of <i>EcoRV</i> site in <i>invA</i>
DG188	GAT TTG AAG GCC GAT ATC ATT GAT GCG GAT	
DG189	ATC CGC ATC AAT GAT ATC GGC CTT CAA ATC	
DG190	CCT CAG TTT TTC AAG CTT TCC TGC GGT ACT	
DG215	AGG TTC AGA TCG CGA CGC GGA AGT	
DG216	ACT TCC GCG TCG CGA TCT GAA CCT	
DG217	TAT AGA ATA CGA ATT CCG GCA GAG T	
DG218	ATA GGG AGA CAA GCT TTC CTG CGG T	
DG218	ATA GGG AGA CAA GCT TTC CTG CGG T	

invA RV. The *NruI* site in the pGEM-4Z *invA* RV was also created by overlap extension. The DG217/DG215 primer pair was used to amplify the pGEM-4Z *invA* RV DNA segment that contained the *NruI* site together with upstream *EcoRI* sequences for subcloning. The DG216/DG218 primer pair was used to amplify the pGEM-4Z *invA* RV DNA segment that contained the *NruI* site downstream *HindIII* sequences for subcloning. The amplified DNAs were purified with the PCRquick-spin™ PCR Product Purification kit, and were used for amplification with the DG217/DG218 pair. The amplified DNA segment was cleaved with *EcoRI/HindIII*, and was cloned into the pGEM-4Z plasmid to get the pGEM-4Z *invA* RV/*NruI*. The pGEM-4Z *invA* RV/*NruI* was cleaved with *EcoRV/NruI*, ligated, and introduced into *E. coli* DH5 α . The colonies were screened for the absence of the *EcoRV* and *NruI* sites by restriction enzyme digestion. After sequencing, the final clone, which had 82 bases deleted, was designated as pGEM-4Z *invA* Δ RV-*NruI*.

Preparation of DNA in artificially inoculated pork meat

The pork meat was homogenized with a Stomacher (AES Laboratoire, France) in distilled water (1:1, w/v) dispensed in bags, and stored at -70°C until use. The guanidine thiocyanate-phenol-chloroform method was used for the preparation of the PCR template³⁹. Briefly, the artificially inoculated 0.1 g of pork meat was extracted with 0.25 mL of solution D (4M guanidine thiocyanate, 0.025M sodium citrate, 0.5% N-Lauroylsarcosine) and 0.5 mL of phenol-chloroform (1:1). The aqueous phase was extracted with 400 μ L of chloroform and the DNA was precipitated with isopropanol and sodium acetate. The resulting pellet was washed once with 70% ethanol, dried, and used for PCR.

Polymerase chain reaction (PCR)

The PCR mixture contained 10 mM Tris-HCl (pH 8.3),

50 mM KCl, 1.5 mM MgCl₂, 200 μ M dNTPs, 2.5 units of *i-StarTaq*™ DNA polymerase, 100 pmol of each primer, and varying amounts of template DNA. The samples were denatured at 94°C for 5 min and subjected to amplification cycles in a thermocycler (My Cycler™ Thermal Cycler, BIO-RAD). Each cycle consisted of a 45-sec denaturation step (94°C), 45-sec annealing step (55°C), and 45-sec extension step (72°C). Finally, the products were extended for 7 min at the completion of 30 amplification cycles.

Results and Discussion

Detection of *Salmonella* spp.

Previously, Rahn et al.²² reported that the detection sensitivity of the *invA* gene was approximately 300 cells per PCR reaction, and Malony et al.⁴⁰ reported 5-50 *Salmonella* colony forming units (cfu) per PCR reaction when they used the boiling method for *Salmonella* DNA extraction. Because the boiling method is usually applied to wash fluid from the surface of a food, it is not adequate to apply it to the direct extraction of *Salmonella* DNA from contaminated meat. Therefore, we decided to use other methods to extract DNA from the meat and compared the modified guanidine thiocyanate-phenol-chloroform method with two commercial kits (G-spin™ genomic DNA extraction kit and Wizard Genomic DNA purification system) for extracting the *Salmonella* DNA (Fig. 1). The modified guanidine thiocyanate-phenol-chloroform method (58-580 cfu) and G-spin™ genomic DNA extraction kit (14.6-146 cfu) were better than the Wizard Genomic DNA purification system (146-1,460 cfu) in detection sensitivity per PCR reaction with primers DG158/DG159 based on the *invA* gene. However, considering the sample size and difficulty in the application to pork meat, the following experiments were performed using the modified guanidine thiocyanate-phenol-chloroform method.




Fig. 1. Gel electrophoresis profile with three template DNA isolation methods. For PCR, the DG158/DG159 primer set was used and 1,031-bp PCR products were compared. The colony-forming units (cfu) of *S. typhimurium* ATCC14028 were determined by MPN. (A) Wizard Genomic DNA purification system (B) Modified guanidine thiocyanate-phenol-chloroform (C) G-spin™ Genomic DNA extraction kit (for bacteria). (A) and (C): Lane M, 100-bp DNA ladder; lane 1, 1.46×10^6 cfu; lane 2, 1.46×10^5 cfu; lane 3, 1.46×10^4 cfu; lane 4, 1.46×10^3 cfu; lane 5, 1.46×10^2 cfu; lane 6, 1.46×10^1 cfu; lane 7, 1.46 cfu (B): Lane M, 100-bp DNA ladder; lane 1, 0.58×10^6 cfu; lane 2, 0.58×10^5 cfu; lane 3, 0.58×10^4 cfu; lane 4, 0.58×10^3 cfu; lane 5, 0.58×10^2 cfu; lane 6, 0.58×10^1 cfu; lane 7, 0.58 cfu.




Fig. 2. Detection sensitivity of *Salmonella* in pork meat. *Salmonella* DNA was extracted using the modified guanidine-thiocyanate-phenol-chloroform method. The cfu of *S. typhimurium* ATCC14028 were determined by MPN. PCR products of 1,031-bp were detected from 1.46×10^6 , 1.46×10^5 , 1.46×10^4 , and 1.46×10^3 cfu of *S. typhimurium* per 0.1 g of pork meat (lanes 1-4). No band was detected from 1.46×10^2 , 1.46×10^1 and 1.46 cfu of *S. typhimurium* per 0.1 g of pork meat (lanes 5-7).

To measure the detection sensitivity in pork meat, homogenized pork meat (0.1 g) was artificially inoculated with 1.46 - 1.46×10^6 cfu of *Salmonella typhimurium* ATCC14028. The DNA was directly extracted with the modified guanidine thiocyanate-phenol-chloroform method and subjected to PCR amplification with primers DG158/DG159. The detection sensitivity was 146-1,460 cfu/0.1 g of pork meat (Fig. 2), which was similar to, or better than, that observed previously; that is, 10^3 cfu of *L. monocytogenes* in 0.5 mL of milk²⁶⁾, 10^3 cfu of *Yersinia enterocolitica* in 0.5 mL of milk²⁸⁾, 860 cfu of *L. monocytogenes* in 0.1 g of pork meat²⁹⁾, 10^3 cfu *Salmonella* spp. per 1 mL milk³⁵⁾ and 17,000 cfu *Salmonella typhimurium* per 0.15 g of wheat grain³⁴⁾. The cycle extension up to 50 cycles, or another 30 cycles of PCR, did not increase the sensitivity. After a 20-fold dilution with LB medium and incubation at 30°C for 15 hr, improved sensitivity was achieved. Five cfu of *S. typhimurium* per 0.1 g of meat was easily detected (data not shown).

Rahn et al.²²⁾ used the primer set 139/141 and amplified the 284-bp *Salmonella invA* specific DNA fragment. Recently, a *Salmonella* spp.-specific conventional PCR assay targeting the *invA* gene was validated in an international collaborative study, and was proposed as an international standard⁴⁰⁾. To evaluate the specificity, we tested 57 *Salmonella* strains comprised of 11 genogroups³⁹⁾, and 24 non-*Salmonella* strains that included eight *E. coli* strains, 13 *Listeria* strains, two *Shigella* strains, and one *Yersinia enterocolitica*. The assay showed a 284-bp *Salmonella* specific band in all 57 *Salmonella* strains, but no false positive reaction occurred in the non-*Salmonella* strains tested (Fig. 3). However, *E. coli*

O119 showed an approximate 450-bp non-specific PCR product (Fig. 3A lane 5). The band was consistent and strong. Rahn et al.²²⁾ mentioned that they also observed this on at least one occasion, in 6 strains of *E. coli*, although they didn't mention the antigen type of *E. coli*, and stated that the non-specific product was obtained inconsistently and showed a weak band. Rahn et al.²²⁾ also mentioned that *S. senftenberg* and *S. litchfield* gave false negative results, whereas Malony et al.⁴⁰⁾ reported that *S. senftenberg* and *S. litchfield* gave positive bands when they were compared using the same primer set 139/141. Our repeated experiments clearly demonstrated that *S. senftenberg* produced a positive *Salmonella* specific 284-bp band (Fig. 3C, lane 45). However, we could not test *S. litchfield* due to the nonavailability of bacteria. At present, it is not clear why this different finding arose. One possible explanation might be due to differences in the PCR conditions between laboratories.

Competitive PCR for the direct enumeration of *Salmonella* spp. in pork meat

Previously we demonstrated that the amplification of radioisotope-labeled *hlyA* of *L. monocytogenes* sequences was still in the exponential range after 30 cycles under the PCR conditions of the study²⁶⁾. Therefore, we used the same conditions for cPCR. For the quantification of *Salmonella* spp., DNA from pork meat that had been artificially inoculated with 1.46×10^4 cfu of *S. typhimurium* ATCC14028 was co-amplified in the presence of known copy numbers of a competitor plasmid, pGEM-4Z *invA* Δ RV-NruI, which binds the same primers (primers 139 and 141) but has been deleted by 82-bp in order to distinguish it from natural sequences. The amplification efficiencies of the two different kinds of DNA should have been almost equal, because the same primers bind the same sequence for amplification. The only difference between the *invA* gene of the bacterial chromosome and the modified gene of pGEM-4Z *invA* Δ RV-NruI was the deleted sequences (82-bp). Since the relative amounts of the cPCR products generated from the amplification reflected the relative initial levels of the two different kinds of DNA, the cfu value of the *Salmonella* spp. was easily estimated by comparing the intensity of the two bands after electrophoresis. In fact, the calculated DNA copy number of the standard DNA and the cfu count deduced from the MPN were almost equal in this procedure using primers 139/141 (Fig. 4).

Several papers have described PCR detection of *Salmonella* after enrichment^{13,19-21,32,34,35,41-44)}, however, reports for direct enumeration of *Salmonella* in food or environmental samples using PCR are rare³⁴⁾. Even though some recent papers have described the automated detection and enumeration of *Salmonella* spp. in composted biosolids³⁶⁾,




Fig. 3. Agarose gel electrophoresis of PCR products obtained from *E. coli*, *Shigella*, *Yersinia*, and *Salmonella* spp. For the detection of *invA*, the DG146/DG147 primer set was used and a 284-bp PCR product is shown. (A) For the detection of 16S rDNA, a DG131/DG132 primer set was used. Lane 1, *E. coli* O26; lane 2, *E. coli* O55; lane 3, *E. coli* O111; lane 4, *E. coli* O114; lane 5, *E. coli* O119; lane 6, *E. coli* O157:H7 ATCC43888; lane 7, *E. coli* O157:H7 ATCC43894; lane 8, *E. coli* K12; lane 9, *Shigella flexneri* ATCC9199; lane 10, *Shigella sonnei* ATCC9120; lane 11, *Y. enterocolitica* ATCC27729; lane 12, *S. typhi* ATCC19430; lane 13, *S. paratyphi* ATCC11511; lane 14, *S. enteritidis* ATCC4391; lane 15, *S. schottmulleri* ATCC10719; lane 16, *S. choleraesuis* ATCC13312. (B) For the detection of the 16S rDNA of *Listeria* spp., the DG75/DG76 primer set was used. Lane 1, *L. monocytogenes* ATCC19113; lane 2, *L. monocytogenes* ATCC19114; lane 3, *L. monocytogenes* ATCC19115; lane 4, *L. monocytogenes* ATCC19117; lane 5, *L. monocytogenes* ATCC19118; lane 6, *L. monocytogenes* (HPB410); lane 7, *L. monocytogenes* ATCC35152; lane 8, *L. ivanovii* ATCC19119; lane 9, *L. innocua* ATCC33090; lane 10, *L. welshimeri* ATCC35897; lane 11, *L. seeligeri* ATCC35967; lane 12, *L. grayi* ATCC19120; lane 13, *L. murrayi* ATCC25401; lane 14, *S. typhimurium* ATCC14028; lane 15, *S. gallinarium* ATCC9184; lane 16, *S. london* ATCC8389. (C) The numbers on each lane are the *Salmonella* strain numbers given in Table 1.

waste water from municipal waste water treatment³⁷), meat⁴⁵), vegetable rinse-water⁴⁶), human feces⁴⁷), and environmental samples^{35,48}) using real-time PCR, the procedure still requires an expensive, special machine and reagents. Depending on the purpose, sometimes simple detection and rough estimation of the cfu are enough for routine control. The procedure described in this study does not provide accurate numbers of contaminated *Salmonella* spp. in pork meat. Nevertheless, it will be useful for rapid enumeration of *Salmonella* spp. under conditions when real-time PCR is not available.

Since the procedure described in this paper is basically PCR-based, it does not differentiate dead cells from viable ones and can thereby yield exaggerated results. A recent paper described the specific detection of viable *Salmonella* spp. using magnetic capture hybridization RT-PCR in seeded soil and chicken manure⁴⁹). Experiments using RT-cPCR to determine viable *Salmonella* spp. in pork meat are ongoing.

In summary, we evaluated that the *invA* gene is unique to *Salmonella* spp., and have shown that primers based on this region can be used to differentiate *Salmonella* and other organisms. We cloned and modified the *invA* gene and




Fig. 4. Agarose gel electrophoresis after cPCR. The upper band (284-bp) was derived from artificially inoculated *S. typhimurium* (ATCC14028) DNA and the lower band (202-bp) was derived from pGEM-4Z *invA* Δ RV-NruI standard DNA. 1.46×10^4 cfu of *S. typhimurium* were artificially inoculated in 0.1g of pork meat. Lane M, 100-bp DNA ladder as a size marker; lane 1, 1.46×10^7 copies of standard DNA (pGEM-4Z *invA* Δ RV-NruI); lane 2, 1.46×10^6 copies of standard DNA; lane 3, 1.46×10^5 copies of standard DNA; lane 4, 1.46×10^4 copies of standard DNA; lane 5, 1.46×10^3 copies of standard DNA; lane 6, 1.46×10^2 copies of standard DNA; lane 7, 1.46×10 copies of standard DNA.

successfully used it with cPCR for the rapid enumeration of *Salmonella* spp. in contaminated pork meat. The results of this study provide the basis for conventional PCR as a rapid, inexpensive, and convenient method for the detection and enumeration of *Salmonella* spp. in a practical field such as HACCP.

요 약

돼지고기에 오염된 살모넬라의 수를 직접 측정하는 방법으로 경쟁적 PCR(competitive PCR)을 사용하였다. 세 가지 DNA추출방법을 비교한 후 guanidine thiocyanate-phenol-chloroform 방법을 일부 수정하여 인위적으로 살모넬라를 오염시킨 돼지고기로부터 DNA를 직접 추출하는 방법으로 선택하였다. Rahn 등(Mol. Cell. Probes 1992년)이 이미 보고한 284-bp *invA* gene의 특이성을 평가하기 위해 살모넬라 57균주와 살모넬라가 아닌 균주 24개를 사용하였다. 시험해 본 모든 살모넬라 균주는 *invA* 양성으로 살모넬라가 아닌 모든 균주는 *invA* 음성으로 판명되었으며 살모넬라가 아닌 균주 중 양성으로 잘못 판명된 경우는 없었다. 돼지고기 0.1 g을 사용할 경우 검출한계는 1,460 cfu였다. 경쟁적 PCR을 위해 *invA* gene 중 82-bp를 결실시킨 DNA조각을 pGEM-4Z백터에 클로닝을 하였다. 인위적으로 오염된 돼지고기로부터 추출한 살모넬라 염색체 DNA와 이와 같은 primer 결합자리를 가진 경쟁 DNA

를 동시에 경쟁적 PCR로 증폭하였다. 경쟁적 PCR을 사용하여 결정한 균수와 MPN방법으로 결정한 균수는 거의 유사하였으며 전체 과정을 수행하는 데 약 5시간이 소요되었다.

Acknowledgements

This study was supported by the Dongguk University. We would like to thank Hyojung Lee and Hyungkun Lim for their technical assistance.

References

- Olsen, J.E., Aabo, S., Hill, W., Notermans, S., Wernars, K., Granum, P.E., Popovic, T., Rasmussen, H.N. and Olsvik, O.: Probes and polymerase chain reaction for the detection of food-borne bacterial pathogens. *International J. Food Microbiol.* **28**, 1-78 (1995).
- Hanai, K., Satake, M., Naakanishi, H. and Venkateswaran, K.: Comparison of commercially available kits for detection of *Salmonella* strains in foods. *Appl. Environ. Microbiol.* **63**, 775-778 (1997).
- Hoorfar, J., Baggesen, D.L. and Porting, P.H.: A PCR-based strategy for simple and rapid identification of rough presumptive *Salmonella* isolates. *J. Microbiol. Methods* **35**, 77-84 (1999).
- Andrews, W.H., Poelma, P.L. and Wilson, C.R.: Isolation and identification of *Salmonella* species. In *Bacterial Analytical manual*, 6th Ed. (US Food and Drug Administration) Association of Official Analytical Chemists, Arlington, VA (1984).
- Van Leusden, F.M., Van Schothorst, M. and Beckers, H.J.: The standard *Salmonella* isolation method. In *Isolation and Identification Methods for Food Poisoning Organisms* (Corry, J.E.L., Roberts, D., and Skinner, F.A.) Academic Press, London, pp. 35-49 (1982).
- Zhu, Q., Lim, C.K. and Chan, Y.N.: Detection of *Salmonella typhi* by polymerase chain reaction. *J. Appl. Bacteriol.* **80**, 244-251 (1996).
- Aznar, E., Alarcon, T., Buendia, Garcia-Penuela and Lopez-Brea, M.: Detection of decreased susceptibility to fluoroquinolones in *Salmonella* spp. by five different methods including real-time PCR. *International J. Antimicrobial Agents* **30**, 67-71 (2007).
- Baumler, A.J., Heffron, F. and Reissbrodt, R.: Rapid detection of *Salmonella enterica* primers specific for *iroB*. *J. Clin. Microbiol.* **35**, 1224-1230 (1997).
- Botteldoorn, N., Coillie, E.V., Grijspeerd, K., Werbrouck, H., haesebrouck, F., Donn, E., D'Haese, E., Heyndrickx, M., Pasmans, F. and Herman, L.: Real-time reverse transcription PCR for the quantification of the *mntH* expression of *Salmonella enterica* as a function of growth phase and phagosomal-like conditions. *J. Microbiol. methods* **66**, 125-135 (2006).
- Cano, R.J., Ramussen, S.R., Fraga, G.S. and Palomares, J.C.:

- Fluorescent detection-polymerase chain reaction (FD-PCR) assay on microwell plates as a screening test for *Salmonellas* in foods. *J. Appl. Bacteriol.* **75**, 247-253 (1993).
11. Chevrier, D., Popoff, M.Y., Dion, M.P., Hermant, D. and Guesdon, J.L.: Rapid detection of *Salmonella* subspecies I by PCR combined with nonradioactive hybridization using covalently immobilized oligonucleotide on a microplate. *FEMS Immunol. Med. Microbiol.* **10**, 245-252 (1995).
 12. Cohen, H.J., Mechanda, S.M. and Lin, W.: PCR amplification of the *fimA* gene sequence of *Salmonella* spp. *Appl. Environ. Microbiol.* **12**, 4303-4308 (1996).
 13. Cortez, A.L.L., Carvalho, A.C.F.B., Ikuno, A.A. and Vidal-Martins, A.M.C.: identification of *Salmonella* spp. isolates from chicken abattoirs by multiplex-PCR. *Research in Veterinary Science* **81**, 340-344 (2006).
 14. Doran, J.L., Collinson, S.K., Buria, J., Sarlos, G., Todd, E.C.D., Munro, C.K., Kay, C.M., Banser, P.A., Peterkin, P.I. and Kay, W.W.: DNA-based diagnostic tests for *Salmonella* species targeting *agfA*, the structural gene for thin, aggregative fimbriae. *J. Clin. Microbiol.* **31**, 2263-2273 (1993).
 15. Fluit, A.C., Widjojatmodjo, M.N., Box, A.T.A., Torensma, R. and Verhoef, J.: Rapid detection of *Salmonellae* in poultry with the magnetic immunopolymerase chain reaction assay. *Appl. Environ. microbiol.* **59**, 1342-1346 (1993).
 16. Hashimoto, Y., Itho, Y., Fuginaga, Y., Khan, A.Q., Sultana, F., Miyake, m., Horose, K., Yamamoto, H. and Ezaki, T.: Development of nested PCR based on the *viaB* sequence to detect *Salmonella typhi*. *J. Clin. Microbiol.* **33**, 775-777 (1995).
 17. Kwang, J., Littledike, E.T. and Keen, J.E.: Use of the polymerase chain reaction for *Salmonella* detection. *Lett. Appl. Microbiol.* **22**, 46-51 (1996).
 18. Mahon, J. and Lax, A.J.: A quantitative polymerase chain reaction method for the detection in avian feces of *Salmonella* carrying the *spvR* gene. *Epidemiol. Infect.* **111**, 455-464 (1993).
 19. Malorny, B., Bunge, C. and Helmuth, R.: A real-time PCR for the detection of *Salmonella* Enteritidis in poultry meat and consumption eggs. *J. Microbiol. methods* **70**, 245-251 (2007).
 20. Murphy, N.M., McLauchlin, J., Ohai, C. and Grant K.A.: Construction and evaluation of a microbiological positive process internal control for PCR-based examination of food samples for *Listeria monocytogenes* and *Salmonella enterica*. *International J. Food Microbiol.* doi:10.1016/j.ijfoodmicro.2007.06.006.
 21. Patel, J.R., Bhagwat, A.A., Sanglay, G.C. and Solomon, M.B.: Rapid detection of *Salmonella* from hydrodynamic pressured-treated poultry using molecular beacon real-time PCR. *Food Microbiol.* **23**, 39-46 (2006).
 22. Rahn, K., De Grandis, S.A., Clarke, R.C., McEwen, S.A., Galan, J.E., Ginocchio, C., Curtiss III, R. and Gyles, C.L.: Amplification of an *invA* gene sequence of *Salmonella typhimurium* by polymerase chain reaction as a specific method of detection of *Salmonella*. *Mol. Cell. Probes* **6**, 271-279 (1992).
 23. Rexach, L., Dilassier, F. and Fach, P.: Polymerase chain reaction for *Salmonella* virulence-associated plasmid genes: a new tool in *Salmonella* epidemiology. *Epidemiol. Infect.* **112**, 33-43 (1994).
 24. Way, J.S., Josephson, K.L., Pillai, S.D., Abbaszadegan, M., Gerba, C.P. and Pepper, I.L.: Specific detection of *Salmonella* spp. by multiplex polymerase chain reaction. *Appl. Environ. Microbiol.* **59**, 1473-1479 (1993).
 25. Yeh, K.-S.Chen, T.-H., Liao, C.-W. and Lo, H.-C.: PCR amplification of the *Salmonella typhimurium fimY* gene sequence to detect the *Salmonella* species. *International J. Food Microbiol.* **78**, 227-234 (2002).
 26. Choi, W.S. and Hong C.-H.: Rapid enumeration of *Listeria monocytogenes* in milk using competitive PCR. *International J. Food Microbiol.* **84**, 79-85 (2003).
 27. Choi, W.S.: Detection of *Salmonella* in milk by polymerase chain reaction. *J. Food Hyg. Safety* **15**, 262-266 (2000).
 28. Choi, W.S.: Detection of *Yersinia enterocolitica* in milk by polymerase chain reaction. *Food Sci. Biotechnol.* **10**, 451-454 (2001).
 29. Lim, H., Hong, C.-H. and Choi, W.S.: Rapid enumeration of *Listeria monocytogenes* in pork meat using competitive PCR. *Food Sci. Biotechnol.* **14**, 387-391 (2005).
 30. Collazo, C.M. and Galn, J.F.: The invasion-associated type III protein secretion system in *Salmonella*.-a review. *Gene* **192**, 51-59 (1997).
 31. Galn, J.E. and Curtis III, R.: Distribution of the *invA*, -B, -C, and -D genes of *Salmonella typhimurium* among other *Salmonella* serovars: *invA* mutants of *Salmonella typhi* are deficient for entry into mammalian cells. *Infect. Immun.* **59**, 2901-2908 (1991).
 32. Hein, I., Flekna, G., Krassnig, M. and Wagner, M.: Real-time PCR for the detection of *Salmonella* spp. in food: An alternative approach to a conventional PCR system suggested by the FOOD-PCR project. *J. Microbiol. Methods* **66**, 538-547 (2006).
 33. Isogai, E., Makungu, C., Yabe, J., Sinkala, P., Nambota, A., Isogai, H., Fukushi, H., Silungwe, M., Mubita, C., Syakalima, M., Hang'ombe, M., Kozaki, S. and Yasuda, J.: Detection of *Salmonella invA* by isothermal and chimeric primer-initiated amplification of nucleic acids (ICAN) in Zambia. *Comparative Immunology, Microbiology & Infectious Diseases* **28**, 363-370 (2005).
 34. Kim, J., Demeke, T., Clear, R.M. and Patrick, S.K.: Simultaneous detection by PCR of *Escherichia coli*, *Listeria monocytogenes* and *Salmonella typhimurium* in artificially inoculated wheat grain. *International J. Food Microbiol.* **111**, 21-25 (2006).
 35. Nam, H.-M., Srinivasan, V., Gillespie, B.E., Murinda, S.E. and Oliver, S.P.: Application of SYBR green real-time PCR assay for specific detection of *Salmonella* spp. in dairy farm environmental samples. *International J. Food Microbiol.* **102**, 161-171 (2005).
 36. Novinscak, A., Surette, C. and Fillion, M.: Quantification of

- Salmonella* spp. in composted biosolids using a TaqMan qPCR assay. *J. Microbiol. Methods* **70**, 119-126 (2007).
37. Shannon, K.E., Lee, D.-Y., Trevors, J.T. and Beaudette, L.A.: Application of real-time quantitative PCR for the detection of selected bacterial pathogens during municipal wastewater treatment. *Science of the Total Environment* **382**, 121-129 (2007).
 38. Sambrook, J. and Russel, D.W.: *Molecular Cloning: A Laboratory Manual*, 3rd Ed. Cold Spring Harbor Press, Cold Spring Harbor, NY, USA (2001).
 39. Lim H., Lee, K.H., Hong, C.-H., Bahk, G.-J. and Choi, W.S.: Comparison of four molecular typing methods for the differentiation of *Salmonella* spp. *International J. Food Microbiol.* **105**, 411-418 (2005).
 40. Malorny, B., Hoorfar, J., Bunge, C. and Helmuth, R.: Multi-center validation of the analytical accuracy of *Salmonella* PCR: towards an international standard. *Appl. Environ. Microbiol.* **69**, 290-296 (2003).
 41. Li, Y., Zhuang, S. and Mustapha, A.: Application of a multiplex PCR for the simultaneous detection of *Escherichia coli* O157:H7, *Salmonella* and *Shigella* in raw and ready-to-eat meat products. *Meat Science* **71**, 402-406 (2005).
 42. Myint, M.S., Johnson, Y.J., Tablante, N.L. and Heckert, R.A.: The effect of pre-enrichment on the sensitivity and specificity of PCR for detection of naturally contaminated *Salmonella* in raw poultry compared to conventional culture. *Food Microbiol.* **23**, 599-604 (2006).
 43. Nowak, B., von Mffling, T., Chaunchom, S. and Hartung, J.: *Salmonella* contamination in pigs at slaughter and on the farm: A field study using an antibody ELISA test and a PCR technique. *International J. Food Microbiol.* **115**, 259-267 (2007).
 44. Wang, X., Jothikumar, N. and Griffiths, M.W.: Enrichment and DNA extraction protocols for the simultaneous detection of *Salmonella* and *Listeria monocytogenes* in raw sausage meat with multiplex real-time PCR. *J. Food Prot.* **67**, 189-192 (2004).
 45. Ellingson, J.L.E., Anderson, J.L., and Carlson, S.A. and Sharma, V.K.: Twelve hour real-time PCR technique for the sensitive and specific detection of *Salmonella* in raw and ready-to-eat meat products. *Mol. Cell. Probes* **18**, 51-57 (2004).
 46. Bhagwat, A.A.: Rapid detection of *Salmonella* from vegetable rinse-water using real-time PCR. *Food Microbiol.* **21**, 73-78 (2004).
 47. Farrell, J.J., Doyle, L.J., Addison, R.M., Reller, L.B., Hall, G.S. and Procop, G.W.: Broad-range (Pan) *Salmonella* and *Salmonella* serotype typhi-specific real-time PCR assays-potential tools for the clinical microbiologist. *Am. J. Clin. Pathol.* **123**, 339-345 (2005).
 48. Klerks, M.M., Van Bruggen, A.H.C., Zijlstra, C. and Donnikov, M.: Comparison of methods of extraction for *Salmonella enterica* Serovar *enteritidis* DNA from environmental substrates and quantification of organism by using a general internal procedural control. *Appl. Environ. Microbiol.* **72**, 3879-3886 (2006).
 49. Jacobsen, C.S. and Holben, W.E.: Quantification of mRNA in *Salmonella* sp. seeded soil and chicken manure using magnetic capture hybridization RT-PCR. *J. Microbiol. Methods* **69**, 315-321 (2007).