

SPECTRAL AREA ESTIMATES FOR NORMS OF COMMUTATORS

MUNEO CHŌ * AND TAKAHIKO NAKAZI **

ABSTRACT. Let A and B be commuting bounded linear operators on a Hilbert space. In this paper, we study spectral area estimates for norms of $A^*B - BA^*$ when A is subnormal or p -hyponormal.

1. Introduction

Let \mathcal{H} be a Hilbert space and $\mathcal{B}(\mathcal{H})$ the set of all bounded linear operators on \mathcal{H} . If T is a hyponormal operator in $\mathcal{B}(\mathcal{H})$ then C. R. Putnam [7] proved that $\|T^*T - TT^*\| \leq \text{Area}(\sigma(T))/\pi$, where $\sigma(T)$ is the spectrum of T . The second named author [5] has proved that if T is a hyponormal operator and K is in $\mathcal{B}(\mathcal{H})$ with $KT = TK$ then

$$\|T^*K - KT^*\| \leq 2\{\text{Area}(\sigma(T))/\pi\}^{1/2}\|K\|.$$

We don't know whether the constant 2 in the inequality is best possible for a hyponormal operator. In §2, we show that the constant is not best possible for a subnormal operator.

When T is a p -hyponormal operator in $\mathcal{B}(\mathcal{H})$, A. Uchiyama [10] generalized the Putnam inequality, that is,

$$\|T^*T - TT^*\| \leq \phi\left(\frac{1}{p}\right) \|T\|^{2(1-p)} \{\text{Area}(\sigma(T))/\pi\}^p.$$

This inequality gives the Putnam inequality when $p = 1$. In §3, we generalize the above inequality for the spectral area estimate of $\|T^*K - KT^*\|$ when $TK = KT$. H. Alexander [1] proved the following inequality for a uniform algebra A . If f is in A then

$$\text{dist}(\bar{f}, A) \leq \{\text{Area}(\sigma(f))/\pi\}^{1/2}.$$

Received January 25, 2006.

2000 *Mathematics Subject Classification*. Primary 47A20.

Key words and phrases. subnormal, p -hyponormal, Putnam inequality.

* This research is partially supported by Grant-in-Aid Scientific Research No.17540139.

** This research is partially supported by Grant-in-Aid Scientific Research No.17540176.

The second named author [5] gave an operator version for the Alexander inequality. This was used in order to estimate $\|T^*K - KT^*\|$ when T is a hyponormal operator and $KT = TK$. We also give an Alexander inequality for a p -hyponormal and we use it to estimate $\|T^*K - KT^*\|$.

In §4, we try to estimate $\|T^*K - KT^*\|$ for arbitrary contraction. In §5, we show a few results about area estimates for p -quasihyponormal operators, restricted shifts and analytic Toeplitz operators.

For $0 < p \leq 1$, T is said to be p -hyponormal if $(T^*T)^p - (TT^*)^p \geq 0$. A 1-hyponormal operator is hyponormal. For an algebra \mathcal{A} in $\mathcal{B}(\mathcal{H})$, let $\text{lat}\mathcal{A}$ be the lattice of all \mathcal{A} -invariant projections. For a compact subset X in \mathcal{C} , $\text{rat}(X)$ denotes the set of all rational functions on X .

2. Subnormal operators

In order to prove Theorem 1, we use the original Alexander inequality.

Theorem 1. *Let T be a subnormal operator in $\mathcal{B}(\mathcal{H})$ and f a rational function on $\sigma(T)$ whose poles are not on it. Then*

$$\|T^*f(T) - f(T)T^*\| \leq \{\text{Area}(\sigma(T))/\pi\}^{1/2}\{\text{Area}(\sigma(f(T)))/\pi\}^{1/2}.$$

Proof. Suppose that $N \in \mathcal{B}(\mathcal{K})$ is a normal extension of $T \in \mathcal{B}(\mathcal{H})$ and P is an orthogonal projection from \mathcal{K} to \mathcal{H} . Then $T = PN \upharpoonright \mathcal{H}$ and so

$$\begin{aligned} & T^*f(T) - f(T)T^* \\ &= PN^*Pf(N)P - Pf(N)PN^*P \\ &= PN^*f(N)P - Pf(N)PN^*P \\ &= Pf(N)N^*P - Pf(N)PN^*P \\ &= Pf(N)(1 - P)N^*P \\ &= Pf(N)(1 - P) \cdot (1 - P)N^*P \end{aligned}$$

because $f(N)P = Pf(N)P$ and $f(N)N^* = N^*f(N)$.

Let F be a rational function in $\text{rat}(\sigma(T))$. Put \mathcal{B}_F = the norm closure of $\{g(F(N)) ; g \in \text{rat}(\sigma(F(N)))\}$ then P belongs to $\text{lat}\mathcal{B}_F$. Hence

$$\begin{aligned} & \|(1 - P)F(N)^*P\| \\ & \leq \text{dist}(F(N)^*, \mathcal{B}_F) \leq \text{dist}(\bar{z}, \text{rat}(\sigma(F(N)))) \\ & \leq \{\text{Area}(\sigma(F(N)))/\pi\}^{1/2} \end{aligned}$$

by the Alexander's theorem [1]. Hence, applying F to $F = z$ or $F = f$

$$\begin{aligned} & \|T^*f(T) - f(T)T^*\| \\ & \leq \|(1 - P)f(N)^*P\| \cdot \|(1 - P)N^*P\| \\ & \leq \{\text{Area}(\sigma(f(N)))/\pi\}^{1/2}\{\text{Area}(\sigma(N))/\pi\}^{1/2} \\ & \leq \{\text{Area}(\sigma(f(T)))/\pi\}^{1/2}\{\text{Area}(\sigma(T))/\pi\}^{1/2}. \end{aligned}$$

□

If T is a cyclic subnormal operator and $KT = TK$ then using a theorem of T. Yoshino [12] we can prove that

$$\|T^*K - KT^*\| \leq \{\text{Area}(\sigma(T))/\pi\}^{1/2} \{\text{Area}(\sigma(K))/\pi\}^{1/2}.$$

The proof is almost same to one of Theorem 1.

3. p -hyponormal operators

In order to prove Theorem 2, we use an operator version of the Alexander inequality for a p -hyponormal operator. Unfortunately Lemma 3 is not best possible for $p = 1$ (see [5]). Lemma 1 is due to W. Arveson [2, Lemma 2] and Lemma 2 is due to A. Uchiyama [11, Theorem 3].

We need the following notation to give Theorem 2 and Proposition 1. Let ϕ be a positive function on $(0, \infty)$ such that

$$\phi(t) = \begin{cases} t & \text{if } t \text{ is an integer} \\ t + 2 & \text{if } t \text{ is not an integer.} \end{cases}$$

We write $\ell^2 \otimes \mathcal{H}$ for the Hilbert space direct sum $\mathcal{H} \oplus \mathcal{H} \oplus \cdots$, and $1 \otimes T$ denotes the operator $T \oplus T \oplus \cdots \in \mathcal{B}(\ell^2 \otimes \mathcal{H})$ for each operator $T \in \mathcal{B}(\mathcal{H})$.

Lemma 1. *Let \mathcal{A} be an arbitrary ultra-weakly closed subalgebra of $\mathcal{B}(\mathcal{H})$ containing 1, and let $T \in \mathcal{B}(\mathcal{H})$. Then*

$$\text{dist}(T, \mathcal{A}) = \sup\{\|(1 - P)(1 \otimes T)P\| ; P \in \text{lat}(1 \otimes \mathcal{A})\}.$$

Lemma 2. *If T is a p -hyponormal operator, then*

$$\|T^*T - TT^*\| \leq \phi\left(\frac{1}{p}\right) \|T\|^{2(1-p)} \{\text{Area}(\sigma(T))/\pi\}^p.$$

Lemma 3. *If T is a p -hyponormal operator then*

$$\text{dist}(T^*, \mathcal{A}) \leq \sqrt{2\phi\left(\frac{1}{p}\right)} \|T\|^{1-p} \{\text{Area}(\sigma(T))/\pi\}^{p/2},$$

where \mathcal{A} is the strong closure of $\{f(T) ; f \in \text{rat}(\sigma(T))\}$.

Proof. Let $S = 1 \otimes T$. Then S is p -hyponormal. In order to prove the lemma, by Lemma 1 it is enough to estimate $\sup\{\|(1 - P)SP\| ; P \in \text{lat}(1 \otimes \mathcal{A})\}$. If $P \in \text{lat}(1 \otimes \mathcal{A})$ then $SP = PSP$ and so

$$\begin{aligned} & \|(1 - P)SP\|^2 \\ &= \|PSS^*P - PSPS^*P\| \\ &= \|PSS^*P - PS^*SP + PS^*SP - PSPS^*P\| \\ &\leq \|P(S^*S - SS^*)P\| + \|(PSP)^*(PSP) - (PSP)(PSP)^*\| \\ &\leq \|S^*S - SS^*\| + \|(PSP)^*(PSP) - (PSP)(PSP)^*\|. \end{aligned}$$

By [11, Lemma 4], PSP is p -hyponormal and so by Lemma 2 we have

$$\begin{aligned} & \|PSS^*P - PSPS^*P\|^2 \\ & \leq \phi\left(\frac{1}{p}\right) \|T\|^{2(1-p)} \{\text{Area}(\sigma(T))/\pi\}^p \\ & \quad + \phi\left(\frac{1}{p}\right) \|PSP\|^{2(1-p)} \{\text{Area}(\sigma(PSP))/\pi\}^p \\ & \leq 2\phi\left(\frac{1}{p}\right) \|T\|^{2(1-p)} \{\text{Area}(\sigma(T))/\pi\}^p \end{aligned}$$

because $\|PSP\| \leq \|S\| = \|T\|$ and $\sigma(PSP) \subset \sigma(S) = \sigma(T)$. By Lemma 1,

$$\text{dist}(T^*, \mathcal{A}) \leq \sqrt{2\phi\left(\frac{1}{p}\right) \|T\|^{1-p} \{\text{Area}(\sigma(T))/\pi\}^{p/2}}.$$

□

Theorem 2. *If T is a p -hyponormal operator in $\mathcal{B}(\mathcal{H})$ and if K is in $\mathcal{B}(\mathcal{H})$ with $KT = TK$, then*

$$\|T^*K - KT^*\| \leq 2\sqrt{2\phi\left(\frac{1}{p}\right) \|T\|^{1-p} \{\text{Area}(\sigma(T))/\pi\}^{p/2}} \|K\|.$$

Proof. When \mathcal{A} is the strong closure of $\{f(T) ; f \in \text{rat}(\sigma(T))\}$, for any $A \in \mathcal{A}$

$$\|T^*K - KT^*\| = \|(T^* - A)K + AK - KT^*\| \leq 2\|T^* - A\| \|K\|.$$

Now Lemma 3 implies the theorem. □

In Theorem 2, if $p = 1$, that is, T is hyponormal then $\|T^*K - KT^*\| \leq 2\sqrt{2}\{\text{Area}(\sigma(T))/2\}^{1/2}\|K\|$. The constant $2\sqrt{2}$ is not best because the second author [5] proved that $\|T^*K - KT^*\| \leq 2\{\text{Area}(\sigma(T))/2\}^{1/2}\|K\|$. If $p = \frac{1}{2}$, that is, T is semi-hyponormal then

$$\|T^*K - KT^*\| \leq 4\|T\|^{1/2} \{\text{Area}(\sigma(T))/\pi\}^{1/4} \|K\|.$$

4. Norm estimates

In general, it is easy to see that $\|T^*T - TT^*\| \leq \|T\|^2$. By Theorem 1, if T is subnormal and f is an analytic polynomial then

$$\|T^*f(T) - f(T)T^*\| \leq \|T\| \|f(T)\|.$$

In this section, we will prove that $\|T^*T^n - T^nT^*\| \leq \|T\|^{n+1}$ for arbitrary T in $\mathcal{B}(\mathcal{H})$.

Theorem 3. *If T is a contraction on \mathcal{H} and f is an analytic function on the closed unit disc \bar{D} then $\|T^*f(T) - f(T)T^*\| \leq \sup_{z \in \bar{D}} |f(z)|$.*

Proof. By a theorem of Sz.-Nagy [6], there exists a unitary operator U on \mathcal{K} such that \mathcal{K} is a Hilbert space with $\mathcal{K} \supseteq \mathcal{H}$ and $T^n = PU^n|_{\mathcal{K}}$ for $n \geq 0$, where P is an orthogonal projection from \mathcal{K} to \mathcal{H} . Then it is known that $U^*P = PU^*P$ and $f(T) = Pf(U)|_{\mathcal{H}}$. Hence

$$\begin{aligned} & T^*f(T) - f(T)T^* \\ &= PU^*Pf(U)P - Pf(U)PU^*P \\ &= PU^*Pf(U)P - Pf(U)U^*P \\ &= PU^*(I - P)f(U)P \end{aligned}$$

because $U^*P = PU^*P$ and $f(U)U^* = U^*f(U)$. Therefore

$$\begin{aligned} & \|T^*f(T) - f(T)T^*\| \\ &= \|PU^*(I - P)f(U)P\| \leq \sup_{z \in D} |f(z)|. \end{aligned}$$

□

Corollary 1. *If T is in $\mathcal{B}(\mathcal{H})$ then for any $n \geq 1$ $\|T^*T^n - T^nT^*\| \leq \|T\|^{n+1}$.*

Proof. Put $A = T/\|T\|$ then A is a contraction and so by Theorem 2 $\|A^*A^n - A^nA^*\| \leq 1$ and so $\|T^*T^n - T^nT^*\| \leq \|T\|^{n+1}$. □

5. Remarks

In this section, we give spectral area estimates for p -quasihyponormal operators, restricted shifts and analytic Toeplitz operators.

For $0 < p \leq 1$, T is said to be p -quasihyponormal if $T^*\{(T^*T)^p - (TT^*)^p\}T \geq 0$. A 1-quasihyponormal operator is called quasihyponormal.

Lemma 4. *Let T be p -quasihyponormal and P be a projection such that $TP = PTP$. Then PTP is also p -quasihyponormal.*

Proof. Since T is p -quasihyponormal, $T^*(T^*T)^pT \geq T^*(TT^*)^pT$. Hence, we have

$$PT^*(T^*T)^pTP \geq PT^*(TT^*)^pTP.$$

Since by the Hansen's inequality [4]

$$\begin{aligned} & PT^*(T^*T)^pTP = (PTP)^*P(T^*T)^pP(PTP) \\ &\leq (PTP)^*(PT^*TP)^p(PTP) \\ &= (PTP)^*\{(PTP)^*(PTP)\}^p(PTP) \end{aligned}$$

and by $0 < p < 1$

$$\begin{aligned} & PT^*(TT^*)^pTP \geq (PT^*P)(TPT^*)^p(PTP) \\ &= (PTP)^*\{(PTP)(PTP)^*\}^p(PTP), \end{aligned}$$

we have

$$(PTP)^*\{(PTP)^*(PTP)\}^p \geq (PTP)^*\{(PTP)(PTP)^*\}^p(PTP).$$

Hence, PTP is p -quasihyponormal. □

Proposition 1. *If T is a p -quasihyponormal operator in $\mathcal{B}(\mathcal{H})$ and if K is in $\mathcal{B}(\mathcal{H})$ with $KT = TK$, then*

$$\|T^*K - KT^*\| \leq 4 \left[\phi \left(\frac{1}{p} \right) \right]^{1/4} \|T\|^{1-p/2} \{\text{Area}(\sigma(T))/\pi\}^{p/4} \|K\|.$$

In particular, if T is quasihyponormal then

$$\|T^*K - KT^*\| \leq 4\|T\|^{1/2} \{\text{Area}(\sigma(T))/\pi\}^{1/4} \|K\|.$$

Proof. We can prove it as in the proof of Theorem 2. By [11, Theorem 6], $\|T^*T - TT^*\| \leq 2\|T\|^{2-p} \sqrt{\phi(\frac{1}{p})} \{\text{Area}(\sigma(T))/\pi\}^{p/2}$. Hence by Lemma 4

$$\text{dist}(T^*, \mathcal{A}) \leq 2\|T\|^{1-\frac{p}{2}} \phi \left(\frac{1}{p} \right)^{\frac{1}{4}} \{\text{Area}(\sigma(T))/\pi\}^{p/4}.$$

This implies the proposition. \square

Let H^2 and H^∞ be the usual Hardy spaces on the unit circle and z the coordinate function. M denotes an invariant subspace of H^2 under the multiplication by z . By the well known Beurling theorem, $M = qH^2$ for some inner function. Suppose N is the orthogonal complement of M in H^2 . For a function ϕ in H^∞ , S_ϕ is an operator on N such that $S_\phi f = P(\phi f)$ ($f \in N$), where P is the orthogonal projection from H^2 to N . For a symbol ϕ in L^∞ , T_ϕ denotes the usual Toeplitz operator on H^2 .

Proposition 2. *Suppose $\Phi = q\bar{\phi}$ belongs to H^∞ . Then*

- (1) $\|S_\phi^* S_\phi - S_\phi S_\phi^*\| \leq \text{Area}(\overline{\Phi(D)})/\pi$;
- (2) $\|S_\phi^* S_\phi^n - S_\phi^n S_\phi^*\| \leq \{\text{Area}(\overline{\Phi(D)})/\pi\}^{n+1}$ for $n \geq 0$.

Proof. By a well known theorem of Sarason [8],

$$\|S_\phi\| = \|\phi + qH^\infty\| = \|\bar{q}\phi + H^\infty\| = \|\bar{\Phi} + H^\infty\|.$$

By Nehari's theorem [6], $\|\bar{\Phi} + H^\infty\| = \|H_{\bar{\Phi}}\|$, where $H_{\bar{\Phi}}$ denotes a Hankel operator from H^2 to $\bar{z}H^2$. Since $\|H_{\bar{\Phi}}\|^2 = \|T_\Phi^* T_\Phi - T_\Phi T_\Phi^*\|$, where T_Φ denotes a Toeplitz operator on H^2 , by the Putnam inequality

$$\|T_\Phi^* T_\Phi - T_\Phi T_\Phi^*\| \leq \text{Area}(\sigma(T_\Phi))/\pi = \text{Area}(\overline{\Phi(D)})/\pi.$$

Now since $\|S_\phi^* S_\phi - S_\phi S_\phi^*\| \leq \|S_\phi\|^2$, (1) follows. (2) is also clear by the proof above and Corollary 1. \square

Proposition 3. *Suppose f and g are in H^∞ . Then*

$$\|T_f^* T_g - T_g T_f^*\| \leq \{\text{Area}(\overline{f(D)})/\pi\}^{1/2} \{\text{Area}(\overline{g(D)})/\pi\}^{1/2}.$$

Proof. It is easy to see that $T_f^*T_g - T_gT_f^* = H_{\bar{g}}^*H_{\bar{f}}$. Hence

$$\|T_f^*T_g - T_gT_f^*\| \leq \|H_{\bar{g}}\| \cdot \|H_{\bar{f}}\|.$$

Since $H_{\bar{f}}^*H_{\bar{f}} = T_f^*T_f - T_fT_f^*$, by the Putnam inequality

$$\|T_f^*T_g - T_gT_f^*\| \leq \{\text{Area}(\overline{f(D)})/\pi\}^{1/2}\{\text{Area}(\overline{g(D)})/\pi\}^{1/2}.$$

□

References

- [1] H. Alexander, *Projections of polynomial hulls*, J. Funct. Anal. **13** (1973), 13–19.
- [2] W. Arveson, *Interpolation problems in nest algebras*, J. Funct. Anal. **20** (1975), no. 3, 208–233.
- [3] M. Chō and M. Itoh, *Putnam's inequality for p -hyponormal operators*, Proc. Amer. Math. Soc. **123** (1995), no. 8, 2435–2440.
- [4] F. Hansen, *An operator inequality*, Math. Ann. **246** (1979/80), no. 3, 249–250.
- [5] T. Nakazi, *Complete spectral area estimates and self-commutators*, Michigan Math. J. **35** (1988), no. 3, 435–441.
- [6] Z. Nehari, *On bounded bilinear forms*, Ann. of Math. (2) **65** (1957), 153–162.
- [7] C. R. Putnam, *An inequality for the area of hyponormal spectra*, Math. Z. **116** (1970), 323–330.
- [8] D. Sarason, *Generalized interpolation in H^∞* , Trans. Amer. Math. Soc. **127** (1967), 179–203.
- [9] B. Sz.-Nagy and C. Foias, *Harmonic Analysis Of Operators On Hilbert Space*, American Elsevier, New York, 1970.
- [10] A. Uchiyama, *Berger-Shaw's theorem for p -hyponormal operators*, Integral Equations Operator Theory **33** (1999), 221–230.
- [11] ———, *Inequalities of Putnam and Berger-Shaw for p -quasihyponormal operators*, Integral Equations Operator Theory **34** (1999), no. 2, 91–106.
- [12] T. Yoshino, *Subnormal operators with a cyclic vector*, Tohoku Math. (2) **21** (1969), 47–55.

MUNEO CHō
 DEPARTMENT OF MATHEMATICS
 KANAGAWA UNIVERSITY
 JAPAN
E-mail address: chiyom01@kanagawa-u.ac.jp

TAKAHIKO NAKAZI
 DEPARTMENT OF MATHEMATICS
 FACULTY OF SCIENCE
 HOKKAIDO UNIVERSITY
 SAPPORO 060-0810, JAPAN
E-mail address: nakazi@math.sci.hokudai.ac.jp