



DdeI Polymorphism in Coding Region of Goat *POU1F1* Gene and Its Association with Production Traits

X. Y. Lan^{1,3,a}, C. Y. Pan^{1,a}, H. Chen^{1,2,*}, C. Z. Lei¹, L. S. Hua¹, X. B. Yang¹, G. Y. Qiu¹
R. F. Zhang¹ and Y. Z. Lun¹

¹ College of Animal Science and Technology, Northwest A & F University,

Shaanxi Key Laboratory of Molecular Biology for Agriculture, Yangling, Shaanxi 712100, P. R. China

ABSTRACT : *POU1F1* is a positive regulator for GH, PRL and TSH β and its mutations associate with production traits in ruminant animals. We described a *DdeI* PCR-RFLP method for detecting a silent allele in the goat *POU1F1* gene: TCT (241Ser)>TCG (241Ser). Frequencies of D₁ allele varied from 0.600 to 1.000 in Chinese 801 goats. Significant associations of *DdeI* polymorphism with production traits were found in milk yield (*p<0.05), litter size (*p<0.05) and one-year-old weight (*p<0.05) between different genotypes. Individuals with genotype D₁D₁ had a superior performances when compared to those with genotype D₁D₂ (*p<0.05). Hence, the *POU1F1* gene was suggested to the potential candidate gene for superior milk performance, reproduction trait and weight trait. Genotype D₁D₁, characterized by a *DdeI* PCR-RFLP detection, was recommended to geneticists and breeders as a molecular marker for better performance in the goat industry. (**Key Words :** Goat, *POU1F1* Gene, Polymorphism, Association)

INTRODUCTION

The goat industry is an important part among the so-called big domestic animals sector in China. The estimated size is more than 157,361,000 mainly reared in the northern China, which belonged to more than twenty native breeds (e.g. dairy, meat and wool breeds). Dairy-goat farming is significant to the economics of the western China with the characteristics of under-development and poverty. Goat's meat from young or adult animals has been consumed throughout the country for recent ten years. Moreover, the wool of goat is used in many ways, e.g. for the wrapping of the dead, the making of clothes (Boyazoglu et al., 2005). Large exports of wool and its related products give great chance for rural and western families to improve the economic situation (Dubeuf et al., 2004). Therefore, the

further improvement and increase of the quantity and quality in goat dairy, meat and wool will better contribute to the Chinese society, particularly in economy, nutrition, tradition and religion. This issue can be resolved by culturing more and better goat breeds. However, it is difficult to culture excellent goat breeds by the traditional genetic and breeding method. So, many breeders mainly focus on DNA markers for animal selection and breeding through marker-assisted selection (MAS).

As a member of the POU-domain family gene, *POU1F1* is a positive regulator for growth hormone (GH) (Zhou et al., 2005), prolactin (PRL) (Li et al., 2006) and thyroid-stimulating hormone β (TSH β) by binding to target DNA promoters as a dimer in mammalian animals (Jacobson et al., 1997). *POU1F1* mutations associated with Snell dwarf (dw) and Jackson dwarf (dw-J) in mice and dwarfism in human (Li et al., 1990; Pfaffle et al., 1992; Reynau et al., 2004). Moreover, polymorphisms of *POU1F1* gene associated with important production traits in cattle (Renaville et al., 1997a; Renaville et al., 1997b; Zhao et al., 2004) and in pig (Yu et al., 1995; Stancekov et al., 1999; Sun et al., 2002). Recently polymorphisms of sheep *POU1F1* gene were firstly reported (Bastos et al., 2006). Few polymorphisms of goat *POU1F1* gene and their associations with production traits had been described.

* Corresponding Author: H. Chen. Tel: +86-29-87091379, Fax: +86-29-87092164, E-mail: chen hong1212@263.net

² Institute of Cellular and Molecular Biology, Xuzhou Normal University, Xuzhou, Jiangsu 221116, P. R. China.

³ Institute of Biochemistry, College of Life Sciences, Zhejiang Sci-Tech University, Hangzhou, Zhejiang 310018, China.

¹ Medical College, Dalian University, Dalian, Liaoning 115522, P. R. China.

^a These authors contributed equally to this paper.

Received August 1, 2006, Accepted April 7, 2007

Table 1. The primer sequences and their information of goat *POU1F1* gene

Gene	The primer sequences	Size	Annealing temperature	Location
<i>POU1F1</i> gene	Forward: 5'-CCATCATCTCCCTTCTT -3'	450 bp	54.5°C	exon 6 and partial intron 5, 3' UTR
	Reverse: 5'-AATGTACAATGTGCCTTCTGAG-3'			

Therefore, it was an interesting and important work to study polymorphisms in *POU1F1* gene and their associations with production traits in goat, which could provide useful genetic markers for animal selection and breeding through marker-assisted selection (MAS). In this paper, we reported the identification of *DdeI* polymorphism at coding region of goat *POU1F1* gene and evaluated its effects on production traits.

MATERIALS AND METHODS

DNA samples

Genomic DNA samples were obtained from 801 goats belonging to nine genetic populations: Inner Mongolia White Cashmere (IMWC, 452), Xinong sannen dairy (Sa, 74), Laoshan dairy (LS, 80), Guanzhong dairy (GZ, 62), Guizhou Black (GB, 21), Matou (MT, 22), Banjiao (BJ, 25), Guizhou White (GW, 31), and Leizhou goat (LZ, 34), which were reared in the province of Inner Mongolia, Shaanxi, Shandong, Guizhou, Hubei, Sichuan and Guangdong (P. R. China). They were all unrelated animals. 1,512 records of milk yield for 216 dairy goats (Sa, LS, GZ), 3,010 records of litter size, 2,004 records of weight traits and one-year-old body sizes for 216 dairy goats (Sa, LS, GZ) and 452 IMWC goats, 4,500 records of lana length, lana thickness and lana yields for 452 IMWC goats were collected, respectively. DNA samples were extracted from leucocytes and ears tissues according to Sambrook et al. (2001).

PCR conditions

Based upon the sequences of sheep *POU1F1* gene (AJ549207) and bovine *POU1F1* gene (Zhao et al., 2004), a pair of primers was designed to amplify the goat *POU1F1* gene (Table 1). The 25 µl volume contained 50 ng genomic DNA, 0.5 µM of each primer, 1×Buffer (including 1.5 mM MgCl₂), 200 µM dNTPs and 0.625 units of *Taq* DNA polymerase (MBI). The cycling protocol was 4 min at 95°C, 35 cycles of denaturing at 94°C for 45 s, annealing at 54.5°C for 45 s, extending at 72°C for 1 min, with a final extension at 72°C for 10 min.

Single stranded conformation polymorphism (SSCP) and sequencing

Aliquots of 5 µl PCR products were mixed with 5 µl denaturing solution (95% formamide, 25 mM EDTA, 0.025% xylene-cyanole and 0.025% bromophenol blue), heated for 10 min at 98°C and chilled on ice. Denatured

DNA was subjected to PAGE (80×73×0.75 mm) in 1×TBE buffer and constant voltage (200 V) for 2.5-3.0 h. The gel was stained with 0.1% silver nitrate (Kim et al., 2005; Hang et al., 2006; Zhou et al., 2006). The seventeen PCR products from different SSCP patterns in different breeds were subcloned to T-vector (Promega) and sequenced in both directions in ABI PRISM 377 DNA sequencer (Perkin-Elmer).

Genotyping of *DdeI* *POU1F1* allele by PCR-RFLP

Aliquots of 20 µl PCR products of *POU1F1* gene were digested with 10 U *DdeI* at 37°C for 5 h. The digested products were detected by 12.0% PAGE electrophoresis and stained with 0.1% silver nitrate (Zhou et al., 2006).

Statistical analysis

The chi-square test was used to analyze the Hardy-Weinberg equilibrium, which was performed by SPSS software (version 13.0). Statistical analysis was performed on the basis of records of production traits in 216 dairy goats and 452 IMWC goats, respectively. All analyses were done in two steps, first using a full animal model and then using a reduced animal model. The full animal model included fixed effects of marker genotype, birth year, season of birth (spring vs. fall), age of dam, sire, farm, sex, breed and random effects of measurements and animal. The reduced model was used in the final analysis. (1) Repeated measurements of the milk yield of 216 dairy goat (74 Sa, 80 LS, 62 GZ) were analyzed by the use of the statistical software SPSS (version 13.0) with the mixed linear model. (2) The adjusted Linear Model I with fixed effects was used to analyze the relationships between genotypes and weight traits and litter size in 668 goats (74 Sa, 62 GZ, 80 LS and 452 IMWC), linear model I: $Y_{ijklm} = \mu + S_i + D_j + A_k + G_l + (SG)_{jl} + E_{ijklm}$, where Y_{ijklm} was the trait measured on each of the $ijklm^{\text{th}}$ animal, μ was the overall population mean, S_i was the fixed effect associated with the i^{th} sire, D_j was the fixed effect associated with j^{th} dam with sire i , A_k was fixed effect due to the k^{th} age, G_l was the fixed effect associated with l^{th} genotype (*POU1F1*/genotype D_1D_1 and D_1D_2), $(SG)_{jl}$ was interaction between the i^{th} sire and the l^{th} genotype and E_{ijklm} was the random error. (3) The adjusted linear model with fixed effects was used to analyze the relationship between genotypes and lana traits in 452 IMWC goats. Lack associated of farm, sex, and season of birth (spring vs. fall) with variability of traits indicated that these factors were not into linear model in IMWC




Figure 1. The structure of goat *POU1F1* gene and SNPs location for exon 6 and the *DdeI* enzyme site. [CTNAG]: the enzyme site of *DdeI* endonuclease;: the forward primer and reverse primer. Shade: the exon 6 of goat *POU1F1* gene.




Figure 2. The DNA electrophoretic patterns on 12.0% PAGE after digestion with *DdeI* endonuclease of the DNA region of the goat *POU1F1* gene. Lane 4: DNA Marker (Tianwei times, China), ladder 100 bp, 200 bp, 300 bp, 400 bp, 500 bp and 600 bp; Lane 1: genotype D₁D₂; Lane 2, 3: genotype D₁D₁. Genotypes D₁D₂ and D₁D₁ had 6 bands (200 bp, 118 bp, 113 bp, 102 bp, 20 bp and 11 bp) and 5 bands (200 bp, 118 bp, 102 bp, 20 bp), respectively. As two small bands (20 bp and 11 bp) were invisible on 12.0% PAGE electrophoresis, only 4 bands (200 bp, 118 bp, 113 bp and 102 bp) and 3 bands (200 bp, 118 bp and 102 bp) were visible for genotypes D₁D₂ and D₁D₁, respectively. Moreover, it was obvious that 113 bp and 102 bp fragments could clearly classify genotypes D₁D₂ and D₁D₁.

populations. The least square means estimates (LSM) with

standard errors and multiple range tests for two *POU1F1* genotypes and production traits were used.

Synonymous codon bias analysis

According to the high homology among bovine, sheep and goat *POU1F1* gene, the exon 6 of goat *POU1F1* gene was analyzed by www.ncbi.nlm.nih.gov/Blastn and www.ebi.ac.uk/clusterw online software. Then, ORF finder was used to identify the amino acid sequence of exon 6 of goat *POU1F1* gene. According to the online software (www.kazusa.or.jp/codon/countcodon.html), the codon frequencies of exon 6 of goat *POU1F1* gene was calculated. Finally, the codon usage frequency was analyzed by the following formula (Kurland et al., 1991; Eyre-Walker et al., 1996; Lavner et al., 2005; Esley et al., 2006; Liu et al., 2006): $F = m \times k / n$, among them, “F” meant codon usage frequency, “m” meant the number of synonymous codon for specific amino acid, “n” meant the number of specific amino acid in analyzed sequences, “k” meant the number of usage for specific synonymous codon.

RESULTS

Exon 6 and its flanking region of goat *POU1F1* locus demonstrated polymorphic patterns in nine populations by

PCR-SSCP. Then seventeen DNA amplification fragments including the exon 6 were sequenced (DQ826397-DQ826413). The comparisons among these sequences revealed four mutations (Figure 1). According to the high homology among bovine, sheep and goat *POUIF1* gene, one DQ826397:g.102T>G mutation in No.60nt of the exon 6 identified a silent allele: 241Ser (TCT)>Ser (TCG) of *POUIF1* protein (291 aa). Interestingly, this mutation can be detected by *DdeI* endonuclease. We named the allele characterized by the presence of T as *POUIF1*-D₁, as well as G for *POUIF1*-D₂ allele. The DQ826397:g.102T>G mutation (CTCAG-to-CGCAG) of the exon 6 region removes a *DdeI* endonuclease restriction site (CTNAG) (Figure 1). Therefore, the amplified DNA fragment with *DdeI* endonuclease digestion showed five fragments (200 bp, 118 bp, 102 bp, 20 bp and 11 bp) for *POUIF1*-D₁ allele and four fragments (200 bp, 118 bp, 113 bp and 20 bp) for *POUIF1*-D₂ allele. Correspondingly, genotype D₁D₁ had five fragments (200 bp, 118 bp, 102 bp, 20 bp and 11 bp) and genotype D₁D₂ had six fragments (200 bp, 118 bp, 113 bp, 102 bp, 20 bp and 11 bp). It was obvious that 113 bp and 102 bp fragments could clearly classify the genotypes by 12.0% PAGE (Figure 2).

Frequencies of *POUIF1*-D₁ allele were 0.875, 0.885,

0.600, 0.847, 1.000, 0.727, 0.920, 0.706 and 0.777 for IMWC, Sa, LS, GZ, GB, MT, BJ, GW, LZ populations reared in China, respectively (Table 2). The genotype distributions of Sa, GZ, GB, MT, BJ populations were in agreement with Hardy-Weinberg equilibrium ($p>0.05$) except LS, GW and LZ populations.

The establishment of relationships between genotype D₁D₁ and D₁D₂ and production traits was attempted (Table 3). Significant statistical results were founded in milk yield ($*p<0.05$), litter size ($*p<0.05$) and one-year-old weight ($*p<0.05$) between genotypes.

The Serine (Ser/S) has 6 synonymous codon (namely, UCU, UCG, UCC, UCA, AGC and AGU). From Table 4, there were 5 Serines in analyzed region. The codon usage frequency for UCU codon meant 2.400, while the codon usage frequency for UCG UCA) meant 0.000, the codon usage frequency (AGC, AGU and UCC) meant 1.200 (Table 5).

DISCUSSION

Goat, sheep and bovine *POUIF1* gene locate in 1q21-22 of chromosomes (Woollard et al., 2000). Chromosome 1 q in ruminant is highly conserved at the gene order and

Table 2. Genotype distribution and allelic frequencies at goat *POUIF1* locus

Breeds	Observed genotypes		Total	Allelic frequencies	
	D ₁ D ₁	D ₁ D ₂		D ₁	D ₂
Inner Mongolia White Cashmere (IMWC)	339	113	452	0.875	0.125
Xinong Sannen dairy (Sa)	57	17	74	0.885	0.115
Laoshan dairy (LS)	16	64	80	0.600	0.400
Guanzhong dairy (GZ)	43	19	62	0.847	0.153
Guizhou Black (GB)	21	0	21	1.000	0.000
Matou (MT)	10	12	22	0.727	0.272
Banjiao (BJ)	21	4	25	0.92	0.080
Guizhou White (GW)	11	20	31	0.706	0.294
Leizhou (LZ)	14	20	34	0.777	0.223

Table 3. Associations of *DdeI* PCR-RFLP with production traits at goat *POUIF1* locus

Production traits	Genotypes of <i>DdeI</i> PCR-RFLP genotyping		p value (two-tailed)
	D ₁ D ₁ (Mean±SE)(number)	D ₁ D ₂ (Mean±SE) (number)	
Milk yields (kg)	574.12 ^b ±13.14 (n = 116)	528.46 ^a ±3.89 (n = 100)	* p<0.05
Litter sizes (lamb)	1.87 ^b ±0.09 (n = 455)	1.39 ^a ±0.15 (n = 213)	* p<0.05
Birth weight (kg)	3.36±0.12 (n = 455)	3.01±0.15 (n = 213)	p>0.05
Nine-month-old weight (kg)	46.93±1.52 (n = 455)	41.00±1.00 (n = 213)	p>0.05
One-year-old weight (kg)	51.70 ^b ±3.21 (n = 455)	45.12 ^a ±1.23 (n = 213)	* p<0.05
One-year-old stature (cm)	68.40±1.32 (n = 116)	66.00±2.08 (n = 100)	p>0.05
One-year-old body size (cm)	78.90±0.77 (n = 116)	76.33±0.88 (n = 100)	p>0.05
One-year-old heart girth (cm)	84.87±1.12 (n = 116)	85.67±1.86 (n = 100)	p>0.05
One-year-old shank girth (cm)	8.31±0.15 (n = 116)	7.67±0.33 (n = 100)	p>0.05
Lana thickness (cm)	5.92±0.17 (n = 339)	5.75±0.17 (n = 113)	p>0.05
Lana length (cm)	18.27±0.61 (n = 339)	18.00±1.32 (n = 113)	p>0.05
Lana yield (g)	604.03±18.44 (n = 339)	549.46±15.03 (n = 113)	p>0.05

^{a,b} Means of traits with different superscripts were significantly different (LSD test, * p<0.05).

Table 4. Codon frequencies of exon 6 of goat *POUIF1* gene

Codon frequency (Number)	Codon frequency (Number)	Codon frequency (Number)	Codon frequency (Number)	Codon frequency (Number)	Codon frequency (Number)
UCU 28.2 (Ser,2)	ACG 0.0(0)	GCU 14.1(1)	UUU 42.3(3)	UAU 14.1(1)	UGU 14.1(1)
UCC 14.1 (Ser,1)	GAC 0.0(0)	GCC 14.1(1)	UUC 0.0(0)	UAC 0.0(0)	UGC 42.3(3)
UCA 0.0 (Ser,0)	AUG 14.1(1)	GCA 0.0(0)	UUA 14.1(1)	UAA 14.1(1)	UGA 0.0(0)
UCG 0.0 (Ser,0)	GUU 14.1(1)	GCG 0.0(0)	UUG 0.0(0)	UAG 14.1(1)	UGG 14.1(1)
AGU 14.1 (Ser,1)	AAG 28.2(2)	CCU 28.2(2)	CUU 14.1(1)	CAU 14.1(1)	CGU 0.0(0)
AGC 14.1 (Ser,1)	GAU 0.0(0)	CCC 0.0(0)	CUC 0.0(0)	CAC 14.1(1)	CGC 0.0(0)
CCA 0.0(0)	AGG 28.2(2)	GUC 0.0(0)	CUA 14.1(1)	CAA 0.0(0)	CGA 14.1(1)
CCG 0.0(0)	GGU 0.0(0)	GUA 0.0(0)	CUG 56.3(4)	CAG 70.4(5)	CGG 14.1(1)
ACU 0.0(0)	GGG 0.0(0)	GUG 42.3(3)	AUU 0.0(0)	AAU 28.2(2)	GAA 84.5(6)
ACC 0.0(0)	AGA 70.4(5)	GGC 0.0(0)	AUC 28.2(2)	AAC 28.2(2)	
ACA 14.1(1)	AAA 42.3(3)	GGA 14.1(1)	AUA 0.0(0)	GAG 56.3(4)	

Table 5. The codon usage frequencies for Serine in the exon 6 of *POUIF1* gene

Synonymous codon	Number of synonymous codon	Number of amino acid (n)	Usage number of codon (k)	Codon usage frequency (F)
UCU	6	5	2	2.400
UCG	6	5	0	0.000
AGC	6	5	1	1.200
AGU	6	5	1	1.200
UCC	6	5	1	1.200
UCA	6	5	0	0.000

cytogenetics levels. Interval mapping to detect QTL revealed significant effects on milk and protein yield associated with chromosome 1 in the region of bovine *POUIF1* (Renaville et al., 1997). Moreover, the *POUIF1* regulates expression of GH, PRL, TSH β gene and itself (Sun et al., 2002). Hence, *POUIF1* gene was considered to have effects on production and will benefit for the goat industry, whose DNA markers will contribute to animal selection and breeding through marker-assisted selection (MAS).

We were aware of few research related to the polymorphisms of goat *POUIF1* gene and their association with production traits. In this paper, four mutations were revealed in the exon 6 of goat *POUIF1* gene by PCR-SSCP and DNA sequencing method. One DQ826397:g.102T>G mutation identified a silent allele (p.S241S) and formed the *DdeI* polymorphism. The frequencies of *POUIF1*-D₁ allele varied from 0.600 to 1.000. Interestingly, we observed that D₂-D₂ genotype which was not detected in genotype analysis. We presumed that the absence of genotype D₂-D₂ associated with "major codon bias". According to the previous papers, the frequencies with which individual synonymous codons were used to code their cognate amino acids was quite variable from genome to genome and within genomes, from gene to gene. One particularly well documented codon bias was that associated with highly expressed genes in bacteria as well as in yeast; this was the so-called major codon bias (Kurland et al., 1991; Lavner et al., 2005; Esley et al., 2006; Liu et al., 2006). As the complete CDS sequence of goat *POUIF1* gene was not available, the exon 6 region was

used to analyze the codon bias. The analysis of codon usage frequency (F) revealed that different synonymous codons for Serine showed codon bias phenomenon. If F (codon) > 2.0, this codon was regarded as "high frequency codon"; while the codon was regarded as "low frequency codon" or "rare codon" if F (codon) = 0.000. Hence, "UCU" was regarded as "high frequency codon", while the "UCG" was called for "rare codon". From Figure 1, the genotype D₁D₁ linked to "T" mutation and complied with the high frequency codon (UCU) which was called for major codon, while the genotype D₂D₂ linked to "G" mutation and complied with the low frequency codon (UCG) which was called for rare codon. Major codon bias (UCU vs. UCG) was not an arrangement for regulating *POUIF1* gene expression. Instead, the similar data suggested that this codon bias, which was correlated with a corresponding bias of tRNA abundance, was a global arrangement for optimizing the growth efficiency of cells (Kurland et al., 1991; Eyre-Walker et al., 1996; Coghlan et al., 2000; Archetti et al., 2004; Lavner et al., 2005; Esley et al., 2006; Liu et al., 2006). Moreover, we presumed that the rare tRNA abundance for UCG may greatly decrease the translation speed from mRNA to amino acid, and seriously restricted the synthesis efficiency of protein and others, then resulted in the absence of D₂-D₂ individual.

The statistical results revealed significant relationships between some traits and genotypes (*p<0.05). The individuals with genotype D₁D₁ had better performance (e.g. milk yield, litter size and weight) than those of the individuals with genotype D₁D₂. Although this silent

mutation didn't change amino acid sequence. it possibly resulted in the change of Serine synonymous codon usage frequency. The DQ826397:g.102T>G mutation changed the frequency of synonymous codon from 2.400 ("UCU", high frequency codon) to 0.000 ("UCG", rare codon). So, we presumed that the codon bias associated with expressed level of *POU1F1*. The genotype D₁D₂ with rare codon (UCG) possibly associated the less expression level of *POU1F1* which regulates expression level of GH, PRL and TSH β gene. thus the genotype D₁D₂ showed junior performance. This presume complied with the previous descriptions (Kurland et al., 1991; Eyre-Walker et al., 1996; Coghlan et al., 2000; Archetti et al., 2004; Lavner et al., 2005; Esley et al., 2006; Liu et al., 2006). Hence, we revealed that *DdeI* polymorphism of coding region of *POU1F1* gene associated with milk performance, reproduction traits and weight traits, which preliminarily implied that *POU1F1* gene has positive effects on them.

In this study, genotype D₁D₂ of *POU1F1* locus characterized by a *DdeI* PCR-RFLP detection was suggested to be molecular marker for junior milk yield, lambs and weight, as well as genotype D₁D₁ for superior performances.

ACKNOWLEDGEMENTS

This work was funded by the National "863" Program of China (No.2006AA10Z197), NSFC (No.30471238), the Natural Science Foundation of Shaanxi province (No.99SM06) and the Outstanding Talents Foundation of Northwest A&F University (No. 01140101). We kindly thank S. Q. Liu and Y. B. Zhang at Sanbei goat-breeding farm (Sanbei country of E'Touke County, Inner Mongolia, P. R. China) for excellent care of White Cashmere goats. We also acknowledge workers at Qianyang, Guanzhong and Lumian goat-breeding farm at Shaanxi and Shandong province (P. R. China) for careful care of dairy goats.

REFERENCES

- Archetti, M. 2004. Codon usage bias and mutation constraints reduce the level of error minimization of the genetic code. *J. Mol. Evol.* 59(2):258-266.
- Bastos, E., I. Santos, I. H. Parentier, J. L. Castrillo, A. Cravador, M. Guedes-Pinto and R. Renaville. 2006. *Ovis aries POU1F1* gene: cloning, characterization and polymorphism analysis. *Genet.* 126:303-314.
- Boyazoglu, J., I. Hatziminaoglou and P. Morand-Fehr. 2005. The role of the goat in society: Past, present and perspectives for the future. *Small Rumin. Res.* 60:13-23.
- Coghlan, A. and K. H. Wolfe. 2000. Relationship of codon bias to mRNA concentration and protein length in *Saccharomyces cerevisiae*. *Yeast*, 16(12):1131-1145.
- Dubeuf, J. P., P. Morand-Fehr, Rubino and R. Situation. 2004. changes and future of goat industry around the world. *Small Rumin. Res.* 51:165-173.
- Esley, M., J. Heizer, Douglas W. Raiford, Michael L. Raymer, Travis E. Doom, Robert V. Miller and Dan E. Krane. 2006. Amino Acid Cost and Codon-Usage Biases in 6 Prokaryotic Genomes: A Whole-Genome Analysis. *Mole. Biol. Evol.* 2399:1670-1680.
- Eyre-Walker, A. 1996. Synonymous codon bias is related to gene length in *Escherichia coli*: selection for translational accuracy? *Mole. Biol. Evol.* 13:864-872.
- Jacobson, E. M., P. Li, A. Leon-del-Pio, M. G. Rosenfeld and A. K. Aggarwa. 1997. Structure of Pit-1 POU domain bound to DNA as dimer: unexpected arrangement and flexibility. *Genes. Dev.* 11:198-212.
- Kim, J. Y., D. H. Yoon, B. L. Park, L. H. Kim, K. J. Na, J. G. Choi, C. Y. Cho, H. K. Lee, E. R. Chung, B. C. Sang, I. J. Cheong, S. J. Oh and H. D. Shin. 2005. Identification of Novel SNPs in Bovine Insulin-like Growth Factor Binding Protein-3 (IGFBP3) Gene. *Asian-Aust. J. Anim. Sci.* 18(1):3-7.
- Kurland, C. G. 1991. Codon bias and gene expression *FEBS Lett.* 285(2):65-69.
- Lavner, Y. and D. Kotlar. 2005. Codon bias as a factor in regulating expression via translation rate in the human genome. *Gene.* 345:127-138.
- Li, J. T., A. H. Wang, P. Chen, H. B. Li, C. S. Zhang and L. X. Du. 2006. Relationship between the Polymorphisms of 5'Regulation Region of Prolactin Gene and Milk Traits in Chinese Holstein Dairy Cows. *Asian-Aust. J. Anim. Sci.* 19(4):459-462.
- Liu, H., X. H. Wang, Y. F. Liu, X. B. Zhao, N. Li and C. X. Wu. 2006. Codon frequency influencing laying performance: Evidence from 3 SNP in chicken prolactin gene, 10(1):196-198 (Abstr.).
- Li, S., III E. B. Crenshaw, E. J. Rawson, D. M. Simmons, L. W. Swanson and M. G. Rosenfeld. 1990. Dwarf locus mutants lacking three pituitary cell types result from mutations in the POU-domain gene Pit-1. *Nature*, 347:528-533.
- Pfaffle, R. W., G. E. DiMattia, J. Parks, M. Brown, J. M. Wit, M. Jansen, N. H. Van der, J. L. Van den Brande, M. G. Rosenfel, H. A. Ingraham. 1992. Mutation of the POU-specific domain of Pit-1 and hypopituitarism without pituitary hypoplasia. *Sci.* 257:1118-1121.
- Renaville, R., N. Gengler, E. Vrench, A. Prandi, S. Massart, C. Corradini, C. Bertozzi, F. Mortiaux, A. Burny and D. Portetelle. 1997a. Pit-1 gene polymorphism, milk yield, and conformation traits for Italian Holstein-Friesian bulls. *J. Dairy Sci.* 80:3431-3438.
- Renaville, R., N. Gengler, I. Parmentier, F. Mortiaux, S. Massart, C. Bertozzi, A. Burny and D. Portetelle. 1997b. Pit-1 gene *HinfI* RFLP and growth traits in double-muscle Belgian Blue Cattle. *J. Anim. Sci.* 75 (Suppl. 1):146 (Abstr.).
- Reynau, R., A. Saveanu, A. Barlier, A. Enjalbert and T. Brue. 2004. Pituitary hormone deficiencies due to transcription factor gene alterations. *Growth Hormone & IGF Research*, 14(6):442-448.
- Sambrook, J. and D. W. Russell. 2001. *Molecular Cloning: A Laboratory Manual*, vol. 3, third ed. Cold Spring Harbor Laboratory Press, New York
- Stancekov, K., D. Vasicek, D. Peskovicov, J. Bull and A. Kubek. 1999. Effect of genetic variability of the porcine pituitary-specific transcription factor (PIT-1) on carcass traits in pigs.

- Anim. Genet. 30:313-315.
- Sun, H. S., L. L. Anderson, T. P. Yu, K. S. Kim, J. Klind and C. K. Tuggle. 2002. Neonatal Meishan pigs show *POU1F1* genotype effects on plasma GH and PRL concentration. Anim. Rep. Sci. 69:223-237.
- Woollard, J., C. K. Tuggle and F. A. Ponce de Leon. 2000. Rapid communication: Localization of *POU1F1* to bovine, ovine, and caprine 1q21-22. J. Anim. Sci. 78:242-243.
- Yu, T. P., C. K. Tuggle, C. B. Schmitz and M. F. Rothschild. 1995. Association of PIT1 polymorphisms with growth and carcass traits in pigs. J. Anim. Sci. 73:1282-1288.
- Zhao, Q., M. E. Davis and H. C. Hines. 2004. Associations of polymorphisms in the Pit-1 gene with growth and carcass traits in Angus beef cattle. J. Anim. Sci. 82(8):2229-2233.
- Zhou, G. L., Q. Zhu, H. G. Jin and S. L. Guo. 2006. Genetic Variation of Growth Hormone Gene and Its Relationship with Milk Production Traits in China Holstein Cows. Asian-Aust. J. Anim. Sci. 19(3):315-318.
- Zhou, L., Y. Y. Yang, Z. H. Li, L. J. Kong, G. D. Xing, H. S. Di and G. L. Wang. 2006. Detection and Characterization of PCR-SSCP Markers of the Bovine Lactoferrin Gene for Clinical Mastitis. Asian-Aust. J. Anim. Sci. 19(10):1399-1403.