

## GENERALIZED STABILITY OF EULER-LAGRANGE TYPE QUADRATIC MAPPINGS

KIL-WOUNG JUN \* AND JEONG-HA OH \*\*

ABSTRACT. In this paper, we investigate the generalized Hyers–Ulam–Rassias stability of the following Euler-Lagrange type quadratic functional equation  $f(ax+by+cz)+f(ax+by-cz)+f(ax-by+cz)+f(ax-by-cz)=4a^2f(x)+4b^2f(y)+4c^2f(z)$ .

### 1. Introduction

In 1940, S. M. Ulam [12] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these was the following question concerning the stability of homomorphisms. *Let  $G$  be a group and let  $G'$  be a metric group with metric  $\rho(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if  $f : G \rightarrow G'$  satisfies  $\rho(f(xy), f(x)f(y)) < \delta$  for all  $x, y \in G$ , then a homomorphism  $h : G \rightarrow G'$  exists with  $\rho(f(x), h(x)) < \epsilon$  for all  $x \in G$ ?*

In 1941, D. H. Hyers [3] considered the case of approximately additive mappings  $f : E \rightarrow E'$ , where  $E$  and  $E'$  are Banach spaces and  $f$  satisfies *Hyers inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon$$

for all  $x, y \in E$ . It was shown that the limit  $L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n}$  exists for all  $x \in E$  and that  $L : E \rightarrow E'$  is the unique additive mapping satisfying

$$\|f(x) - L(x)\| \leq \epsilon.$$

Let  $E_1$  and  $E_2$  be real vector spaces. A function  $f : E_1 \rightarrow E_2$ , there exists a quadratic function if and only if  $f$  is a solution function of the

---

Received October 30, 2007.

2000 Mathematics Subject Classification: Primary 39B62; Secondary 39B82.

Key words and phrases: Euler-Lagrange type quadratic mapping, generalized Hyers–Ulam stability, functional inequality.

This work was supported by the second Brain Korea 21 Project in 2006.

quadratic functional equation

$$(1.1) \quad f(x+y) + f(x-y) = 2f(x) + 2f(y).$$

A stability problem for the quadratic functional equation (1.1) was solved by F. Skof [11] for mapping  $f : E_1 \rightarrow E_2$ , where  $E_1$  is a normed space and  $E_2$  is a Banach space.

In 1978, Th. M. Rassias [7] provided a generalization of Hyers' Theorem which allows the *Cauchy difference to be unbounded*. S. Czerwik [2] proved the Hyers-Ulam-Rassias stability of quadratic functional equation (1.1). Let  $E_1$  and  $E_2$  be a real normed space and a real Banach space, respectively, and let  $p \neq 2$  be a positive constant. If a function  $f : E_1 \rightarrow E_2$  satisfies the inequality

$$\|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p)$$

for some  $\epsilon > 0$  and for all  $x, y \in E_1$ , then there exists a unique quadratic function  $q : E_1 \rightarrow E_2$  such that

$$\|f(x) - q(x)\| \leq \frac{2\epsilon}{|4 - 2^p|} \|x\|^p$$

for all  $x \in G$ . In particular, we note that J.M. Rassias introduced the Euler-Lagrange quadratic mappings, motivated from the following pertinent algebraic equation

$$(1.2) \quad |ax + by|^2 + |bx - ay|^2 = (a^2 + b^2)[|x|^2 + |y|^2].$$

Thus the second author of this paper introduced and investigated the stability problem of Ulam for the relative Euler-Lagrange functional equation

$$(1.3) \quad f(ax + by) + f(bx - ay) = (a^2 + b^2)[f(x) + f(y)]$$

in the publications[8-10].

Recently, S.M. Jung [5] and J. Bae, K. Jun and S. Jung [1] have generalized the equation (1.1) to

$$(1.4) \quad \begin{aligned} f(x+y+z) &+ f(x-y+z) + f(x+y-z) + f(x-y-z) \\ &= 4f(x) + 4f(y) + 4f(z) \end{aligned}$$

and then have investigated the general solution and the stability problem for the functional equation.

Now, we consider the following functional equations

$$(1.5) \quad \begin{aligned} f(ax + by + cz) &+ f(ax + by - cz) \\ &+ f(ax - by + cz) + f(ax - by - cz) \\ &= 4a^2 f(x) + 4b^2 f(y) + 4c^2 f(z), \end{aligned}$$

where  $a, b, c \neq 0$  are real numbers.

In this paper, we will establish the general solution and the generalized Hyers-Ulam-Rassias stability problem for the equation (1.5) in Banach spaces.

## 2. Euler-Lagrange type quadratic mapping in Banach spaces

LEMMA 2.1. *Let  $X$  and  $Y$  be vector spaces. If a mapping  $f : X \rightarrow Y$  satisfies  $f(0) = 0$  and*

$$\begin{aligned} f(ax + by + cz) &+ f(ax + by - cz) \\ (2.1) \quad &+ f(ax - by + cz) + f(ax - by - cz) \\ &= 4a^2 f(x) + 4b^2 f(y) + 4c^2 f(z) \end{aligned}$$

for all  $x, y, z \in X$ , then the mapping  $f$  is quadratic and  $f(\lambda^n x) = \lambda^{2n} f(x)$ , where  $\lambda = a, b$  or  $c$ .

*Proof.* Letting  $x = y$  in (2.1), we get

$$\begin{aligned} f((a+b)x + cz) &+ f((a+b)x - cz) \\ (2.2) \quad &+ f((a-b)x + cz) + f((a-b)x - cz) \\ &= 4a^2 f(x) + 4b^2 f(x) + 4c^2 f(z) \end{aligned}$$

for all  $x, z \in X$ . Setting  $y = -x$  in (2.1), we obtain

$$\begin{aligned} f((a-b)x + cz) &+ f((a-b)x - cz) \\ (2.3) \quad &+ f((a+b)x + cz) + f((a+b)x - cz) \\ &= 4a^2 f(x) + 4b^2 f(-x) + 4c^2 f(z). \end{aligned}$$

By (2.2) and (2.3), we conclude that  $f$  is even. And by setting  $y = 0$  and  $z = 0$  in (2.1), we get  $f(ax) = a^2 f(x)$  for all  $x \in X$ . So, it is easy to verify  $f(a^n x) = a^{2n} f(x)$  by induction. Similarly, we have the identity for  $b$  and  $c$ . Now, substituting 0 for  $z$  in (2.1), one obtains

$$\begin{aligned} f(ax + by) + f(ax - by) &= 2a^2 f(x) + 2b^2 f(y) \\ &= 2f(ax) + 2f(by). \end{aligned}$$

for all  $x, y \in X$ . Hence  $f$  is quadratic.  $\square$

The mapping  $f : X \rightarrow Y$  given in the statement of Lemma 2.1 is called an *Euler-Lagrange type quadratic mapping*. Putting  $z = 0$  in (2.1) with  $a = 1 = b$ , we get the quadratic mapping  $f(x+y) + f(x-y) = 2f(x) + 2f(y)$ .

From now on, Let  $X$  and  $Y$  be a normed vector space and a Banach space, respectively.

For a given mapping  $f : X \rightarrow Y$ , we define

$$\begin{aligned} Df(x, y, z) &:= f(ax + by + cz) + f(ax + by - cz) + f(ax - by + cz) \\ &\quad + f(ax - by - cz) - 4a^2f(x) - 4b^2f(y) - 4c^2f(z) \end{aligned}$$

for all  $x, y, z \in X$

**THEOREM 2.2.** *Let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  for which there exists a function  $\phi : X^3 \rightarrow [0, \infty)$  such that*

$$(2.4) \quad \Phi(x, y, z) := \sum_{j=1}^{\infty} a^{2j} \phi\left(\frac{x}{a^j}, \frac{y}{a^j}, \frac{z}{a^j}\right) < \infty,$$

$$(2.5) \quad \|Df(x, y, z)\| \leq \phi(x, y, z)$$

for all  $x, y, z \in X$ . Then there exists a unique Euler-Lagrange type quadratic mapping  $Q : X \rightarrow Y$  such that  $DQ(x, y, z) = 0$  and

$$(2.6) \quad \|f(x) - Q(x)\| \leq \frac{1}{4a^2} \Phi(x, 0, 0)$$

for all  $x \in X$ .

*Proof.* Letting  $y = 0$  and  $z = 0$  in (2.5), we get

$$\|f(ax) - a^2f(x)\| \leq \frac{1}{4}\phi(x, 0, 0)$$

for all  $x \in X$ . So

$$\left\|f(x) - a^2f\left(\frac{x}{a}\right)\right\| \leq \frac{1}{4}\phi\left(\frac{x}{a}, 0, 0\right)$$

for all  $x \in X$ . Hence

$$\begin{aligned} \left\|a^{2l}f\left(\frac{x}{a^l}\right) - a^{2m}f\left(\frac{x}{a^m}\right)\right\| &\leq \sum_{j=l+1}^m \left\|a^{2(j-1)}f\left(\frac{x}{a^{j-1}}\right) - a^{2j}f\left(\frac{x}{a^j}\right)\right\| \\ (2.7) \quad &\leq \sum_{j=l+1}^m \frac{1}{4}a^{2(j-1)}\phi\left(\frac{x}{a^j}, 0, 0\right) \end{aligned}$$

for all  $x \in X$ . It means that a sequence  $\{a^{2n}f(\frac{x}{a^n})\}$  is Cauchy for all  $x \in X$ . Since  $Y$  is complete, the sequence  $\{a^{2n}f(\frac{x}{a^n})\}$  converges. So one can define a mapping  $Q : X \rightarrow Y$  by  $Q(x) := \lim_{n \rightarrow \infty} a^{2n}f(\frac{x}{a^n})$  for all  $x \in X$ .

By (2.4) and (2.5),

$$\begin{aligned}\|DQ(x, y, z)\| &= \lim_{n \rightarrow \infty} a^{2n} \left\| Df\left(\frac{x}{a^n}, \frac{y}{a^n}, \frac{z}{a^n}\right) \right\| \\ &\leq \lim_{n \rightarrow \infty} a^{2n} \phi\left(\frac{x}{a^n}, \frac{y}{a^n}, \frac{z}{a^n}\right) = 0\end{aligned}$$

for all  $x, y, z \in X$ . So  $DQ(x, y, z) = 0$ . By Lemma 2.1, the mapping  $Q : X \rightarrow Y$  is quadratic.

Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (2.7), we get the approximation (2.6) of  $f$  by  $Q$ .

Now, let  $Q' : X \rightarrow Y$  be another quadratic mapping satisfying (2.6). Then we obtain

$$\begin{aligned}\|Q(x) - Q'(x)\| &= a^{2n} \left\| Q\left(\frac{x}{a^n}\right) - Q'\left(\frac{x}{a^n}\right) \right\| \\ &\leq a^{2n} \left[ \left\| Q\left(\frac{x}{a^n}\right) - f\left(\frac{x}{a^n}\right) \right\| + \left\| Q'\left(\frac{x}{a^n}\right) - f\left(\frac{x}{a^n}\right) \right\| \right] \\ &\leq \frac{1}{2} a^{2(n-1)} \Phi\left(\frac{x}{a^n}, 0, 0\right),\end{aligned}$$

which tends to zero as  $n \rightarrow \infty$ . So we can conclude that  $Q(x) = Q'(x)$  for all  $x \in X$ . This proves the uniqueness of  $Q$ . Hence the mapping  $Q : X \rightarrow Y$  is a unique quadratic mapping satisfying (2.6).  $\square$

**COROLLARY 2.3.** *Let  $p$  and  $\theta$  be positive real numbers such that either  $p > 2$  and  $|a| > 1$  or  $p < 2$  and  $|a| < 1$ , and let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  and*

$$(2.8) \quad \|Df(x, y, z)\| \leq \theta(\|x\|^p + \|y\|^p + \|z\|^p),$$

for all  $x, y, z \in X$ . Then there exists a unique Euler-Lagrange type quadratic mapping  $Q : X \rightarrow Y$  such that

$$(2.9) \quad \|f(x) - Q(x)\| \leq \frac{\theta \cdot \|x\|^p}{4(|a|^p - a^2)}$$

for all  $x \in X$ .

*Proof.* Define  $\phi(x, y, z) = \theta(\|x\|^p + \|y\|^p + \|z\|^p)$ , and apply Theorem 2.2.  $\square$

**THEOREM 2.4.** *Let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  for which there exists a function  $\phi : X^3 \rightarrow [0, \infty)$  such that*

$$(2.10) \quad \Phi(x, y, z) := \sum_{j=0}^{\infty} \frac{1}{a^{2j}} \phi(a^j x, a^j y, a^j z) < \infty,$$

$$(2.11) \quad \|Df(x, y, z)\| \leq \phi(x, y, z)$$

for all  $x, y, z \in X$ . Then there exists a unique Euler-Lagrange type quadratic mapping  $Q : X \rightarrow Y$  such that  $DQ(x, y, z) = 0$  and

$$(2.12) \quad \|f(x) - Q(x)\| \leq \frac{1}{4a^2} \Phi(x, 0, 0)$$

for all  $x \in X$ .

*Proof.* Letting  $y = 0$  and  $z = 0$  in (2.11), we get

$$\|f(ax) - a^2 f(x)\| \leq \frac{1}{4} \phi(x, 0, 0)$$

for all  $x \in X$ . So

$$\left\| f(x) - \frac{1}{a^2} f(ax) \right\| \leq \frac{1}{4a^2} \phi(x, 0, 0)$$

for all  $x \in X$ .

Hence

$$(2.13) \quad \begin{aligned} \left\| \frac{1}{a^{2l}} f(a^l x) - \frac{1}{a^{2m}} f(a^m x) \right\| &\leq \sum_{j=l+1}^m \left\| \frac{1}{a^{2(j-1)}} f(a^{j-1} x) - \frac{1}{a^{2j}} f(a^j x) \right\| \\ &\leq \sum_{j=l+1}^m \frac{1}{4a^{2j}} \phi(a^{j-1} x, 0, 0) \end{aligned}$$

for all  $x \in X$ . It means that a sequence  $\{\frac{1}{a^{2n}} f(a^n x)\}$  is Cauchy for all  $x \in X$ . Since  $Y$  is complete, the sequence  $\{\frac{1}{a^{2n}} f(a^n x)\}$  converges. So one can define a mapping  $Q : X \rightarrow Y$  by  $Q(x) := \lim_{n \rightarrow \infty} \frac{1}{a^{2n}} f(a^n x)$  for all  $x \in X$ .

By (2.10) and (2.11),

$$\begin{aligned} \|DQ(x, y, z)\| &= \lim_{n \rightarrow \infty} \frac{1}{a^{2n}} \|Df(a^n x, a^n y, a^n z)\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{a^{2n}} \phi(a^n x, a^n y, a^n z) = 0 \end{aligned}$$

for all  $x, y, z \in X$ . So  $DQ(x, y, z) = 0$ . By Lemma 2.1, the mapping  $Q : X \rightarrow Y$  is a quadratic.

Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (2.13), we get the approximation (2.12) of  $f$  by  $Q$ .

Now, let  $Q' : X \rightarrow Y$  be another quadratic mapping satisfying (2.12). Then we obtain

$$\begin{aligned} \|Q(x) - Q'(x)\| &= \frac{1}{a^{2n}} \|Q(a^n x) - Q'(a^n x)\| \\ &\leq \frac{1}{a^{2n}} [\|Q(a^n x) - f(a^n x)\| + \|Q'(a^n x) - f(a^n x)\|] \\ &\leq \frac{1}{2a^{2(n+1)}} \Phi(a^n x, 0, 0), \end{aligned}$$

which tends to zero as  $n \rightarrow \infty$ . So we can conclude that  $Q(x) = Q'(x)$  for all  $x \in X$ . This proves the uniqueness of  $Q$ . Hence the mapping  $Q : X \rightarrow Y$  is a unique quadratic mapping satisfying (2.12).  $\square$

**COROLLARY 2.5.** *Let  $p$  and  $\theta$  be positive real numbers with either  $p < 2$  and  $|a| > 1$  or  $p > 2$  and  $|a| < 1$ , and let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  and*

$$(2.14) \quad \|Df(x, y, z)\| \leq \theta(\|x\|^p + \|y\|^p + \|z\|^p),$$

*for all  $x, y, z \in X$ . Then there exists a unique Euler-Lagrange type quadratic mapping  $Q : X \rightarrow Y$  such that*

$$(2.15) \quad \|f(x) - Q(x)\| \leq \frac{\theta \cdot \|x\|^p}{4(a^2 - |a|^p)}$$

*for all  $x \in X$ .*

*Proof.* Define  $\phi(x, y, z) = \theta(\|x\|^p + \|y\|^p + \|z\|^p)$ , and apply Theorem 2.4.  $\square$

**COROLLARY 2.6.** *Let  $f : X \rightarrow Y$  be a mapping satisfying  $f(0) = 0$  for which there exists a nonnegative number  $\theta$  such that*

$$(2.16) \quad \|Df(x, y, z)\| \leq \theta$$

*for all  $x, y, z \in X$ . If  $|a| \neq 1$ , then there exists a unique Euler-Lagrange type quadratic mapping  $Q : X \rightarrow Y$  such that*

$$(2.17) \quad \|f(x) - Q(x)\| \leq \frac{\theta}{4|1 - a^2|}$$

*for all  $x \in X$ .*

*Proof.* Define  $\phi(x, y, z) = \theta$ , and apply Theorem 2.2 and Theorem 2.4.  $\square$

## References

- [1] J. H. Bae, K. W. Jun and S.M. Jung, *On the stability of a quadratic funtional equation*, Kyungpook Math. J. **43** (2003), 415–423.
- [2] S. Czerwik, *On the stability of the mapping in normed spaces*, Abh. Math. Sem. Univ. Hamburg, **62**(1992), 59–64.
- [3] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [4] K. Jun and Y. Lee, *A generalization of the Hyers–Ulam–Rassias stability of Jensen’s equation*, J. Math. Anal. Appl. **238** (1999), 305–315.
- [5] S. M. Jung, *Quadratic functional equations of Perider thpe*, Interant. J. Math. Math. Sci. **24** (2000), 351–359.
- [6] C. Park, W. Park, *On the Jensen’s equation in Banach modules*, Taiqanese J. Math. **6** (2002), 523–300.
- [7] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [8] J. M. Rassisa, *On the stability of the Euler-Lagrange functional equation*, Chinese J. Math. **20** (1992), 185–190.
- [9] J. M. Rassias, *On the stability of the non-linear Euler-Lagrange functional equation in real normed linear spaces*, J. Math. Phys. Sci. **28** (1994), 231–235.
- [10] J. M. Rassias, *On the stability of the general Euler-Lagrange functional equation*, Demonstratio Math. **29** (1996), 755–766.
- [11] F. Skof, *Local properties and approximations of operators(Italian)*, Rend. Sem. Mat. Fis. Milano, **53** (1983), 113–129.
- [12] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ. New York, 1960.

\*

Department of Mathematics  
 Chungnam National University  
 Daejeon, 305–764, Republic of Korea  
*E-mail:* kwjun@@math.cnu.ac.kr

\*\*

Department of Mathematics  
 Chungnam National University  
 Daejeon, 305–764, Republic of Korea  
*E-mail:* jhoh@cnu.ac.kr