

김휘종¹, 김현석², 마정은², 이승준², 함현석², 조유지², 정이영², 전경녀³, 김호철², 이종덕², 황영실²

Clinical Characteristics of Endobronchial Tuberculosis that Develops in Patients over 70 Years of Age

Hwi Jong Kim, M.D.¹, Hyeon Sik Kim, M.D.², Jeong Eun Ma, M.D.², Seung Jun Lee, M.D.², Hyoun Seok Ham, M.D.², Yu Ji Cho, M.D.², Yi Yeong Jeong, M.D.², Kyoung-Nyeo Jeon, M.D.³, Ho Cheol Kim, M.D.², Jong Deok Lee, M.D.², Young Sil Hwang, M.D.²

¹Department of Internal Medicine, Ulsan Hospital, Ulsan, Departments of ²Internal Medicine and ³Diagnostic Radiology, College of Medicine, Gyeongsang National University, Jinju, Korea

Background: The possibility of developing pulmonary tuberculosis usually increases with increasing age. Therefore, the incidence of endobronchial tuberculosis in older people may increase. We evaluated the clinical characteristics in patients with endobronchial tuberculosis above the age of 70 years.

Methods: We enrolled 74 patients (12 males and 62 females; mean age 64.6 ± 16.2 years) that were diagnosed with endobronchial tuberculosis from March 2003 to July 2006 at Gyeongsang University Hospital. We retrospectively evaluated the clinical characteristics of endobronchial tuberculosis for patients 70 years or older (older group) and for patients below the age of 70 years (younger group).

Results: The number of patients in the older group was 41 (55%). Cough was the most common symptom in the two groups of patients and dyspnea on exertion was more common in the older group of patients than in the younger group of patients (31.7% vs. 12.1%). The actively caesating type of disease was more common in the younger group of patients than in the older group of patients (66.7% vs. 39%). The edematous type of disease was more common in the older group of patients than in the younger group of patients (53.7% vs. 27.2%) ($p < 0.05$). Tracheal and main bronchial involvement of lesions were more common for the younger group of patients than for the older group of patients (30.3% vs. 9.7%) ($p < 0.05$).

Conclusion: Endobronchial tuberculosis was commonly observed in patients older than 70 years and this group of patients had some clinical characteristics that were different from the younger group of patients. (*Tuberc Respir Dis* 2007;63:412-416)

Key Words: Endobronchial tuberculosis, Older people, Clinical characteristics

서 론

가

3

가

1.

가

10~40%

2

가

가

4.

Address for correspondence: Ho Cheol Kim, M.D.
Department of Internal Medicine, College of Medicine,
Gyeongsang National University, 92, Chilam-dong, Jinju
660-741, Korea
Phone: 82-55-740-8068, Fax: 82-55-740-8618
E-mail: hochkim@gshp.gsnu.ac.kr

Received: Oct. 12, 2007

Accepted: Oct. 29, 2007

가

가

가

5-8가

70

대상 및 방법

2003 3 2006 6

70

가

1,2

7가

Table 1. Age distribution

Age	Male	Female	Total (%)
<20	1	0	1 (1.4)
20~	0	4	4 (5.4)
30~	0	4	4 (5.4)
40~	2	1	3 (4.2)
50~	3	3	6 (8.1)
60~	2	13	15 (20.2)
70~	3	32	35 (47.2)
>80	1	5	6 (8.1)
Total	12	62	74 (100)

Table 2. The clinical manifestation

Symptoms and signs	Age<70 (n, %)	Age≥70 (n, %)
Cough	20 (60.6)	21 (51.2)
Sputum	12 (36.4)	10 (24.4)
Dyspnea	4 (12.1)	13 (31.7)*
Hemoptysis	3 (9.1)	4 (9.8)
Weakness	2 (6.1)	2 (4.9)
Fever	1 (3)	0 (0)
Hoarseness	1 (3)	0 (0)

*p<0.05 compared with age<70.

12

SPSS 10.0 version

t- chi-square test , p

0.05

결 과

74 74 12 , 62
 64.6±16.2 . 60 가
 56 76% 70
 41 (55%) (Table 1).
 가 , 70
 (Table 2).
 가
 (Table 3). 70
 22 (66.7%)
 70 16 (39.0%) ,
 70 22 (53.7%) 70
 9 (27.2%) (Table 4).
 70
 3 (9.1%) , 70

Table 3. The radiologic finding

Chest X-ray findings	Age<70 (n, %)	Age≥70 (n, %)
Infiltration	22 (66.7)	25 (61.0)
Atelectasis	7 (21.2)	15 (36.6)
Fibrostreaky	4 (12.1)	4 (9.5)
Mass shadow	1 (3)	5 (11.9)
Effusion	1 (3)	1 (2.4)
Cavity	0 (0)	2 (4.8)
Normal	1 (3)	1 (2.4)

Table 4. The classification of endobronchial tuberculosis according to Jung's criteria

EBTB classification	Age<70 (n, %)	Age≥70 (n, %)
Active caseating	22 (66.7)	16 (39.0)*
Edematous hyperemic	9 (27.2)	22 (53.7)*
Tumorous	2 (6.1)	2 (4.9)
Ulcerative	0	1 (2.4)

*p<0.05 compared with age<70.

13. Toichi E, Hanada K, Hosokawa T, Higuchi K, Hosokawa M, Imamura S, et al. Age-related decline in humoral immunity caused by the selective loss of TH cells and decline in cellular immunity caused by the impaired migration of inflammatory cells without a loss of TDTH cells in SAMP1 mice. *Mech Ageing Dev* 1997; 99:199-217.
 14. Kim HO, Ma JE, Lee SJ, Cho YJ, Jeong YY, Jeon KN, et al. Causes of right middle lobe syndrome: recent experience in local tertiary hospital for several years. *Tuberc Respir Dis* 2007;62:192-6.
 15. An JY, Lee JE, Park HW, Lee JH, Yang SA, Jung SS, et al. Clinical and bronchoscopic features in endobronchial tuberculosis. *Tuberc Respir Dis* 2006;60:532-9.
 16. Kim SW, Kim IS, Park DH, No TM, Joeng JK, Jung SW, et al. The clinical significances of bronchial anthracofibrosis in the patients with endobronchial tuberculosis. *Tuberc Respir Dis* 2004;56:495-504.
 17. Chung MP, Lee KS, Han J, Kim H, Rhee CH, Han YC, et al. Bronchial stenosis due to anthracofibrosis. *Chest* 1998;113:344-50.
 18. Lee HS, Maeng JH, Park PG, Jang JG, Park W, Ryu DS, et al. Clinical features of simple bronchial anthracofibrosis which is not associated with tuberculosis. *Tuberc Respir Dis* 2002;53:510-8.
-