

## 폐에 발생한 다형태 암종 환자에서 전이부위 절제술 후 장기 생존

이진구\* · 정경영\* · 박인규\* · 김대준\* · 변천성\* · 조상호\*\*

### Long-term Survival following Surgical Resection for Recurrence of Pulmonary Pleomorphic Carcinoma

Jin Gu Lee, M.D.\*, Kyung Young Chung, M.D.\*, In-Kyu Park, M.D.\*, Dae-Joon Kim, M.D.\*, Cheon-Sung Byeon, M.D.\*, Sang Ho Cho, M.D.\*\*

Pulmonary pleomorphic carcinomas are rare malignant tumors that account for 0.1 to 0.4% of all lung cancers. They are notable for their aggressive clinical behavior and poor prognosis. We report here on a patient who is alive and disease-free 12 years after receiving surgical treatment for the rib recurrence of pulmonary pleomorphic carcinoma.

(Korean J Thorac Cardiovasc Surg 2007;40:587-589)

**Key words:** 1. Lung neoplasms  
2. Survival  
3. Recurrence

### CASE REPORT

A 48-year-old male patient present in 1990 with a 2-month history of cough and blood-tinged sputum. He denied any history of pulmonary disease. Physical examination and routine laboratory studies were unremarkable. A chest X-ray and computed tomogram revealed a 5 cm-sized mass obstructing the left lower lobe bronchus. Needle aspiration biopsy showed that the mass was suspicious of non-small-cell lung cancer or sarcoma. A whole-body bone scan and abdominal ultrasonogram were unremarkable.

Left pneumonectomy was performed through postero-lateral thoracotomy after the 6<sup>th</sup> left rib was resected. The pathological diagnosis indicated pulmonary pleomorphic carcinoma and the resection margin and all lymph nodes were negative

for malignancy; the pathologic stage was Ib. Adjuvant chemotherapy and radiotherapy was planned, but after five weeks of radiation the patient refused further treatment. Four years after the first operation, metastasis of the 7<sup>th</sup> and 8<sup>th</sup> left ribs was found through a chest computed tomogram and whole-body bone scan. Metastasectomy was planned, and an en bloc resection was performed including the 7<sup>th</sup> and 8<sup>th</sup> ribs and a 7×8 cm-sized pleural cavity mass connected to those ribs. Pleomorphic carcinoma cells were detected on the resected ribs and aspirated fluid from the cavity mass (Fig. 1). After 1 cycle of adjuvant chemotherapy, the patient refused further treatment. Twelve years after the last resection, the patient is doing well with a satisfying quality of life and no restriction to his daily activities. A recent chest X-ray showed no sign of recurrence (Fig. 2).

\*연세대학교 의과대학 흉부외과학교실

Department of Thoracic and Cardiovascular Surgery, Yonsei University College of Medicine

\*\*연세대학교 의과대학 병리학교실


Department of Pathology, Yonsei University College of Medicine

논문접수일 : 2007년 3월 6일, 심사통과일 : 2007년 7월 2일

책임저자 : 정경영 (120-752) 서울시 서대문구 신촌동 134번지, 연세대학교 의과대학 흉부외과학교실

(Tel) 02-2228-2140, (Fax) 02-393-6012, E-mail: kychu@yumc.yonsei.ac.kr

본 논문의 저작권 및 전자매체의 지적소유권은 대한흉부외과학회에 있다.



**Fig. 1.** Pleomorphic carcinoma: (A) The spindle cell component shows cytotypic pleomorphism but epithelioid morphology (H&E staining,  $\times 200$ ). (B) The spindle cells stain strongly for epithelial membrane antigen ( $\times 200$ ).



**Fig. 2.** Recent chest X-ray shows no sign of recurrence.

## DISCUSSION

Pleomorphic carcinoma is a rare, epithelial malignant tumor. It has been reported in several human organs, including the lung parenchyma[1]. The 2004 World Health Organization (WHO) classification identified pulmonary pleomorphic carcinoma

as a subgroup of lung cancer among sarcomatoid carcinoma[2], this category of tumors was previously called “carcinoma with pleomorphic, sarcomatoid or sarcomatous elements” in the 1999 WHO classification[3]. Prior to 1999, pleomorphic carcinoma was considered to be a variant of other well-known lung carcinomas because of its biphasic appearance and its frequent association with the other histologic types[4]. Diagnosis of this neoplasm has been problematic and confusing because of the lack of uniform diagnostic criteria. This neoplasm was first diagnosed as carcinosarcoma in 1990, but now, according to the 2004 WHO classification, the diagnosis should be pleomorphic carcinoma.

Patients with these tumors tend to demonstrate a despondent clinical course and their prognosis is generally poor because surgical, irradiation and chemotherapy treatments are usually ineffective. The reported overall survival is 5 to 19 months[1,5,6]. The cause of death is local recurrence and distant metastasis and the prognosis of this tumor seems to depend on the sarcoma component of the tumor. The most frequent metastatic sites are the lymph nodes, but metastases also occur in many other sites, most commonly the kidney, bone, liver, lung, spleen, and gastrointestinal tract[7]. According to the American Society of Clinical Oncology (ASCO)

guidelines for unresectable non-small cell lung cancer[8], resection for distant metastases is recommended only for solitary metastasis of the brain and adrenal gland, while the surgical efficacy of other metastasized sites is doubtful.

To our knowledge, there is no previous report of a long-term survivor after resection of recurrence with pleomorphic carcinoma. In the current case, the patient has been alive and disease-free for 12 years after resection for rib recurrence of pulmonary pleomorphic carcinoma.

In conclusion, the current case stresses that aggressive resection of recurrent pulmonary pleomorphic carcinomas can result in long-term survival with good quality of life in appropriately selected patients.

## REFERENCES

1. Chang YL, Lee YC, Shih JY, Wu CT. *Pulmonary pleomorphic (spindle) cell carcinoma; Peculiar clinicopathologic manifestations different from ordinary non-small cell carcinoma*. Lung Cancer 2001;34:91-7.
2. Beasley MB, Brambilla E, Travis WD. *The 2004 World Health Organization classification of lung tumors*. Semin Roentgenol 2005;40:90-7.
3. Brambilla E, Travis WD, Colby TV, Corrin B, Shimosato Y. *The new World Health Organization classification of lung tumours*. Eur Respir J 2001;18:1059-68.
4. World Health Organization. *Histological typing of lung tumors*. Geneva: WHO. 1981.
5. Raveglia F, Mezzetti M, Panigalli T, et al. *Personal experience in surgical management of pulmonary pleomorphic carcinoma*. Ann Thorac Surg 2004;78:1742-7.
6. Rossi G, Cavazza A, Sturm N, et al. *Pulmonary carcinomas with pleomorphic, sarcomatoid, or sarcomatous elements: a clinicopathologic and immunohistochemical study of 75 cases*. Am J Surg Pathol 2003;27:311-24.
7. Koss MN, Hochholzer L, Frommelt AR. *Carcinosarcomas of the lung: a clinicopathologic study of 66 patients*. Am J Surg Pathol 1999;23:1514-26.
8. Pfister DG, Johnson DH, Azzoli CG, et al. *American society of clinical oncology treatment of unresectable non-small cell lung cancer guideline: update 2003*. J Clin Oncol 2004;22:330-53.

### =국문 초록=

폐에서 다형태 암종은 전체 폐암의 0.1~0.4%를 차지하는 매우 드문 악성 종양으로 나쁜 임상경과와 좋지 않은 예후로 알려져 있다. 저자들은 폐에 발생한 다형태 암종 치료 후 발생한 늑골전이에서 외과적 치료를 통해 12년의 무병 장기 생존을 얻었기에 보고하는 바이다.

중심 단어 : 1. 폐종양  
2. 생존  
3. 재발