

Molecular phylogeny of parasitic Platyhelminthes based on sequences of partial 28S rDNA D1 and mitochondrial cytochrome c oxidase subunit I

Soo-Ung LEE¹⁾, Ha-Chung CHUN²⁾ and Sun HUH^{3)*}

¹⁾*Department of Environmental and Tropical Medicine, School of Medicine, Konkuk University, Chungju, 380-701,*

²⁾*Department of Radiation Oncology, College of Medicine, Hanyang University, Seoul 133-731,*

³⁾*Department of Parasitology, College of Medicine, Hallym University, Chuncheon 200-702, Korea*

Abstract: The phylogenetic relationships existing among 14 parasitic Platyhelminthes in the Republic of Korea were investigated via the use of the partial 28S ribosomal DNA (rDNA) D1 region and the partial mitochondrial cytochrome c oxidase subunit 1 (mCOI) DNA sequences. The nucleotide sequences were analyzed by length, G + C %, nucleotide differences and gaps in order to determine the analyzed phylogenetic relationships. The phylogenetic patterns of the 28S rDNA D1 and mCOI regions were closely related within the same class and order as analyzed by the PAUP 4.0 program, with the exception of a few species. These findings indicate that the 28S rDNA gene sequence is more highly conserved than are the mCOI gene sequences. The 28S rDNA gene may prove useful in studies of the systematics and population genetic structures of parasitic Platyhelminthes.

Key words: Platyhelminthes, molecular phylogeny, 28S ribosomal DNA, mitochondrial cytochrome c oxidase subunit 1

INTRODUCTION

The interrelationships of the Platyhelminthes have attracted attention from both phylogeneticists restricting their attention to morphological characters and those relying solely on molecular data. Recently a number of authors have utilized partial and complete ribosomal DNA and mitochondrial DNA sequences to resolve parasitic helminths phylogenetic relationships

within intraspecies and interspecies (Olson et al., 2001, 2003; Tkach et al., 2001; Vilas et al., 2005).

The first phylogenetic analysis providing explicit character matrices for trematodes was not attempted until Cribb et al. (2001) combined a newly coded morphological matrix with new molecular data from the small subunit ribosomal RNA gene (ssrRNA) from 75 digenean species representing 55 families. Also, recent efforts based primarily on comparative analysis of morphology, and to the lesser extent on molecules, have advanced our understanding of tapeworm systematics and evolution considerably within the phylum Platyhelminthes (Hoberg et al., 1999; Littlewood and Bray, 2001; Olson et al., 2001). However, Platyhelminthes remain still unclear about their morphological features in some species among the experts

• Received 9 March 2007, accepted after revision 10 August 2007.

• This work was supported by the Korea Research Foundation Grant (KRF-2003-015-E00113).

Nucleotide sequences data reported in this paper are available in the GenBank™ database under the accession numbers provided in Tables 2-3.

*Corresponding author (e-mail: shuh@hallym.ac.kr)

(Halanych, 2004).

However, PCR technology and DNA sequencing techniques permit the identification of species, strains, and populations from any stage in their life history to distinguish among morphologically similar parasites (Olson et al., 2001, 2003; Lockyer et al., 2003). So far, many nuclear rDNA and mitochondrial DNA (mtDNA) genes have come into a common use in intraspecies or interspecies molecular systematics on Platyhelminthes (Blaxter et al., 1998; Vilas et al., 2005). The nuclear rDNA gene repeat unit harbors different regions that evolve at varying rates, and thus has also been extensively used in the study of variation and phylogeny at a number of different taxonomic levels. The region incorporated into the first 3 variable domains of 28S rDNA (D1-D3) has been extensively utilized in estimation of the relationships existing within and among the Platyhelminthes (Olson et al., 2003). Mitochondrial DNA has proven to be quite useful for discrimination of closely related organisms, and has been utilized in a number of studies to investigate the phylogenetic relationships existing among eukaryotes at many levels, due to its relatively rapid rate of evolution as compared to the nuclear genome (Lee et al., 2004a, 2004b; Vilas et al., 2005). Mitochondrial

cytochrome c oxidase subunit I (mCOI) is the most highly conserved among 3 genes coding for cytochrome oxidase, and has been employed in phylogenetic studies.

The molecular sequence analysis and phylogenetic relationships among the parasitic helminths remain still unclear in the Republic of Korea. Here we would like to provide additional resolution for the interpretation of helminths evolution. The objective of this paper is to compare the utility of a prospecting DNA marker of ribosomal 28S rDNA D1 region and the mCOI gene sequence for genetic variation and phylogenetic relationships among 14 parasitic Platyhelminthes.

MATERIALS AND METHODS

Parasites

Table 1 shows a taxonomic listing of 14 species (2 classes, 6 orders, 8 families; 9 species of trematodes and 5 species of cestodes) and a recovery source.

Isolation of genomic DNA

Genomic DNA was extracted from the whole adult worms. Each specimen was fixed using 100% ethanol

Table 1. Taxonomic listing of parasites used in this study

Class	Order	Family	Species	Recovery source (strain)	
Trematoda	Opisthorchiida	Opisthorchiidae	<i>Clonorchis sinensis</i>	Rat (S-D) ^{c)}	
		Heterophyidae	<i>Heterophyes nocens</i> ^{a)}	Domestic cat	
			<i>Metagonimus takahashii</i>	Rat (S-D)	
			<i>Metagonimus yokogawai</i>	Rat (S-D)	
			<i>Pygidiopsis summa</i> ^{a)}	Domestic cat	
	Plagiorchiida	Paragonimidae	<i>Paragonimus westermani</i>	Rat (S-D)	
			<i>Plagiorchis muris</i> ^{a)}	Rat (S-D)	
	Echinostomida	Echinostomatidae	<i>Echinostoma hortense</i> ^{b)}	Rat (S-D)	
	Strigeidida	Gymnophallidae	<i>Gymnophalloides seoi</i>	Mouse (BALB/c)	
	Cestoda	Pseudophyllidea	Diphyllobothriidae	<i>Diphyllobothrium latum</i>	Human
<i>Spirometra erinacei</i>				Human	
Cyclophyllidea		Hymenolepididae	<i>Hymenolepis diminuta</i>	Rat (S-D)	
			Taeniidae	<i>Taenia saginata</i>	Human
				<i>Taenia taeniaeformis</i>	Domestic cat

^{a)}Samples were supplied by Dr. Woon-Mok Sohn.

^{b)}Sample was provided by Dr. Jong-Yil Chai.

^{c)}Sprague-Dawley strain.

at room temperature or frozen at -80°C until prepared for DNA extraction. The ethanol in the tissue samples was replaced with 1 M Tris-EDTA buffer (pH 8.0). The frozen worms were lyophilized and lysed with lysis buffer containing 1% SDS, proteinase K (500 $\mu\text{g}/\text{ml}$), and RNase at 37°C for 2-3 hr or overnight. The DNA was extracted in phenol/chloroform, as reported by Sambrook and Russell (2002).

PCR

The PCR of the 28S rDNA D1 region was conducted in a final volume of 50 μl , consisting of extracted DNA as a 50-100 ng template, 200 μM of each dNTP, 10 μmoles of each primer, and 0.5 unit of Ex Taq enzyme (TaKaRa Ex Taq Kit, TAKARA Shuzo Co., Shiga, Japan). The reactions were conducted in a GeneAmp PCR System 4800 thermal cycler (Perkin Elmer, Foster City, California, USA). The PCR reaction cycles consisted of denaturation for 20 sec at 95°C , 30 sec of annealing at 55°C , and 30 sec of extension at 72°C , followed by a final extension of 6 min at the end of 40 cycles. The forward primer (JB10, 5' GATTACCCGCT-GAACTTAAGCATA 3') consisted of the conserved region from the 21st-45th sites of the 28S rDNA sequence of *Schistosoma mansoni*. The reverse primer (JB9, 5' GCTGCATTCACAAACACCCCGACTC 3') was complementary to the 278th-302nd sites of the same gene. The mCOI PCR was conducted using the above method. The forward primer was JB3 (5' TTTTTTGGGCATCCTGAGGTTTAT 3') (2575) and the reverse primer was JB4.5 (5' TAAAGAAAGAA-CATAATGAAAATG 3') (3021). The numbers in brackets refer to the position of the 5' end of the primer from the sequence of *Fasciola hepatica* (Bowles and McManus, 1994; Lee et al., 2004a, 2004b). The PCR conditions were the same as described above, except that annealing was conducted for 30 sec at 48°C .

Cloning of 28S rDNA D1 and mCOI

The PCR products were purified with a Gel Extraction Kit (QIAGEN Co., Valencia, California, USA) and ligated into a T cloning vector (pT7Blue Perfectly Blunt Cloning Kit, Novagen Co., Madison, Wisconsin, USA). The clones were generated via the

transformation of *E. coli* NovaBlue competent cells, which were provided by the supplier. The positive recombinant clones were selected and grown in 2 ml of LB broth, in the presence of 50 $\mu\text{g}/\text{ml}$ ampicillin, overnight at 37°C . The positive plasmids were purified using the QIAprep Spin Plasmid kit (QIAGEN) in accordance with the manufacturer's instructions. The recombinant plasmid was screened using isopropyl- β -thiogalactoside (IPTG) and 5-bromo-4 chloro-3-indolyl- β -D-galactoside (X-gal). Plasmids harboring inserts were digested with the appropriate restriction enzymes for 2 hr at 37°C , and examined with ethidium bromide-stained 1% agarose gels.

Sequencing

DNA sequencing was conducted via the dideoxy chain termination method (Sanger et al., 1977) using the Sequenase kit (ABI Prism Dye Terminator Cycle Sequencing Core Kit, Perkin Elmer) and an automated DNA sequencer (Applied Biosystems model 373A, Perkin Elmer) in accordance with the manufacturer's instructions. The sequencing primers included the T3 and T7 universal primers for both directions. At least 3 clones were sequenced per isolate, with additional clones sequenced as necessary in order to resolve any ambiguous sites.

Sequence analysis and alignment

Sequence analysis was conducted using the nucleotide BLAST program in the NCBI database (National Center for Biotechnology Information, NIH, Bethesda, Maryland, USA) for similarity and nucleotide length (Tatusova and Madden, 1999). We also calculated the fractional GC contents of the nucleic acid sequences, using the EMBOSS GEECEE program provided by Dr. Richard Bruskiwich (Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK) (<http://bioweb.pasteur.fr/seqanal/interfaces/geecce.html>). The nucleotide sequences were compared using the Multalign program (Multiple sequence alignment, European Bioinformatics Institute, <http://www.ebi.ac.uk/clustalw/>) for nucleotide differences and gaps (Corpet, 1988). The resulting aligned output was manually adjusted in

order to improve the homology statements. The alignment gaps were treated as missing data. Sequences corresponding to the PCR amplification primers were excluded prior to multiple sequence alignment and phylogenetic analysis, as primer incorporation during amplification masks potential mismatches (substitutions) that might occur within the priming sites. The confirmed sequences were then deposited in the EMBL/GenBank Data Libraries of the NCBI.

Phylogenetic analysis

All phylogenetic analyses were conducted using the PAUP (Swofford, 2001, PAUP; Phylogenetic Analysis Using Parsimony Version 4.0 d81, Sinauer Associates, Sunderland, Massachusetts, USA). Each molecular data partition was analyzed by both maximum parsimony (MP) and neighbor-joining (NJ) analysis (Saitou and Nei, 1987). The analyses were run using a heuristic search strategy (100 search replicates) with tree bisection-reconnection branch swapping, equally weighted characters and gaps treated as missing data. Bootstrap analysis was performed using a fast heuristic search strategy (1,000 replicates where $\geq 50\%$) for the 28S rDNA D1 and mCOI data. A neighbor-joining analysis was performed using the Kimura 2-parameter model for distance correction; corrected for multiple substitutions (Kimura, 1980).

The 28S rDNA D1 and mCOI sequences used for the analysis and tree construction of *Clonorchis sinensis*, *Plagiorchis muris*, *Metagonimus takahashii*, *Metagonimus yokogawai*, and *Pygidiopsis summa*, as determined for comparative purposes, were the same as those previously reported by Lee and Huh (2004) and Lee et al. (2004a, b). In addition, we utilized the 28S rDNA D1 and mCOI sequences of other species, which have not been previously reported. The sequences from *Gastrocotyle trachuri* (GenBank accession No.: AF131727 for 28S rDNA D1, AY009167 for mCOI) as Monogenea were employed as an outgroup for phylogenetic analysis.

RESULTS

PCR for 28S rDNA D1 and mCOI

We acquired PCR-amplified products from genomic DNA using each of the primer sets. The product length differed between genes or within species. Each of the PCR products was approximately 280-300 base pairs (bp) for 28S rDNA D1 gene and 360-480 bp for mCOI gene in length. No differences in band size between the 28S rDNA D1 and mCOI were observed among the species, with the exception of a few (data not shown).

Sequence analysis

The results of sequence analysis indicate that variation occurs not only in terms of sequence length and G + C percentage, but also in terms of nucleotide differences and gaps (nucleotide insertions, deletions, and substitutions). In the case of 28S rDNA D1 and mCOI, nucleotide differences and gaps were found to be variable within the same class, and several nucleotide insertions, deletions, and substitutions were detected with gaps in the same or different positions (Tables 2-3).

For 28S rDNA D1, the nucleotide sequences ranged from 278 to 313 bp with primer sequences (trematodes; 293-304 bp, cestodes; 278-313 bp) in length among all specimens. The longest nucleotide sequence was 304 bp (*P. muris* in trematodes) and 313 bp (*Taenia saginata* in cestodes) (Tables 2-3). The 28S rDNA D1 nucleotide sequences evidenced a GC content of 50-57% (trematodes; 50-55%, cestodes; 53-57%). The highest G + C percent was 55% (*C. sinensis* in trematodes) and 57% (*T. saginata* in cestodes) (Tables 2-3). The nucleotide differences among species ranged between 0-60 bp in trematodes and 38-155 bp in cestodes when adjusted for gaps. The largest nucleotide difference observed for the 28S rDNA D1 gene was 60 bp between *Gymnophalloides seoi* and *P. muris* in trematodes and 155 bp between *Spirometra erinacei* and *T. saginata* in cestodes (Tables 2-3). The gap differences among species ranged between 0-10 bp in trematodes and 10-29 bp in cestodes when adjusted for gaps. Large gaps of the 28S rDNA D1 gene were 10 bp

Table 2. Comparison of aligned sequence analysis for different trematode species

Gene	Species	Aligned length (bp) ^{a)}	G + C %	GenBank accession No.	Nucleotide differences and gaps ^{b)}								
					1	2	3	4	5	6	7	8	9
28S D1	<i>Clonorchis sinensis</i>	295	55	AF0955330 ^{c)}	0	2	4	1	2	2	1	7	1
	<i>Echinostoma hortense</i>	299	54	AF0955328	40	0	6	2	2	2	1	7	1
	<i>Gymnophalloides seoi</i>	293	50	AF181883	52	49	0	5	6	6	5	10	5
	<i>Heterophyes nocens</i>	298	54	AF181890	37	3	48	0	1	1	0	6	0
	<i>Metagonimus takahashii</i>	299	51	AF095332 ^{c)}	29	45	51	42	0	0	1	5	0
	<i>Metagonimus yokogawai</i>	299	51	AF095331 ^{c)}	29	45	52	42	1	0	1	5	0
	<i>Paragonimus westermani</i>	298	53	AF095329	28	40	49	36	13	13	0	6	0
	<i>Plagiorchis muris</i>	304	52	AF096222 ^{c)}	27	47	60	49	37	37	38	0	6
	<i>Pygidiopsis summa</i>	298	53	AF181885	28	39	49	37	13	13	0	38	0
mCOI	<i>Clonorchis sinensis</i>	444	41	AF181889 ^{c)}	0	19	1	79	4	2	20	50	0
	<i>Echinostoma hortense</i>	463	36	AF096227	114	0	20	102	16	23	37	41	19
	<i>Gymnophalloides seoi</i>	443	35	AF096234	97	105	0	87	5	3	21	65	1
	<i>Heterophyes nocens</i>	366	36	AF188119	242	252	237	0	77	82	100	121	77
	<i>Metagonimus takahashii</i>	448	44	AF096231 ^{c)}	108	131	107	134	0	6	24	52	44
	<i>Metagonimus yokogawai</i>	446	45	AF096230 ^{c)}	103	131	109	235	51	0	18	54	22
	<i>Paragonimus westermani</i>	464	48	AF096228	149	105	134	261	135	127	0	43	20
	<i>Plagiorchis muris</i>	488	48	AF096236 ^{c)}	253	258	257	224	255	255	251	0	48
	<i>Pygidiopsis summa</i>	444	40	AF181884	3	115	97	241	107	110	146	272	0

^{a)}Aligned nucleotide lengths were counted without gaps.

^{b)}Below diagonal is nucleotide differences number adjusted for gaps, above diagonal is nucleotide gaps (insertion, deletion or substitution).

^{c)}This sequences data has been cited from previously reported by Lee and Huh (2004) and Lee et al. (2004a, 2004b).

Table 3. Comparison of aligned sequence analysis for different cestode species

Gene	Species	Aligned length (bp) ^{a)}	G + C %	GenBank accession No.	Nucleotide differences and gaps ^{b)}				
					1	2	3	4	5
28S D1	<i>Diphyllobothrium latum</i>	278	54	AF004719	0	29	20	28	17
	<i>Hymenolepis diminuta</i>	288	53	AF004722	76	0	17	27	21
	<i>Spirometra erinacei</i>	278	54	AF004717	38	64	0	18	10
	<i>Taenia saginata</i>	313	57	AF096224	141	69	155	0	16
	<i>Taenia taeniaeformis</i>	299	54	AF004721	63	39	47	46	0
mCOI	<i>Diphyllobothrium latum</i>	454	32	AF096240	0	1	5	5	28
	<i>Hymenolepis diminuta</i>	444	33	AF096244	86	0	0	4	27
	<i>Spirometra erinacei</i>	453	38	AF096238	101	105	0	4	27
	<i>Taenia saginata</i>	454	34	AF096241	77	99	107	0	18
	<i>Taenia taeniaeformis</i>	473	31	AF096243	58	115	129	131	0

^{a)}Aligned nucleotide lengths were counted without gaps.

^{b)}Below diagonal is nucleotide differences number adjusted for gaps, above diagonal is nucleotide gaps (insertion, deletion or substitution).

between *G. seoi* and *P. muris* in trematodes originally due to nucleotide length, and 29 bp between *Diphyllobothrium latum* and *Hymenolepis diminuta* in

cestodes (Tables 2-3).

For the mCOI gene sequence, the nucleotide sequences ranged from 366 to 488 bp with the primer

Table 4. Character status and tree statistics for phylogenetic analysis using PAUP 4.0

		28S rDNA D1	mCOI
Character status	No. of total characters	367	504
	No. of constants	201	88
	No. of variable characters	66	174
	No. of parsimony-informative characters	100	242
Tree statistics	Tree length	329	1165
	Consistency index (CI)	0.7325	0.6395
	CI excluding uninformative characters	0.6535	0.5459
	Homoplasy index (HI)	0.2675	0.3605
	HI excluding uninformative characters	0.3465	0.4541
	Retention index (RI)	0.7521	0.4897
	Rescaled consistency index (RC)	0.5509	0.3131

sequences (trematodes; 366-488 bp, cestodes; 444-473 bp) in length among all specimens. The longest nucleotide sequence was 488 bp (*P. muris* in trematodes) and 473 bp (*Taenia taeniaeformis* in cestodes) (Tables 2-3). The nucleotide sequences evidenced a GC content of 31-48% (trematodes; 35-48%, cestodes; 31-38%). The highest G + C percentages were 48% (*Paragonimus westermani* and *P. muris* in trematodes), and 38% (*S. erinacei* in cestodes) (Tables 2-3).

For the mCOI sequences, the nucleotide differences among species ranged between 3-272 bp in trematodes and 58-131 bp in cestodes when adjusted for gaps. The largest nucleotide difference was 272 bp between *P. muris* and *P. summa* in trematodes and 131 bp between *T. saginata* and *T. taeniaeformis* in cestodes (Tables 2-3). In addition, gaps among trematode species were more variable than those observed in other cestode species. The gap differences among species ranged from 1-121 bp in trematodes and 0-28 bp in cestodes when adjusted for gaps among classes in proportion to each nucleotide difference. The largest gap was 121 bp between *Heterophyes nocens* and *P. muris* in trematodes, and 28 bp between *D. latum* and *T. taeniaeformis* in cestodes (Tables 2-3). We deposited each sequence into the GenBank database at NCBI under the accession numbers shown in Tables 2-3.

Phylogenetic analysis

The results of the 28S rDNA D1 and mCOI phylo-




Fig. 1. Phylogenetic tree of 28S rDNA D1 gene by maximum parsimony (MP) analysis using the PAUP 4.0 program. Nucleotide sequences were used to construct this tree. Sequences from *Gastrocotyle trachuri* (GenBank accession No.: AF131727) were employed as an outgroup for phylogenetic analysis. Above branch: MP bootstrap values, below branch: NJ bootstrap values, when 1,000 bootstrap replicates where $\geq 50\%$, respectively.

genic trees are provided in Table 4, and Figs. 1-2, respectively. Table 4 shows the results of character status and tree statistics resulting from the individual maximum parsimony analyses. Analyses using the heuristic search algorithm implemented in PAUP 4.0 (Table 4). *G. trachuri* as Monogenea was utilized as an outgroup for both genes in our phylogenetic analysis.

For 28S rDNA D1 gene, in the trematode groups, *C. sinensis* was in the same clade of the phylogenetic tree as *P. muris*. *P. summa* was located in the same clade of the phylogenetic tree as *P. westermani*. However,




Fig. 2. Phylogenetic tree of mCOI gene by maximum parsimony (MP) using the PAUP 4.0 program. Nucleotide sequences were used to construct this tree. Sequences from *Gastrocotyle trachuri* (GenBank accession No.: AFAY009167) were employed as an outgroup for phylogenetic analysis. Above branch: MP bootstrap values, below branch: NJ bootstrap values, when 1,000 bootstrap replicates where $\geq 50\%$, respectively.

Echinostoma hortense and *H. nocens* were genetically close, but different clade with other trematode species. In the cestode groups, *D. latum* was quite closely related to *S. erinacei*. *H. diminuta* and *T. taeniaeformis* were in the same clade of the phylogenetic tree as was *T. saginata* (Fig. 1).

For mCOI gene, in the trematode groups, *C. sinensis* was closely related to *P. summa*, *P. westermani* was in the same clade as the *Metagonimus* species. *E. hortense* was closely related to *G. seoi*, but both of these species were genetically distinct from other trematode groups. *H. nocens* and *P. muris* were generically same clade in our phylogenetic tree. However, these 2 species were located in the different clade from other trematode species. In the cestode groups, *D. latum* was quite closely related to *T. taeniaeformis*. However, *T. saginata*, *H. diminuta*, and *S. erinacei* were genetically distinct from *D. latum* and *T. taeniaeformis* (Fig. 2).

DISCUSSION

So far, only a few is known about the relative divergence rates of nuclear versus mitochondrial DNA in the Platyhelminthes (Morgan and Blair, 1998; Vilas et

al., 2005). Therefore, here we established the utility as markers of interspecific variation and phylogenetic relationship in the Platyhelminthes. In a comparison of genetic variations of the partial 28S rDNA D1 and mCOI gene sequences, our result suggests that the 28S rDNA D1 gene was mostly conserved relative to the mCOI sequences by each sequence analysis. This reason was that mitochondrial DNA may evolve more rapidly than ribosomal DNA, and found that the majority of the nucleotide indels, substitutions occurred at the same or different sites under a variety of biological or environmental conditions. Our data showed consistent patterns with other researcher groups (Vilas et al., 2005). They reported that mitochondrial DNA sequences of the Platyhelminthes accumulate nucleotide substitutions at a much higher rate than internal transcribed spacer (ITS) gene in comparisons of genetic distances.

From our data, we observed genetic variation among species in parasitic Platyhelminthes. The each variation is very variable based on each gene among species. We think that these differences in the nucleotide lengths, GC percentage, nucleotide differences, and number of gaps of each gene can be attributed largely to varying numbers of repeats, copy numbers, deletions, alignment gaps, and base substitutions and additions. Base substitutions and additions are characterized by very high C content which can, at times, represent pure poly-C structures. In addition, this may be a consequence of mutations in the lineage. Also, we may think that other reasons, which mitochondrial DNA may be error by pseudogenes, frame-shift mutations in environmental conditions. Morgan and Blair (1998) reported that the pseudogenes in mitochondrial and ITS genes have been found in parasitic helminths. Because frame-shift mutations or premature stop codons are common in pseudogenes, protein-coding genes of the mtDNA can be error-checked by translating the nucleotide sequences (Benasson et al., 2001).

The phylogenetic patterns of the 28S rDNA D1 gene differed from those of the mCOI. The phylogenetic patterns of the 28S rDNA D1 and mCOI regions were quite closely related among classes, with the exception

of a few species. According to our 28S rDNA D1 data, in the trematode groups, *G. seoi* (Gymnophallidae) was separated from other family species among the class. *M. takahashii* was very closely related to *M. yokogawai* as previously reported by Lee and Huh (2004) and Lee et al. (2004a, 2004b), and other groups evidenced similar results for the ITS1 gene using the PCR-RFLP technique (Yu et al., 1997). In the cestode groups, *H. diminuta* was very closely related to *Taenia* species, due to the smallest nucleotide differences among taxa, despite the disparities in host species and morphology.

The majority of the trematode taxa groups evidenced different patterns for the mCOI gene, with the exception of the *Metagonimus* species. *M. takahashii* was quite close to *M. yokogawai*. This result is consistent with results for the mCOI gene as assessed by PCR-RFLP (Yu et al., 1997). A unique result revealed in the present study is the observation that *H. nocens* and *P. muris* were quite divergent from other trematode groups, possibly as the result of their having a shorter nucleotide sequence than other trematode species.

In the cestodes, the majority of taxa groups for mCOI did not evidence identical 28S rDNA D1 patterns. *D. latum* was in the same clade as the other *Taenia* species group's taxa, but was not in the same clade as *S. erinacei*, even though they are both the same Diphylobothriidae. These observations are inconsistent with the phylogenetic relationship observed between *D. latum* and *S. erinacei* in the 28S rDNA D1 tree. *T. taeniaeformis* was related closely to *T. saginata*. However, Gasser et al. (1999) showed similarity difference or variations between *T. taeniaeformis* and *T. saginata* using the same primers. This mCOI phylogenetic tree did not support the current phylogram on the basis of morphological evidence for within class separation.

We compared both gene sequences by MP and NJ analysis for phylogenetic analysis, and then we acquired the same tree pattern. We also analyzed the 28S rDNA D1 gene sequence by likelihood method (LM) for phylogenetic analysis. This result showed the same tree pattern results, like as MP and NJ methods

(data not shown). However, due to the effects of the large number of taxa and high levels of homoplasy of the mCOI data partition, the LM analysis was stopped prior to completion of the initial heuristic search and a consensus was based on the equally parsimonious trees. Also, we could not do combined analyses between 28S rDNA D1 and mCOI genes, because these genes were different in protein non-coding (28S rDNA) or coding (mCOI).

In conclusion, our data revealed that the 28S rDNA D1 genes are conserved to a higher degree relative to the mCOI sequences. The mCOI resultant phylogenetic tree did not support the current hypotheses on the basis of morphological evidence for the separation of classes, in comparison to the 28S rDNA D1 region. Although more work is needed to determine which factors are involved, we carefully suggest that the 28S rDNA gene may prove useful for studies of systematics and population genetic structures of parasitic Platyhelminthes, although the interrelationships of many families were left unresolved. In addition, further studies will be required for a more adequate elucidation of the phylogeny of Platyhelminthes using other ribosomal DNA and full mitochondrial genome sequences in the future. This data will indirectly aid investigations into the dynamics of morphological and developmental evolution, as well as the biology of parasitism.

ACKNOWLEDGMENTS

The authors wish to thank Dr. Jong-Yil Chai, Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and Dr. Woon-Mok Sohn, Department of Parasitology, Gyeongsang National University College of Medicine, Korea for supplying the specimens.

REFERENCES


- Benasson D, Zhang DX, Hartl DL and Hewitt GM (2001) Mitochondrial pseudogenes: evolution's misplaced witnesses. *Trends Ecol Evol* **16**: 314-321.
- Blaxter ML, De Ley P, Garey JR, Liu LX, Scheldeman P,

- Vierstraete JR, Vanfleteren JR, Mackey LY, Dorris M, Frisse LM, Vida JT, Thomas WK (1998) A molecular evolutionary framework for the phylum Nematoda. *Nature* **392**: 71-75.
- Bowles J, McManus DP (1994) Genetic characterization of the Asian *Taenia*, a newly described taeniid cestode of humans. *Am J Trop Med Hyg* **50**: 33-44.
- Corpet F (1988) Multiple sequence alignment with hierarchical clustering. *Nucl Acids Res* **16**: 10881-10890.
- Cribb TH, Bray RA, Littlewood DTJ, Pichelin SP, Herniou, EA (2001) The Digenea. In *Interrelationships of the Platyhelminthes*, Littlewood DTJ and Bray RA (eds.), Taylor and Francis, London, UK.
- Gasser RB, Zhu X, McManus DP (1999) NADH dehydrogenase subunit 1 and cytochrome c oxidase subunit I sequences compared for members of the genus *Taenia* (Cestoda). *Int J Parasitol* **29**: 1965-1970.
- Halanych KM (2004) The new view of animal phylogeny. *Ann Rev Ecol Evol Syst* **35**: 229-256.
- Hoberg EP, Brooks DR, Mariaux J (1999) Phylogeny of the orders of the eucestoda: morphological and molecular evidence. *Syst Parasitol* **42**: 12-37.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**: 111-120.
- Lee SU, Huh S (2004) Variation of nuclear and mitochondrial DNAs in Korean and Chinese isolates of *Clonorchis sinensis*. *Korean J Parasitol* **42**: 145-148.
- Lee SU, Huh S, Sohn WM (2004a) Molecular phylogenetic location of the *Plagiorchis muris* (Digenea, Plagiorchiidae) based on sequences of partial 28S rDNA and mitochondrial cytochrome c oxidase subunit I. *Korean J Parasitol* **42**: 71-75.
- Lee SU, Huh S, Sohn WM, Chai JY (2004b) Sequence comparisons of 28S ribosomal DNA and mitochondrial cytochrome c oxidase subunit I of *Metagonimus yokogawai*, *M. takahashii* and *M. miyatai*. *Korean J Parasitol* **42**: 129-135.
- Littlewood DTJ and Bray RA (2001) *Interrelationships of the Platyhelminthes*. Taylor and Francis, London, UK.
- Lockyer AE, Olson PD, Littlewood DTJ (2003) Utility of complete large and small subunit rRNA genes in resolving the phylogeny of the Neodermata (Platyhelminthes): implications and a review of the cercomer theory. *Biol J Linn Soc* **78**: 155-171.
- Morgan JAT and Blair D (1998) Relative merits of nuclear ribosomal internal transcribed spacer and mitochondrial COI and ND1 genes for distinguishing among *Echinostoma* species (Trematoda). *Parasitology* **116**: 289-297.
- Olson PD, Cribb TH, Tkach VV, Bray RA and Littlewood DTJ (2003) Phylogeny and classification of the digenea (Platyhelminthes: Trematoda). *Int J Parasitol* **33**: 733-755.
- Olson PD, Littlewood DT, Bay RA, Mariaux J (2001) Interrelationships and evolution of the tapeworms (Platyhelminthes: Cestoda). *Mol Phylogenet Evol* **19**: 443-467.
- Saitou N, Nei M (1987) The neighbor joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**: 406-425.
- Sambrook J, Russell DW (2002) *Molecular cloning: a laboratory manual*, 3rd ed. Cold Spring Harbor Laboratory Press, New York, USA.
- Sanger F, Nicklen S, Coulson AR (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* **74**: 5463-5467.
- Swofford DL (2001) *Phylogenetic analysis using parsimony (*and other methods)*. Version 4. Sunderland, MA: Sinauer.
- Tkach VV, Pawlowski J, Swiderski Z (2001) Molecular phylogeny of the suborder Plagiorchiata and its position in the system of Digenea. In *Interrelationships of Platyhelminthes*, Littlewood DTJ and Bray RA (eds.), Taylor and Francis, London, UK.
- Tatusova TA, Madden TL (1999) BLAST 2 sequences, a new tool for comparing protein and nucleotide sequences. *FEMS Microbiol Lett* **177**: 187-188.
- Vilas R, Criscione CD, Blouin MS (2005) A comparison between mitochondrial DNA and the ribosomal internal transcribed regions in prospecting for cryptic species of platyhelminth parasites. *Parasitology* **131**: 839-846.
- Yu JR, Chung JS, Huh S, Lee SH, Chai JY (1997) PCR-RFLP pattern of three kinds of *Metagonimus* in Korea. *Korean J Parasitol* **35**: 271-276.

IN MEMORIAM

Professor SHIN-YONG KANG

(1946-2007)



Professor Shin-Yong Kang passed away on 7 June 2007, just 1 week before his son's wedding. He was born on 22 September 1946 and grown up in Seoul, Korea. He graduated from College of Veterinary Medicine, Seoul National University in 1970 and obtained Ph.D. from Seoul National University in 1983.

He began to study parasitology as a Teaching Assistant in Seoul National University College of Medicine in 1973. He became an instructor in 1979 and promoted to a Professor in the Department of Parasitology, College of Medicine, Chung-Ang University, and served there during his life time. He was a visiting scientist of Animal Parasitology Institute at Beltsville Agricultural Research Center, Beltsville, Maryland, USA from 1987 to 1988. He studied on pathobiology of metagonimiasis, and also participated in researches on filariasis epidemiology on Jeju Island, serodiagnosis of paragonimiasis and cysticercosis, and molecular biology of parasitism.

Professor Kang was honored by receiving an Academic Award from The Korean Society for Parasitology in 1983 in recognition of his scientific achievements in histopathological studies on metagonimiasis. He served as the Vice President (1994-1997) and President (1998-1999) of The Korean Society for Parasitology. His memory leaves those who knew him with a sense of affection and great admiration. He will not be forgotten for his mentorship and achievements by those, both young and old, who follow him in the challenges of parasitology research.

[Korean Society for Parasitology]