

Antinociceptive and antidiarrhoeal activities of *Sonneratia caseolaris*

F Ahmed, IZ Shahid*, B Baksi and SK Sadhu

Pharmacy Discipline, Khulna University, Khulna-9208, Bangladesh

SUMMARY

The crude ethanol extract of leaves of *Sonneratia caseolaris* Linn. (Sonneratiaceae) was screened for its antinociceptive and antidiarrhoeal activities. The extract produced significant writhing inhibition in acetic acid induced writhing in mice at dose of 250 and 500 mg/kg body weight ($P < 0.01$) comparable to the standard drug diclofenac sodium at the dose of 25 mg/kg of body weight. When tested for its antidiarrhoeal effects on castor oil induced diarrhea in mice, it increased mean latent period and decreased the frequency of defecation significantly at the dose of 500 mg/kg body weight ($P < 0.05$) comparable to the standard drug loperamide at the dose of 50 mg/kg of body weight. The overall results tend to suggest the antinociceptive and antidiarrhoeal activities of the extract.

Key words: Antinociceptive; Antidiarrhoeal; *Sonneratia caseolaris*

INTRODUCTION

Sonneratia caseolaris Linn. (Sonneratiaceae), locally known as ora, choila, etc., is a small tree distributed in the tidal creek and mangrove swamps of Bangladesh, India, Ceylon, Malay, etc. Its fruit is traditionally used in sprains and swellings, hemorrhage, and in the treatment of piles (Kirtikar and Basu, 1987). Based on the traditional usage of this plant, the crude ethanol extract was tested for antioxidant activity using DPPH-radical scavenging effect both quantitatively and qualitatively on TLC in which the extract showed potent antioxidant activity (Ahmed *et al.*, 2006). Although fatty acids, hydrocarbons, steroids, pectin and sugars were previously isolated from this plant (Rollet, 1981; Hogg and Gillan, 1984), by taking DPPH-radical scavenging effect as the isolation guide, Sadhu *et al.* (2006) isolated a flavone, luteolin and its 7-O- β -

glucoside (cynaroside) from the crude extract. However, no other biological activity has yet been reported. The objective of the present study was to investigate the antinociceptive and antidiarrhoeal activity of the crude extracts of leaves of *Sonneratia caseolaris* (*S. caseolaris*).

MATERIALS AND METHODS

Plant material collection and extraction

The plant parts were collected from the Sundarbans' Mangrove Forests, Bangladesh and were taxonomically identified by experts at the Bangladesh National Herbarium (accession no.: 29787). About 400 g of powdered plant material was taken in a clean, flat-bottomed glass container and soaked in 1300 ml of 80% ethanol. The container with its contents was sealed and kept for a period of 7 days accompanying occasional shaking and stirring. The whole mixture then underwent a coarse filtration by a piece of clean, white cotton material followed by a filtration through whatmann filter paper and the filtrate

*Correspondence: IZ Shahid, Pharmacy Discipline, Khulna University, Khulna-9208, Bangladesh. Tel: +880417201713, ext. 252; Fax: +88041731244; E-mail: shahid_israt@yahoo.com

thus obtained was concentrated by using a rotary evaporator (Bibby RE200, Sterilin Ltd., UK) to get the crude extract.

Drugs

Diclofenac Sodium (Oponin Chemical Industries Ltd, Bangladesh), Loperamide (Square Pharmaceuticals Ltd., Bangladesh).

Preliminary phytochemical analysis

The ethanol extract of leaves of *S. caseolaris* was subjected to a preliminary phytochemical screening for major chemical groups. In each test, 10% (w/v) solution of the extract in ethanol was used unless otherwise specified in individual test (Evans, 1989; Ghani, 1998).

Tests for reducing sugar

Benedict's Test: 0.5 ml of the extract was placed in a test tube and then 5 ml Benedict's solution was added to it, boiled for 5 min and allowed to cool spontaneously.

Fehling's Test (Standard Test): 2 ml of the extract was added in 1 ml of a mixture of equal volumes of Fehling's solutions A and B, and was boiled for few min.

Combined Reducing Sugar test: 1 ml of the extract was boiled with 2 ml of diluted hydrochloric acid for 5 min. After cooling the mixture was neutralized with sodium hydroxide solution and then Fehling's test was performed as described above.

Tests for tannins

Ferric Chloride Test: 5 ml of the extract was placed in a test tube and then 1 ml of 5% Ferric chloride solution was added to it.

Potassium dichromate test: 5 ml of the extract was placed in a test tube and then 1 ml of 10% potassium dichromate solution was added.

Test for flavonoids

A few drops of concentrated hydrochloric were

added to 5 ml of the extract.

Test for saponins

1 ml of the extract was placed in a graduated cylinder and was diluted to 20 ml with distilled water and shaken gently for 15 min.

Test for gums

5 ml of the extract was placed in a test tube and then Molish's reagent and sulphuric acid were added to it.

Tests for steroids

Libermann-Burchard test: 1 ml of the extract was placed in a test tube and then 2 ml Libermann-Burchard reagent was added to it.

Sulphuric acid test: 1 ml of the extract was placed in a test tube and 1 ml sulphuric acid was added to it.

Tests for alkaloids

Mayer's test: 2 ml of the extract and 0.2 ml of dilute hydrochloric acid were taken in a test tube and 1 ml of Mayer's reagent was added to it.

Dragendroff's test: 2 ml of the extract and 0.2 ml of dilute hydrochloric acid were placed in a test tube and then 1 ml Dragendroff's reagent was added.

Wagner's test: 2 ml of the extract and 0.2 ml of dilute hydrochloric acid were placed in a test tube. Then 1 ml of iodine solution (Wagner's reagent) was added.

Hager's test: 2 ml solution of the extract and 0.2 ml of dilute hydrochloric acid were placed in a test tube. Then 1 ml of picric acid solution (Hager's reagent) was added.

Animals

Young Swiss-albino mice of either sex, weighing 20 - 25 g, purchased from the Animal Research Branch of the International Centre for Diarrhoeal Disease and Research, Bangladesh (ICDDR, B) were used for the tests. The animals were kept at

animal house (Pharmacy Discipline, Khulna University) for adaptation after their purchase under standard laboratory conditions (relative humidity 55 - 65%, room temperature 25.0 ± 2.0 °C and 12 h light: dark cycle) and fed with standard diets (ICDDR, B formulated) and had free access to tap water.

Pharmacological studies

Antinociceptive activity

Antinociceptive activity of the crude ethanol extract of *S. caseolaris* was tested using the model of acetic acid induced writhing in mice (Whittle, 1964; Ahmed *et al.*, 2004). The experimental animals were randomly divided into four groups, each consisting of five animals. Group I was treated as 'control group' which received 1% (v/v) Tween-80 in water by per oral (p.o.) route at the dose of 10 ml/kg of body weight; group II was treated as 'positive control' and was given the standard drug diclofenac sodium at dose of 25 mg/kg of body weight; group III and group IV were test groups and were treated with the extracts at dose of 250 and 500 mg/kg of body weight respectively. Control vehicle, standard drug and extracts were administered orally, 30 min prior to acetic acid (0.7%) injection. Then after an interval of 15 min, the number of writhes (squirms) was counted for 5 min.

Antidiarrhoeal activity

Antidiarrhoeal activity of the ethanol extract of *S. caseolaris* was tested using the model by castor oil induced diarrhoea in mice (Chatterjee, 1993). The mice were all screened initially by giving 0.5 ml of castor oil and only those showing diarrhoea were selected for the final experiment. The test animals were randomly chosen and divided into three groups having five mice in each. Group-I was kept as control and received 1% Tween-80 at the dose of 10 ml/kg of body weight; group II was treated as 'positive control' and was given the standard drug loperamide at dose of 50 mg/kg of body weight; group III was test group and was treated with the extract at dose of 500 mg/kg of body weight.

Control vehicle, standard drug and the extract were administered orally, 1 h prior to the oral administration of castor oil at a dose of 0.5 ml per mouse. Individual animals of each group were placed in separate cages having adsorbent paper beneath and examined for the presence of diarrhoea every hour in five hours study after the castor oil administration. Number of stools or any fluid material that stained the adsorbent paper were counted at each successive hour during the experiment (5 h). The latent period of each mouse was also counted. At the beginning of each hour new papers were placed for the old ones.

Statistical analysis

Student's *t*-test was used to determine a significant difference between the control group and experimental groups.

RESULTS

Preliminary phytochemical analysis

Results of different chemical tests on the ethanol extract of *S. caseolaris* showed the presence of flavonoids, reducing sugars, gums, saponins and tannins (Table 1).

Antinociceptive activity

Table 2 showed the effect of the ethanol extract of *S. caseolaris* on acetic acid induced writhing in mice. At the dose of 250 and 500 mg/kg of body weight, the extract produced about 21 and 48% writhing inhibition in test animals, respectively. The results were statistically significant ($P < 0.01$) and were comparable to the standard drug diclofenac sodium, which showed 40% writhing inhibition at the dose of 25 mg/kg ($P < 0.01$).

Antidiarrhoeal activity

Antidiarrhoeal activity of the ethanol extract of *S. caseolaris* was tested by castor oil induced diarrhoea in mice. The extract caused an increase in latent period (1.58 h) i.e. delayed the onset of diarrhoeal

Table 1. Chemical groups present in ethanol extract of *S. caseolaris*

| Chemical group | Test solution | Observation | Inference |
|-----------------|---|--|------------------------------|
| Alkaloids | 0.1 ml of Mayer's reagent | Yellowish buff colored precipitate was not obtained | Alkaloids were absent |
| | 0.1 ml of Dragendroff's reagent. | Orange brown precipitate was not observed | |
| | 0.1 ml of iodine solution (Wagner's reagent). | Reddish brown precipitate was not obtained | |
| | 0.1 ml of picric acid solution (Hager's reagent). | Yellowish precipitate was not obtained | |
| Steroids | 2 ml Libermann-Burchard reagent | Reddish purple color was obtained | Steroids were absent |
| | 1 ml sulfuric acid | Chloroform layer had not acquired reddish brown color and acid layer had not showed green fluorescence | |
| Flavonoids | a) 1 ml dilute Ammonia solution | Greenish yellow color was obtained | Flavonoids were present |
| | b) 1 ml dilute sodium carbonate solution | Pale yellow color was obtained | |
| | c) 1 ml dilute sodium hydroxide solution | Yellow color was obtained | |
| Saponins | Shaken in a graduated cylinder for 15 min | One centimeter layer of foam | Saponins were present |
| Reducing sugars | 5 ml Fehling's solution | Brick red colored precipitate | Reducing sugars were present |
| | 5 ml Benedict's reagent | Brick red colored precipitate | |
| | 2 drops of 5% alpha naphthol solution and 1 ml of sulfuric acid | Violet colored ring was formed at the junction of two liquids. | |
| Tannins | 1 ml of 5% Ferric chloride solution | Greenish black precipitate | Tannins were present |
| | 1 ml of 10% Lead acetate solution | Yellow precipitate | |
| | 1 ml of 10% potassium dichromate solution | Yellowish brown precipitate | |
| | 1 ml of aqueous bromine solution | Brown precipitate | |
| Gums | Molish reagent and sulfuric acid | Red -violet ring was produced at the junction of two liquids | Gums were present |

Table 2. Effect of ethanol extract of *S. caseolaris* on acetic acid induced writhing in mice

| Animal Group/Treatment | Number of writhes (% writhing) | Inhibition (%) |
|--|--------------------------------|----------------|
| Group-I (Control) 1% tween-80 solution in water, p.o. | 14.7 ± 0.35 (100) | - |
| Group-II (Positive control) Diclofenac sodium 25 mg/kg, p.o. | 8.8 ± 0.46* (59.86) | 40.14 |
| Group-III Ethanol extract 250mg/kg, p.o. | 11.6 ± 0.37** (78.91) | 21.09 |
| Group-IV Ethanol extract 500 mg/kg, p.o. | 7.6 ± 0.51* (51.70) | 48.30 |

Values are expressed as mean ± S.E.M. (Number of animals, n = 5). * indicates $P < 0.001$, ** indicates $P < 0.01$ vs. control; p.o.: per oral.

episode at the dose of 500 mg/kg body of weight significantly ($P < 0.05$) which was comparable to the standard drug loperamide at the dose of 50 mg/kg body weight in which the value was 1.57 h ($P < 0.05$) (Table 3a). The extract also decreased the frequency of defecation at the same dose where the mean numbers of stool at the 1st, 2nd, 3rd, 4th and 5th

h of study were 0.6, 1.2, 1.4, 0.6 and 0.4 respectively and in standard drug the values were 0.4, 1.6, 2.2, 0.8 and 0.4 respectively (Table 3b).

DISCUSSION

Since *S. caseolaris* belongs to the coastal forests, part

Table 3a. Effect of *S. caseolaris* on castor oil induced diarrhoea in mice (Latent period)

| Animal Group/Treatment | Dose (/kg, p.o.) | Latent period (h) |
|---|------------------|-------------------|
| Group-I (Control) (1% tween-80 solution in water) | 10 ml | 0.766 ± 0.128 |
| Group-II (Positive control) Loperamide. | 50 mg | 1.57 ± 0.191* |
| Group-III Et. Extract | 500 mg | 1.58 ± 0.330* |

Values are expressed as mean ± S. E. M. (n = 5). *indicates $P < 0.05$ vs. control; Et., ethanol; p.o., per oral.

Table 3b. Effect of *S. caseolaris* on castor oil induced diarrhoea in mice (Number of stools)

| Animal Group/Treatment | Dose (/kg, p.o.) | Period of study (h) | Total number of stool |
|--|------------------|---------------------|-----------------------|
| Group-I (Control) (1% tween-80 solution in water) | 10 ml | 1 | 1.8 ± 0.374 |
| | | 2 | 2.8 ± 0.800 |
| | | 3 | 3.0 ± 0.447 |
| | | 4 | 2.6 ± 0.510 |
| | | 5 | 1.8 ± 0.374 |
| Group-II (Positive control) Loperamide | 50 mg | 1 | 0.4 ± 0.400 |
| | | 2 | 1.6 ± 0.678 |
| | | 3 | 2.2 ± .969 |
| | | 4 | 0.8 ± 0.583 |
| | | 5 | 0.4 ± 0.250* |
| Group -III Et. Extract | 500 mg | 1 | 0.6 ± 0.247 |
| | | 2 | 1.2 ± 0.450 |
| | | 3 | 1.4 ± 0.230* |
| | | 4 | 0.6 ± 0.230* |
| | | 5 | 0.4 ± 0.240* |

Values are expressed as mean ± S.E.M. (n = 5). *indicates $P < 0.05$ vs. control; Et., ethanol ; p.o., per oral.

of the plant constituents may be polar in nature. Ethanol was used which has a wide range of solubility in both polar and non-polar region. To avoid any solvent effect on the experimental animals, the solvent was evaporated completely to dryness.

Antinociceptive activity of the ethanol extract of *S. caseolaris* was tested by acetic acid induced writhing model in mice. Acetic acid induced writhing model represents pain sensation by triggering localized inflammatory response. Acetic acid, which is used to induce writhing, causes algia by liberation of endogenous substances, which in turn excite the pain nerve endings (Taesotikul *et al.*, 2003). Increased levels of PGE₂ and PGF_{2α} in the peritoneal fluid have been reported to be responsible for pain sensation caused by intraperitoneal administration of acetic acid (Derardt *et al.*, 1980). The ethanol extract of *S. caseolaris* produced significant writhing inhibition comparable to the

standard drug diclofenac sodium (Table 2). On the basis of this result it can be concluded that the ethanol extract of *S. caseolaris* possesses antinociceptive activity.

Antidiarrhoeal activity of the ethanol extract of *S. caseolaris* was tested by using the model of castor oil induced diarrhoea in mice (Chatterjee, 1993). Castor oil, which is used to induce diarrhoea in mice, mixes with bile and pancreatic enzymes and liberates ricinoleic acid from the triglycerides upon oral administration. Most of the ricinoleic acid remains in the intestine and produces its anti absorptive or secretory effect. The ricinoleic acid thus liberated readily forms ricinoleate salts with sodium and potassium in the lumen of the intestine. The salt formed as such behaves like a soap or surfactant within the gut and at the mucosal surface. Most agreed view is that ricinoleate salts stimulates the intestinal epithelial cell's adenylyl cyclase (Racusen *et al.*, 1979) or release

prostaglandin (Beubler *et al.*, 1979). The extract caused an increase in latent period (1.58 h) i.e. delayed the onset of diarrhoeal episode and decreased the frequency of defecation as well as the number of stool. On the basis of the result of castor oil induced diarrhoea, it can be concluded that the ethanol extract of *S. caseolaris* possesses antidiarrhoeal activity.

In conclusion, it could be suggested that the crude ethanol extract of *S. caseolaris* possesses antinociceptive and antidiarrhoeal. However, further studies are necessary to find out the active principles responsible for these activities.

REFERENCES

- Ahmed F, Selim MST, Das AK, Choudhuri MSK. (2004) Anti-inflammatory and antinociceptive activities of *Lippia nodiflora* Linn. *Pharmazie* **59**, 329-330.
- Ahmed F, Shahid IZ, Razzak MA, Rahman MM, Hoque T, Rahman MT, Sadhu SK. (2006) Free radical scavenging activity of some Mangroves available in Bangladesh. *Orient. Pharm. Exp. Med.* **6**, 58-64.
- Beubler E, Juan H. (1979) Effect of Ricinoleic acid and other Laxatives in Net Water Flux and Prostaglandin E release by the Rat Colon. *J. Pharm. Pharmacol.* **31**, 681-685.
- Chatterjee TK. (1993) *Handbook of laboratory Mice and Rats*. 1st edition, pp. 133-139. Jadavpur University, India.
- Derardt R, Jougney S, Delevalcee F, Falhout M. (1980) Release of prostaglandins E and F in an algogenic reaction and its inhibition. *Eur. J. Pharmacol.* **51**, 17-24.
- Evans WC. (1989) *Trease and Evan's Textbook of Pharmacognosy*. 13 th edition, p. 546. Cambridge University Press, London.
- Ghani A. (1998) *Medicinal Plants of Bangladesh*. The Asiatic Society of Bangladesh, Dhaka, Bangladesh, 1st edition.
- Hogg RW, Gillan FT. (1984) Fatty acids, sterols and hydrocarbons in the leaves from eleven species of Mangrove. *Phytochemistry* **23**, 93-97.
- Kirtikar KR, Basu BD. (1987) *Indian Medicinal Plants*. 2nd edition, pp. 1082-1083. International Book Distributors, India.
- Rollet B. (1981) *Bibliography of Mangrove research. 1600-1975*. UNESCO Paris. p. 479. Pub. Information Retrieval Ltd., London.
- Racusen LC, Binder HJ. (1979) Ricinoleic acid stimulation of active anion secretion in colonic mucosa of the rat. *J. Clin. Invest.* **63**, 743-749.
- Sadhu SK, Ahmed F, Ohtsuki T, Ishibashi. (2006) Flavonoids from *Sonneratia caseolaris*. *J. Nat. Med.* (In press)
- Taesotikul T, Panthong A, Kanjanapothi D, Verpoorte R, Scheffer JJC. (2003) Anti-inflammatory, antipyretic and antinociceptive activities of *Tabernaemontana pandacaqui* Poir. *J. Ethnopharmacol.* **84**, 31-33.
- Whittle BA. (1964) The use of changes in capillary permeability in mice to distinguish between narcotic and non-narcotic analgesics. *Br. J. Pharmacol. Chemother.* **22**, 246-253.