

<Report>

## Biology of *Platydema nigroaeneum* Motschulsky (Coleoptera: Tenebrionidae) from Korea: Life History and Fungal Hosts

Jung, Boo Hee<sup>1\*</sup> and Jin Ill Kim<sup>2</sup>

<sup>1</sup>Laboratory of Behavior & Ecology, Division of Ecoscience, Ewha Womans University, Seoul 120-750, Korea

<sup>2</sup>Department of Biology, Sungshin Women's University, Seoul 136-742, Korea

**ABSTRACT:** We conducted field and laboratory observations of the life histories and fungal hosts of the darkling beetle, *Platydema nigroaeneum* Motschulsky. *P. nigroaeneum*, a fungivorous tenebrionid beetle, was a wide-spread inhabitant of fungi on deciduous trees (*Quercus*, *Salix*, *Alnus* and *Carpinus* etc.) in Korea. Development from egg to adulthood took 4~11 months in nature and about 66 days in the laboratory at 25.5~26.1°C and 63.5~64.5% relative humidity. Both larvae and adults overwintered in their host fungi or beneath the bark of the host tree near the host fungi. Sporophores of *Coriolus* Quélet, *Bjerkandera* Karst., *Daedaleopsis* Schroet. and *Merulius* were common feeding and breeding sites in Korea.

**Key words:** Host fungi, Korea, Life history, *Platydema nigroaeneum*, Tenebrionidae

### INTRODUCTION

Fungi of the order Aphyllophorales, especially the woody and perennial forms, provide microhabitats in which many arthropod species spend their entire lives (Klimaszewski and Peck 1987, Leschen 1990). These fungi provide their inhabitants with food, shelter, and breeding places either directly or indirectly (Graves 1960). The fungivorous tenebrionids, which spend their whole lives in the sporophores associated with dead trees, play an important role in their forest communities (Jung et al. 2007).

It is difficult to study the ecology of fungivorous tenebrionids and other fungivorous beetles because they usually feed on the same sporophores until the fungi are completely destroyed and they also breed in the fruiting bodies of fungi, concealing their life histories. Thus, the biology of mycophagous tenebrionids is poorly understood, although there have been several studies of their life histories and larval development stages (e.g., Lily 1956, Kompantseva 1987a, Leschen 1990, Nadovornaya and Nadvornyy 1991).

*Platydema nigroaeneum* Motschulsky, a fungivorous tenebrionid in the tribe Diaperini (Coleoptera: Tenebrionidae), is common in Korea, and is found in various habitat types from mountain forests to urban gardens. *P. nigroaeneum* is morphologically recognized by its oblong-oval body with a shiny, convex dorsum, and its head with a pair of straight horns in males and a pair of blunt tubercles in females (Kim and Jung 2005). This species is associated with host fungi, generally Aphyllophorales but rarely Agaricales, throughout its life span (Kim and Jung 2005, Jung 2008).

The purpose of this study was to describe the biological characteristics of Korean *P. nigroaeneum*, and to provide comprehensive information on *P. nigroaeneum* life cycles and host-fungi relationships by rearing *P. nigroaeneum* in the laboratory and observing it in the field. This study is the first research on mycophagous tenebrionids associated with fungi in Korea. We provide a description of *P. nigroaeneum*'s life history, fungal hosts, and developmental characteristics.

### MATERIALS AND METHODS

#### Sampling

We collected 212 fruiting bodies of fungi associated with *P. nigroaeneum* from 2005 to 2007. To collect material effectively, we removed the host fungi from the substrate on the forest-floor with cloth or vinyl cloth and quickly transferred them into vinyl bags and sealed the bags. Eggs, larvae, and adults found on the fungi were brought to the laboratory and transferred to plastic containers (30 × 25 × 20 cm) or zippered vinyl bags, which were covered with black cloth to maintain darkness, and were subsequently sorted, identified, counted, and reared.

#### Identification of Fungi

Host fungi collected for this study were identified using several illustrated mushroom books (Lincoff and Nehring 1981, Breitenbach and Kränzlin 1986, Imazeki and Hongo 1987, 1989, Lee 1988, Kim et al. 2004) with the assistance of a mushroom taxonomist, Dr. Seok S. J. (NIASST).

\* Corresponding author; Phone: +82-2-3277-4513, e-mail: starylight12@hanmail.net

## Rearing and Observation

*P. nigroaeneum* was reared in the laboratory in petri dishes (diameter = 9 cm, height = 1.5 cm) at room temperature (25.5~26.1 °C) and at a relative humidity of 63.5~64.5%, and provided with pieces of the host fungus. Behavioral and developmental observations were made in the petri dishes. Head width and body length of specimens were measured using an ocular grid micrometer. Field observations were also carried out once a week from April to August of 2006 in the Kildong Ecological Park, Seoul. We observed the behavior and development of *P. nigroaeneum* in host fungi associated with rotten trees in the forest, and collected specimens and cast exuviae, which were then preserved in 70% ethanol and dry-mounted and deposited in Sungshin Women's University Insect Collection (Seoul, Korea). We used a Stereomicroscope (MZ APO; Leica, Switzerland), a digital camera (Nikon D 200, Japan), and a thermometer (Extech 4465CF) for measurements.

## RESULTS AND CONCLUSION

### The Egg Stage

Freshly laid eggs ( $n = 10$ ) were 0.1~0.2 mm (mean = 0.15 mm) in length and 0.05~0.06 mm (mean = 0.05 mm) in diameter, oval or oblong in shape, opaque white in color, and had a smooth, shiny surface. Egg development under laboratory conditions lasted 3~5 days (Table 1).

### Larvae and Feces

#### The Larval Stage

Most larvae passed through 4 instars, or rarely 3 or 5 instars. The body length of newly hatched instars ( $n = 10$ ) averaged 1.8 mm, and the head capsule averaged 0.2 mm in width. The size of the last instar was an average of 11.25 mm in length and 0.98 mm in

head width (Table 2). The larval stage took about 41 days in the laboratory: the first larval stage took 8~10 days (mean = 9 days), the second larval stage took 9~11 days (mean = 10 days), the third larval stage took 9~13 days (mean = 12 days), and the last larval stage took 11~12 days (mean = 12 days; Table 1). The early first instar does not feed, is not sclerotized in its mouthparts, and the inner part of its body ranges from transparent to visible. The last instar is cylindrical in shape (Fig. 2) and strongly sclerotized on its surface, with a well-developed anal tube in the ninth abdominal tergite. After hatching, *P. nigroaeneum* in the first instar remained under the egg capsules for a while before moving into the fruiting body, after which they moved freely and actively in the fruiting body. While molting, instars of each developmental stage moved rapidly and fed during the trophokinesis period and moved slowly, did not feed, and evacuated food residues during the apolysis period. The post-molting period was characterized by a pale color and less sclerotization (Fig. 2).

#### Feces

Larvae principally lived and fed in the hymenium of the fruiting body. After feeding, larvae produced hair-like filament-feces about 0.2 mm in diameter and over 10 cm in length that formed a 3-dimensional mass. As a result, the host fungi of *P. nigroaeneum* became densely covered with filament-feces, which were produced through all larval stages. We examined feces of *P. nigroaeneum* that had been fed the host fungus *Daedaleopsis tricolor* (Bull.: Fr.) Bond. et Sing. under the optical microscope and did not detect spores and clumps of the hyphal structure in the feces. This suggests that *P. nigroaeneum* larvae may have selected sterile structures rather than fertile hymenium including spores (Fig. 1). Their dung resembled the color of their host fungi: they produced the brown feces after feeding on *D. tricolor* and pink feces after feeding on *Merulius tremellosus* Fr. The filamentous feces might confuse predators and

Table 1. Average developmental period of *P. nigroaeneum* under laboratory conditions from early June to early August, 2006

| Stage           | Egg | Larval stage    |                 |                 |                 | Prepupa | Pupa | Pigmentation | Life cycle (d) |
|-----------------|-----|-----------------|-----------------|-----------------|-----------------|---------|------|--------------|----------------|
|                 |     | 1 <sup>st</sup> | 2 <sup>nd</sup> | 3 <sup>rd</sup> | 4 <sup>th</sup> |         |      |              |                |
| Duration (days) | 4   | 9               | 10              | 11              | 11              | 4       | 9    | 8            | 66             |

Table 2. Size of each larval and pupal stage of *Platydemus nigroaeneum*

( $N = 10$ )

|                  | Larval stage    |                 |                 |                 | Pupae       |
|------------------|-----------------|-----------------|-----------------|-----------------|-------------|
|                  | 1 <sup>st</sup> | 2 <sup>nd</sup> | 3 <sup>rd</sup> | 4 <sup>th</sup> |             |
| Body length (mm) | 1.83 ± 0.06     | 3.47 ± 0.46     | 8.50 ± 1.00     | 11.25 ± 0.87    | 6.20 ± 0.20 |
| Head width (mm)  | 0.20 ± 0.10     | 0.27 ± 0.12     | 0.85 ± 0.10     | 0.98 ± 0.17     | 3.90 ± 0.10 |

at the same time create a 3-dimensional medium into which the agile *P. nigroaeneum* larva can escape, as well as a barrier to entry for other insects, thus restricting competition for food and space in the chamber (Borden et al. 1969).

### The Pupal Stage

#### Prepupae

The C-shaped prepupal body was smaller than the last instar, and developed in the laboratory for 3~5 days (mean = 4 days; Table 1). *P. nigroaeneum* prepupae rarely moved, and did not feed. If touched, they stretched their C-shaped bodies and then twisted from side to side and shook violently.

#### Pupae

The body size of pupae ranged from 6.0~6.5 mm in length (mean = 6.2 mm,  $N=10$ ; Fig. 2), and their development in the laboratory took 8~10 days (mean = 9 days). Pupae did not move or

feed, but if disturbed, they became very active, rapidly rotating clockwise and counter-clockwise and thrashing their abdomens. With the approach of eclosion, their eyes, mouth parts, elytra, tarsal claw and pronotum developed a dark brownish tinge and became visible through the integument.

### The Adult Stage

#### Imagines

Young adults emerging from pupae were pale with a soft integument and gradually became yellowish-brown, reddish-brown and then shiny and black (Fig. 2). Normal sclerotization and color were acquired within 7~9 days (mean = 8 days; Table 1). During this period, they usually remained in the area of overlap of the host fungi and the mass of filament-feces. If disturbed, they rapidly retreated and hid in a dark place.

#### Mating and Oviposition

We were unable to observe mating patterns in detail or to determine the number of eggs deposited during a season in nature because reproductive events were concealed in fungal fruiting bodies. The occurrence of fungi at any time of the year and the concurrent collection of instars of different larval stages suggests that *P. nigroaeneum* mate and deposit eggs repeatedly during the season. Mating was first observed in April~May in nature, while in the laboratory, *P. nigroaeneum* copulated year-round. Most mating occurred at night in this nocturnal species, but mating also occurred in daytime under overlapping fungi, in the bottom of fungi, and in crevices in the tree bark attached to the fungi. In the laboratory, *P. nigroaeneum* copulated for a few minutes in dark areas formed by overlapping fungi. After mating, they deposited their eggs in fungal tissue one at a time. We dissected *P. nigroaeneum* ovaries and found 15~23 eggs.

### Ecological Characteristics

#### Overwinter

*P. nigroaeneum* was observed overwintering in both the larval and adult stages in rotten wood and beneath the bark attaching fungal fruiting bodies and mycelia to trees. Overwintering larvae and adults were collected beginning in late October to November in 2005~2007 in Gyeonggi-do province (Table 3).

#### Generation Time

Development from egg to adulthood in nature took 4~11 months and developmental patterns could be classified into three types. Overwintering adults took about 4 months to complete their development




Fig. 1. Feces of *Platydema nigroaeneum*.

A: After feeding on *Merulius tremellosus*, B: After feeding on *Daedaleopsis tricolor*, C~D: After feeding on *Daedaleopsis tricolor* (20  $\mu\text{m}$ ), E: After feeding on *Bjerkandera adusta* (20  $\mu\text{m}$ ).




Fig. 2. *Platydema nigroaeneum*.

A: After molting, B: Last instar, C: Pupae in the dung, D: Imago, E: Adults.

Table 3. Observation of overwintering *Platydema nigroaeneum*

| Date       | Distribution                          | Host fungi                    | Host tree                 | Life stage   |
|------------|---------------------------------------|-------------------------------|---------------------------|--------------|
| 21-10-2006 | Donggureung, Guri-si, Gyeonggi-do     | <i>Bjerkandera adusta</i>     | <i>Carpinus laxiflora</i> | Adult/Larvae |
| 24-10-2006 | Gwangreung, Namyangju-si, Gyeonggi-do | <i>Bjerkandera adusta</i>     | <i>Carpinus laxiflora</i> | Adult/Larvae |
| 30-10-2006 | Donggureung, Guri-si, Gyeonggi-do     | <i>Bjerkandera adusta</i>     | <i>Carpinus laxiflora</i> | Adult/Larvae |
| 31-10-2006 | Mt. Yongmunasn, Gyeonggi-do           | <i>Bjerkandera adusta</i>     | <i>Alnus japonica</i>     | Adult        |
| 01-11-2006 | Jije-myeon, Gyeonggi-do               | <i>Daedaleopsis conragosa</i> | <i>Alnus japonica</i>     | Adult/Larvae |

Table 4. Larval stage of *Platydema nigroaeneum*

|          | Jan. | Feb. | Mar. | Apr. | May | Jun. | Jul. | Aug. | Sep. | Oct. | Nov. | Dec. |
|----------|------|------|------|------|-----|------|------|------|------|------|------|------|
| I type   |      |      |      |      |     |      |      |      |      |      |      |      |
| II type  |      |      |      |      |     |      |      |      |      |      |      |      |
| III type |      |      |      |      |     |      |      |      |      |      |      |      |

(type I, Table 4), while individuals hatching from the spring eggs of overwintering larvae took 11 months to complete their development (type II, Table 4). Finally, larvae hatching from early summer eggs deposited by overwintering larvae took 10 months to complete their development (type III, Table 4). In the laboratory, development from egg to adulthood took about 66 days.

#### Habits

Instars of all larval stages freely moved around the fruiting bodies of fungi without forming chambers and tunnels. They rapidly moved back and forth in the fruiting body while developing an anal tube on their abdominal tergite. When disturbed, they shook violently and fled to dark places in their host fungi or the fecal mass. Dung of adults is granular, while larval dung forms a filament. Adults and larvae display negative phototropism and have cryptic lifestyles, preventing detailed observation of their lives.

Behavior. Pupae were observed feeding on larvae in laboratory cultures, and multiple adults typically colonized the same host fungus. We recorded an average of 8 individuals per  $2.5 \times 4$  cm section of fungus ( $n = 15$ ) from the bottom of fruiting body. *P. nigroaeneum* was nocturnal, and was most active and frequently observed feeding and mating from 8:00 pm to 4:00 am. During the daytime they mostly hid under their host fungi, in the filament-feces, and in crevices of tree bark attached to their host fungi. They are believed to occasionally move to other nearby locations by walking, and more rarely to fly to more distant places (Heatwole and Heatwole 1967).

Relationships with host fungi. *P. nigroaeneum* was observed feeding and breeding on sporophores of *Daedaleopsis tricolor* (Bull.

Fr.) Bond. et Sing, *Bjerkandera adusta* (Willd.: Fr.) Karst., *Merulius tremellosus* Fr., *Cerrena unicolor* (Fr.) Murr., *Coriolus versicolor* (L.: Fr.) Pilát, *Coriolus brevis* (Berk.) Aoshi., and *Stereum gausapatum* Fr.: Fr. associated with old dead trees. *P. nigroaeneum* was most frequently found on *Daedaleopsis* and *Bjerkandera*, and was observed mainly in old broad-leaf forests dominated by *Quercus*, *Salix*, *Carpinus* and *Alnus* (Krasutskii 2007). Larvae and adults were also observed feeding on *Lampteromyces japonicus* (Kawam.) Sing and *Amillaria* (= *Armillariella*) *mellea* (Vahl: Fr.) P. Kummer of Agaricales on rare occasions but have not been observed breeding in these host species.

#### ACKNOWLEDGMENTS

We would like to express our sincere gratitude to Dr. Hae Chul Park of the National Institute of Agricultural Science and Technology for critical comments on manuscript.

#### LITERATURE CITED

- Borden JH, McClaren M, Horta MA. 1969. Fecal filaments produced by fungus-infesting larvae of *Platydema oregonense*. Ann Entomol Soc Am 62: 444-456.
- Breitenbach J, Kränzlin F. 1986. Fungi of Switzerland, Volume 2 Non gilled fungi (Heterobasidiomycetes, Aphylophorales, Gastromycetes). Verlag Mykologia, Switzerland.
- Graves RC. 1960. Ecological observations on the insects and other inhabitants of woody shelf fungi (Basidiomycetes: Polyporaceae) in the Chicago area. Ann Entomol Soc Am 53: 61-78.
- Heatwole H, Heatwole A. 1968. Movements, host-fungus preferences,

- and longevity of *Bolitotherus cornutus* (Col.: Tene.). *Ann Entomol Soc Am* 61: 18-23.
- Imazeki R, Hongo H. 1987. Colored Illustrations of Mushrooms of Japan Vol. I. Hiokusha Publishing Co., Ltd, Japan.
- Imazeki R, Hongo H. 1989. Colored Illustrations of Mushrooms of Japan Vol. II. Hiokusha Publishing Co., Ltd, Japan.
- Jung BH, Kim SY, Kim JI. 2007. Taxonomic review of the tribe Bolitophagini in Korea (Tenebrionidae: Tenebrioninae). *Entomol Res* 37: 190-196.
- Jung BH. 2008. A taxonomy of Korean Tenebrionida and ecology of fungivorous tenebrionids. Sungshin Women's University, Seoul. (doctoral dissertation).
- Kim JI, Jung BH. 2005. A Taxonomic Review of the Genus *Platydema* Laporte & Brulle in Korea (Coleoptera, Tenebrionidae, Diaperinae). *Entomol Res* 35 (1): 9-15.
- Klimaszewski J, Peck SB. 1987. Succession and phenology of beetles faunas (Col.) in the fungus *Polyporellus squamosus* (Huds.: Fr.) Karst. (Polyporaceae) in Silesia, Poland. *Can J Zool* 65: 542-550.
- Kompantseva TV. 1987a. Ecological peculiarities of xylophilous and mycetophilous darkling beetles (Col. Tene.). In: *Ecology and Morphology of Insects Inhabiting Fungal Substrates* (Pravdin FN, ed). Nauka, Moscow, pp 45-56 (in Russian).
- Krasutskiy BV. 2007. Beetles (Coleoptera) Associated with the Polypore *Daedaleopsis conragosa* (Bolton: Fr.) J. Schrot (Casidiomycetes, Aphyllophorales) in Forests of the Urals and Transurals. *Entomol Rev* 87(5): 512-523.
- Lee JY. 1988. Colored Korean Mushrooms (I). Academy, Seoul.
- Leschen RAB. 1990. Tenebrionoid-Basidiomycete relationships with comments on feeding ecology and the evolution of fungal mycophagy (Col. Hymenomycetes). *Univ Kansas Sci Bull* 54: 165-177.
- Liles M. 1956. A study of the life history of the forked fungus beetle, *Bolitotherus cornutus* (Panzer) (Coleoptera: Tenebrionidae). *Ohio J Sci* 56(6): 329-337.
- Nadovornaya, Nadvornyy. 1991. Biology of the Darkling beetles *Bolitophagus reticulatus* L. and *Uloma culinaria* L. (Coleoptera: Tenebrionidae) in the Forest Steppe Zone of the Ukraine. *Entomologicheskoye Obozneniye* 2: 349-354 (In Russian).
- National Institute of Agricultural Science and Technology Rural Development Administration. 2004. The mushrooms of Korea. Dongbang Media, Seoul.

(Received August 15, 2008; Accepted August 28, 2008)