

## A New Neolignan Glycoside from *Pteris multifida* Poir.

Xu-dong Zheng,<sup>\*</sup> Hao-bin Hu, and Huai-sheng Hu

College of Chemistry and Chemical Engineering, Longdong University, Qingyang 745000, P.R. China

<sup>\*</sup>E-mail: zxd1585@sohu.com

Received November 5, 2007

**Key Words :** *Pteris multifida*, rel-(7*S*,8*S*)- $\Delta^7$ -2,9'-Dihydroxy-5'-methoxy-7,3'-dioxy-8,4'-neolignan-4-*O*- $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside. Multifidoside A

*Pteris multifida* Poir. is widely distributed in the south and southwest district of China (Chinese name "fengweicao"),<sup>1</sup> which has been mainly used as a traditional Chinese folk drug for the treatment of eczema, haematemesis, enteritis, diarrhea, bacillary dysentery cold and are also known to have anticancer and antibacterial effects.<sup>2</sup> However, to the best of our knowledge, very little is known about its chemical constituents except for antimutagenic activity.<sup>3</sup> Our previous paper reported the isolation and characterization of six compounds from EtOAc fraction obtained by partition of the EtOH extract.<sup>4</sup> In a continuation of the phytochemical research on this plant, we now report the isolation and structural elucidation of a new neolignan glycosides, multifidoside A (**1**) from the *n*-BuOH fraction of the EtOH extract, along with four known compounds, scaphopetalone (**2**),<sup>5</sup> (-)-isolariciresinol 3 $\alpha$ -*O*- $\beta$ -apiofuranosyl-(1 $\rightarrow$ 2)-*O*- $\beta$ -glucopyranoside (**3**),<sup>6</sup> 6,7-dihydroxy-3'-methoxy-4',5'-methylenedioxyisoflavone 6-*O*- $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (**4**),<sup>7</sup> polyporusterone I (**5**).<sup>8</sup>

Compound **1**, to which we assigned the name multifidoside A, was obtained as white amorphous powder and has a molecular formula of C<sub>30</sub>H<sub>38</sub>O<sub>15</sub> determined by HRFAB-MS which showed a quasi-molecular formula ion peak at

*m/z*: 639.2289 [M+H]<sup>+</sup> (calcd. for C<sub>30</sub>H<sub>38</sub>O<sub>15</sub>, 638.2211). This formula indicated 12 degrees of unsaturation. The <sup>13</sup>C-NMR and DEPT spectra of **1** clearly displayed 30 carbon signals (2  $\times$  CH<sub>3</sub>, 4  $\times$  CH<sub>2</sub>, 16  $\times$  CH, 8  $\times$  C), of which 11 carbons could be assigned to a glucose unit ( $\delta_C$  104.5, 74.8, 77.5, 71.1, 77.2, 67.8) and an apiose unit ( $\delta_C$  111.1, 77.8, 80.4, 75.0, 65.8), and the remaining 19 carbon signals were assigned to the aglycone. The UV spectrum showed the absorption bands at 208, 266 nm. Its IR spectrum (KBr) showed the absorption bands at 3328 (hydroxyl), 1630 (olefinic C=C), 1601, 1516 (phenyl). The <sup>1</sup>H and <sup>13</sup>C-NMR spectra of **1** showed the presence of two *meta*-coupling aromatic protons signals [ $\delta_H$  6.98 (1H, d, *J* = 1.7 Hz) and 6.83 (1H, d, *J* = 1.7 Hz),  $\delta_C$  110.8 and 116.8], three *asym*-coupling aromatic protons signals [ $\delta_H$  6.42 (1H, d, *J* = 2.4 Hz), 6.44 (1H, dd, *J* = 7.9, 2.4 Hz) and 6.96 (1H, d, *J* = 7.9 Hz),  $\delta_C$  103.9, 108.7 and 116.2], one methoxyl group [ $\delta_H$  3.76 (3H, s),  $\delta_C$  55.5], a (*E*)-coniferyl alcohol signal [ $\delta_H$  4.03 (2H, br d, *J* = 5.7 Hz), 6.39 (1H, d, *J* = 15.3 Hz) and 6.20 (1H, dt, *J* = 15.3, 5.7 Hz),  $\delta_C$  61.5, 128.8 and 126.7],<sup>9</sup> two methine signals [ $\delta_H$  4.79 (1H, d, *J* = 8.0 Hz) and 4.33 (1H, dq, *J* = 8.0, 6.4 Hz),  $\delta_C$  79.5 and 72.9], a methyl signal [ $\delta_H$  1.19 (3H, d, *J* = 6.6 Hz),  $\delta_C$  17.2], one hydroxyl signal [ $\delta_H$




Figure 1. The structure of compounds 1-5.




Figure 2. The key HMBC correlations of compound 1.

9.68 (1H, s, HO-2),  $\delta_C$  154.8 (C-2)], and two anomeric protons of sugars [ $\delta_H$  4.81 (1H, d,  $J = 7.5$  Hz, H-1'') and 5.28 (1H, d,  $J = 2.2$  Hz, H-1'''), the corresponding anomeric carbon signals at  $\delta_C$  104.5 (C-1'') and 111.1 (C-1''')]. Comparison of the  $^1\text{H}$  and  $^{13}\text{C}$ -NMR data of 1 with those of eusiderin E<sup>10</sup> indicated that 1 is a 7,3'-dioxy-8,4'-neolignan glycoside. In HMBC experiment, the correlations of  $\delta_C$  145.8 (C-4) with  $\delta_H$  4.81 (H-1'' of Glc)/6.42 (H-3)/6.44 (H-5)/6.96 (H-6);  $\delta_C$  131.2 (C-1') with  $\delta_H$  6.39 (H-7'')/6.83 (H-6'')/6.98 (H-2'');  $\delta_C$  149.0 (C-5') with  $\delta_H$  3.76 (-OMe)/6.83 (H-6''); and  $\delta_C$  154.8 (C-2) with  $\delta_H$  6.42 (H-3)/6.96 (H-6), suggested that the site of attachment of the disaccharide chain, (*E*)-coniferyl alcohol side-chain, the methoxyl and hydroxyl groups were at C-4, C-1', C-5' and C-2 of the aglycone, respectively.

On acid hydrolysis, compound 1 gave *D*-glucose and *D*-apiose on the basis of *co*-TLC with authentic sample and rotational analysis according to Hudson rules,<sup>11</sup> showing the presence of *D*-glucose and *D*-apiose units. In addition, they were deduced from the FAB-MS spectral observation of *m/z* 507 [ $\text{M}+\text{H}-132$ ]<sup>+</sup> and *m/z* 345 [ $\text{M}+\text{H}-132-162$ ]<sup>-</sup> fragment ions, arising from the elimination of an apiose and a glucose unit, indicating the apiose was terminal sugar and the glucose was attached to the aglycone. Comparison of  $^{13}\text{C}$ -NMR data of the sugar moieties with literature values<sup>12</sup> revealed that the glucose was present in pyranoside form and the apiose was in furanoside form. The HMBC experiment

of 1 showed long-range correlations (Fig. 2) between the H-1''' ( $\delta_H$  5.28) of apiose and the C-6'' ( $\delta_C$  67.8) of glucose as well as between the H-6'' ( $\delta_H$  4.05/3.96) of glucose and the C-1''' ( $\delta_C$  111.1) of apiose, thus suggesting the linkage of apiose-(1 $\rightarrow$ 6)-glucose. The relative stereochemistry of 1 was determined based on the  $^{13}\text{C}$ -NMR spectra data and the  $J$  values measured in the  $^1\text{H}$ -NMR spectrum. The  $\beta$ -configuration on C-1''' anomeric orientation of apiose was confirmed by comparing the  $^{13}\text{C}$ -NMR spectra data of 1 with those of  $\alpha$ -*D*-( $\delta_C$  104.5) and  $\beta$ -*D*-apiofuranosides ( $\delta_C$  111.5), respectively,<sup>13</sup> and the glucose had the  $\beta$ -configuration according to the coupling constant ( $J = 7.5$  Hz) of H-1'' of glucose. The coupling constants observed between H-7' and H-8' ( $J = 15.3$  Hz) suggested that the (*E*)-coniferyl alcohol side-chain had a *trans*-configuration. The signals of H-7 and H-8 in the  $^1\text{H}$ -NMR spectrum appeared at slightly lower fields ( $\delta_H$  4.79 and 4.33, respectively) than verticillatoside A,<sup>14</sup> and with a larger coupling constant ( $J = 8.0$  Hz) indicating a *trans*-orientation (axial-axial) of H-7 and H-8 pair in 1.<sup>15</sup> On these grounds, multifidoside A was elucidated as *rel*-(7*S*,8*S*)- $\Delta^7$ -2,9'-dihydroxy-5'-methoxy-7,3'-dioxy-8,4'-neolignan-4-*O*- $\beta$ -*D*-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -*D*-glucopyranoside.

The known compounds were identified by comparing their spectral data with reported values in the literature or their melting points and  $R_f$  values with authentic samples.

## Experimental

**General Procedures.** Melting points were observed with a Chinese X-4 melting point apparatus (uncorrected). Optical rotations were measured with Perkin-Elmer 241 digital polarimeter. UV and IR (KBr disks) spectra were obtained on Shimadzu UV-300 (double beam) and Alpha-Centari FT-TR spectrophotometer.  $^1\text{H}$  and  $^{13}\text{C}$ -NMR (DEPT) spectra were recorded on Bruker AM-400 NMR spectrometer. Mass

Table 1.  $^1\text{H}$  and  $^{13}\text{C}$ -NMR spectral data of compound 1 (400 and 100 MHz,  $J_{\text{Hz}}$ , DMSO- $d_6$ , TMS )

No.	$\delta_H$	$\delta_C$	DEPT	HMBC (H $\rightarrow$ C)	No.	$\delta_H$	$\delta_C$	DEPT	HMBC (H $\rightarrow$ C)
1		131.2	C	3,6,5,7, HO-2	9'	4.03 (2H, brd, 5.7)	61.5	CH <sub>2</sub>	8', HO-9'
2		154.8	C	3,6,7, HO-2	HO-2	9.68 (1H, s)			
3	6.42 (1H, d, 2.4)	103.9	CH	HO-2,5	HO-9				
4		145.8	C	1'',3,5,6	MeO-5'	3.76 (3H, s)	55.5	CH <sub>3</sub>	
5	6.44 (1H, dd, 7.9, 2.4)	108.7	CH	3,6	Glc-1''	4.81 (1H, d, 7.5)	104.5	CH	
6	6.96 (1H, d, 7.9)	116.2	CH	5,7	2''	3.82 (1H, dd, 9.1, 7.4)	74.8	CH	
7	4.79 (1H, d, 8.0)	79.5	CH	8,6	3''	3.77 (1H, dd, 9.1, 8.5)	77.5	CH	
8	4.33 (1H, dq, 8.0, 6.4)	72.9	CH	7,9	4''	3.94 (1H, dd, 9.9, 8.5)	71.1	CH	
9	1.19 (3H, d, 6.6)	17.2	CH <sub>3</sub>	8	5''	3.82 (1H, ddd, 9.9, 6.0, 1.6)	77.2	CH	
1'		131.2	C	2',6',7'	6''	4.05 (1H, dd, 11.3, 1.6) 3.96 (1H, dd, 11.3, 6.0)	67.8	CH <sub>2</sub>	1'''
2'	6.98 (1H, d, 1.7)	110.8	CH	7',6'	Api-1'''	5.28 (1H, d, 2.2)	111.1	CH	6''
3'		143.6	C	2',7'	2'''	4.29 (1H, d, 2.2)	77.8	CH	
4'		135.5	C	6',8'	3'''		80.4	C	
5'		149.0	C	CH <sub>3</sub> O-,6'	4'''	3.75 (1H, d, 9.4) 3.96 (1H, d, 9.4)	75.0	CH <sub>2</sub>	
6'	6.83 (1H, d, 1.7)	116.8	CH	2',7'	5'''	3.69 (2H, s)	65.8	CH <sub>2</sub>	
7'	6.39 (1H, d, 15.3)	128.8	CH	2',6'					
8'	6.20 (1H, dt, 15.3, 5.7)	126.7	CH	9'					

spectra analyses were carried out with ZAB-HS and MAT-112 mass spectrometer, respectively. Separation and purification were performed by column chromatography on silica gel (100-200, 200-300 mesh). TLC was performed on silica gel GF<sub>254</sub> plates. The spots were visualized by UV (254 nm) and EtOH-H<sub>2</sub>SO<sub>4</sub>.

**Plant Material.** The roots of *P. multifida* Poir. were collected in August 2002, from Pingjiang district of Hunan Province, China. It was identified by Prof. Yun-Shan Lian (Department of Biology, Northwest Normal University). A voucher specimen (No. 107083) of the plant is deposited in the Herbarium of the Botany Department, Northwest Normal University, Lanzhou, 730070, China.

**Extraction and Isolation.** The air-dried and powdered roots of *P. multifida* Poir. (5.0 kg) were soaked in 95% EtOH (15 L, 7 d × 3) at room temperature. After removing the solvent, the extract (250 g) was suspended in warm water and partitioned with petroleum ether (60-90 °C), CHCl<sub>3</sub>, EtOAc, and *n*-BuOH, successively. The *n*-BuOH-soluble fraction was evaporated to give 78.5 g of residues, which was chromatographed on a silica gel column eluting with CHCl<sub>3</sub>-MeOH (8:0→1:5) in increasing polarity and the eluates combined by monitoring with TLC to give three fractions (A, B and C). Fraction A (3.9 g) was further fractionated over silica gel column and eluted with CHCl<sub>3</sub>-MeOH (4:1) to obtain **5** (21 mg). Fraction B (2.6 g) was purified by a silica gel column using CHCl<sub>3</sub>-MeOH (3:1→1:1) as elution gradient to afford **1** (15 mg). Fraction C (3.1 g) was rechromatographed over a silica gel column eluting with EtOAc-MeOH (3:1→2:1) to yield **2** (9 mg) and subfraction. Subfraction was further purified by preparative TLC (silica gel) developed with CHCl<sub>3</sub>-MeOH (1:1) to provide compound **3** (13 mg) and **4** (11 mg).

**Compound 1:** White amorphous powder (MeOH), mp. 216-218 °C; [ $\alpha$ ]<sub>D</sub><sup>20</sup> -11.2° (c = 0.45, MeOH); HRFAB-MS: *m/z* 639.2289 [M+H]<sup>+</sup> (calcd. for C<sub>30</sub>H<sub>38</sub>O<sub>15</sub>, 638.2211); UV

$\lambda_{\max}^{\text{MeOH}}$  (nm): 208, 266; IR  $\nu_{\max}^{\text{KBr}}$  (cm<sup>-1</sup>): 3328 (OH), 1630 (olefinic C=C), 1601, 1516 (phenyl); FAB-MS: *m/z* 639 [M+H], 507 [M+H-132]<sup>+</sup> and 345 [M+H-162-132]<sup>-</sup>; <sup>1</sup>H and <sup>13</sup>C-NMR data see Table 1.

**Acknowledgements.** To Prof. Shang-zhen ZHENG (College of Chemistry and Chemical Engineering, Northwest Normal University) for assistance during my experiment, and to the Educational Foundation of Gansu Province of China (Grant No. 2003039-04) for financial support.

## References

1. Delectis Florae Reipublicae Popularis Sinicae Agendae Academiae Sinicae Edita *Flora Reipublicae Popularis Sinicae*. Science Press: Beijing, 1990; Vol. 3, p 40.
2. Jiangsu New Medical College *Dictionary of Traditional Chinese Drug*. Shanghai Science and Technology Press: Shanghai, 1985; Vol. 1, p 487.
3. Lee, H.; Lin, J. Y. *Mutat. Res/Genet. Toxicol.* **1988**, *204*, 229.
4. Hu, H. B.; Zheng, X. D. *Nat. Prod. Res. Dev.* **2004**, *16*, 379.
5. Vardamides, J. C.; Azebaze, A. G. B.; Nkengfack, A. E.; Van Heerden, F. R.; Fomum, Z. T.; Ngando, T. M.; Conrad, J.; Vogler, B.; Kraus, W. *Phytochemistry* **2003**, *62*, 647.
6. Kanchanapoom, T.; Chumsri, P.; Kasai, R.; Otsuka, H.; Yamasaki, K. *Phytochemistry* **2003**, *63*, 985.
7. Wang, D. Y.; Zheng, Z. Z.; Xu, S. Y.; Zheng, S. Z. *J. Asian Nat. Prod. Res.* **2002**, *4*, 303.
8. Zheng, S. Z.; Yang, H. P.; Ma, X. M.; Shen, X. W. *Nat. Prod. Res.* **2004**, *18*, 403.
9. Hattoti, M. X.; Yang, W.; Shu, Y. Z.; Kakiuchi, N.; Tezuka, Y.; Kikuchi, T.; Namba, T. *Chem. Pharm. Bull.* **1988**, *36*, 648.
10. Cavalcante, S. H.; Yoshida, M.; Gottlieb, O. R. *Phytochemistry* **1985**, *24*, 1051.
11. Capon, B. *Adv. Carbohydr. Chem.* **1960**, *15*, 11.
12. Agrawal, P. K. *Phytochemistry* **1992**, *31*, 3307.
13. Kitagawa, I.; Hori, K.; Sakagami, M.; Hashiuchi, F.; Yoshikawa, M.; Ren, J. *Chem. Pharm. Bull.* **1993**, *41*, 1350.
14. Su, B. N.; Yang, L.; Jia, Z. J. *Phytochemistry* **1997**, *45*, 1271.
15. Fang, J. M.; Lee, C. K.; Cheng, Y. S. *Phytochemistry* **1992**, *31*, 3659.