

Structure and Functional Effect of Tryptophan Mutants of *Nicotiana tabacum* Acetohydroxyacid Synthase

Im-Joung La, Masud Karim, and Moon-Young Yoon*

Department of Chemistry, Hanyang University, Seoul 133-791, Korea. *E-mail: myyoon@hanyang.ac.kr

Received June 23, 2008

Key Words : Acetohydroxyacid synthase. Circular dichroism. Tryptophan quenching. Herbicide inhibition

The branched chain amino acids (BCAAs) are synthesized by plants, algae, fungi, bacteria and archaea, but not by animals. Therefore, the enzymes of the BCAA biosynthetic pathway are potential targets for the development of herbicides, fungicides and antimicrobial compounds. Most interest has focused on the first enzyme in this pathway, acetohydroxyacid synthase (AHAS, EC 2.2.16) because it is the target of several herbicides including the sulfonylureas (SUs),¹ imidazolinones (IMs),² triazolopyrimidines³ and pyri-midinyl oxybenzoates.^{4,5} AHAS catalyzes the condensation of two molecules of pyruvate to give rise to 2-acetolactate in the first step of valine and leucine biosynthesis, and in parallel, it also catalyzes the condensation of pyruvate and 2-ketobutyrate to yield 2-aceto-2-hydroxybutyrate in the second step of isoleucine biosynthesis.⁶ AHAS requires thiamine diphosphate (ThDP), flavin adenine dinucleotide (FAD), and a divalent metal ion, Mg²⁺ or Mn²⁺, as cofactors for its catalytic function.

In the past, research on plant AHAS has been hindered due to its instability and low abundance in plants. The AHAS genes from *Arabidopsis thaliana*⁷ and tobacco⁸ have been functionally expressed in *E. coli* and purified as glutathione S-transferase fusion proteins. Recently, we have reported several herbicide-resistant mutants of tobacco AHAS⁹ and also identified the role of lysine,¹⁰ arginine,¹¹ histidine,¹² methionine,¹³ aspartic acid¹⁴ and tryptophan residues.¹⁵ The inhibition effect of herbicides was also characterized for several AHAS sources.¹⁵⁻¹⁷ In the tobacco AHAS two tryptophan residues (W573 and W490) were reported as critical for enzyme activity.¹⁸ The yeast crystal model suggests that the pyrimidine ring of ThDP would stack on the indole ring of W586 (W573 in tobacco).¹⁹ Within the γ -domain of AHAS three tryptophan residues were identified (W490, W503, and W573) which could potentially participate in hydrophobic stacking interactions.

To determine the catalytic role of tryptophan residues (W266, W439, W490, W503 and W573) in tobacco AHAS, we carried out site-directed mutagenesis in this study. The effect of the mutations on the kinetic parameters, cofactor binding, structure of the enzyme and inhibition by herbicides were analyzed.

All mutants were expressed and successfully purified to homogeneity. SDS-PAGE analysis showed that all mutants had the same molecular mass as that of wild type enzyme (data not shown). The ability of wild type and tryptophan

mutant proteins to produce acetolactate in the absence of 2-ketobutyrate was measured and the results are shown in Table 1. The acetolactate formation also serves as an indirect measure of the affinity of the wild type and various mutant proteins for the substrate. The kinetic parameters were obtained by analysis of Michaelis-Menten plots of initial rates for AHAS under standard cofactor conditions. Among the five tryptophan mutants, the W573F showed a 30-fold reduction in enzyme activity whereas W490F had completely abolished the enzyme activity. The K_m value for W573F was approximately 25-fold higher than that of wild type, indicating the perturbation of the substrate binding site. The kinetic parameters of W503F, W266F and W439F were not significantly altered relative to those of wild type. These data indicate that W490 and W573 are the most important catalytic tryptophan residues involved in the catalysis. The results were identical with previous reports by Chong *et al.*¹⁸

The cofactors-dependent quenching of intrinsic protein fluorescence leads to a reduction in equilibrium fluorescence and is a measure of the cofactors binding affinity to AHAS. Molecular modeling studies of the γ -domain intermediate complex revealed that W573 is largely buried within the structure in a nonpolar pocket and W490 is also in a largely nonpolar region sandwiched between helices within the β -domain. To investigate the function of these two tryptophan residues (W490 and W573), fluorescence quenching studies were carried out with varying ThDP concentrations. The quenching of tryptophan fluorescence provides a direct tool for the determination of the dissociation constant of a protein-ligand complex. However, the sensitivity of the method depends on the number and location of the fluorophores.

Table 1. Kinetic parameters and IC₅₀ values of wild type AHAS and tryptophan mutants

Enzyme	K_m for pyruvate (mM)	V_{max} (U/mg)	IC ₅₀ value for sulfonylureas		
			SMM (nM)	CE (nM)	PSM (nM)
Wild type	14.7	6.47	2.31 ± 0.42	6.12 ± 0.73	5.46 ± 0.21
W266F	45.6	8.11	3.45 ± 0.51	4.23 ± 0.22	7.34 ± 0.71
W439F	25.17	5.66	4.35 ± 0.21	8.16 ± 0.26	4.68 ± 0.64
W490F	NA	NA	NA	NA	NA
W503F	53.22	4.65	3.56 ± 0.61	27.05 ± 0.28	9.45 ± 0.13
W573F	424.27	1.31	87.1 ± 0.23	73.32 ± 0.45	68.17 ± 0.82

NA: No enzymatic activity

Tobacco AHAS exhibited intrinsic fluorescence characteristics due to the presence of tryptophan residues, very likely buried in the protein, as revealed by a low fluorescence emission maximum (340-345 nm) upon excitation at 300 nm.²⁰ Replacement of each of the three tryptophan residues within the catalytic domain followed by titration with ThDP revealed that substitution of both W490 and W573 residues alleviated the substrate-induced fluorescence quenching (data not shown). In order to further investigate this quenching mechanism, titration of a number of tryptophan replacement mutant proteins with ThDP was conducted. When W573 was replaced with phenylalanine in the AHAS protein's active site, the alleviation of the observed quenching upon ThDP binding was the largest among the mutant proteins tested. The largest effect was observed upon substitution of W573 (a change from 42% to 25% quenching).²¹ The reduction in the fluorescence quenching was smaller but significant for the W490F mutant (42% to 36% quenching; Figure 1), whereas W266F, W439F and W503F had quench-




Figure 1. The effect of ThDP on the fluorescence emission spectrum of wild type and mutant AHAS (W490F). Each 0.1 mg/mL of wild type AHAS (A) and the same concentration of W490F (B) enzyme was pre-incubated with a buffer containing 100 mM potassium phosphate buffer pH 7.4, 20 μ M FAD, 2 mM MgCl₂, and the following indicated concentrations of ThDP were added. 0 (1), 0.06 (2), 0.1 (3), 0.12 (4), 0.2 (5), 0.25 (6), 0.4 (7), 0.5 (8), 0.8 (9) and 1.0 (10) mM (the curve numbers were indicated through an arrow from 01 to 10). The wavelength of excitation was 300 nm and the emission spectra were recorded in the range 310-450 nm.

ing similar to the wild type protein (data not shown). The dissociation constants were determined by fitting the data to Equation (1) where F_0 is the intrinsic fluorescence of enzyme in the absence of quencher, F_1 is the observed fluorescence at a given $[Q]$ (concentration of quencher), f_a is the fractional degree of fluorescence and K_d is the dissociation constant.

$$\left(1 - \frac{F_1}{F_0}\right) = \frac{f_a \cdot [Q]}{K_d + [Q]} \quad (1)$$

Fluorescence quenching of W490F and W573F²¹ by ThDP gives dissociation constants (K_d) of 0.54 ± 0.01 mM and 0.40 ± 0.01 mM respectively, values that are virtually identical with wild type protein $K_d = 0.34$ mM. These results imply that the ThDP binding itself into a deep channel of the active site was not affected by the single mutations (W573F or W490F). The titration and intrinsic fluorescence quenching of W573F by pyruvate imply that pyruvate binding in the active site was not affected by the substitution at position 573.²¹ The absorption and fluorescence emission peaks of W490F corresponding to the cofactor FAD indicated that the W490F mutation prevents this cofactor binding to AHAS with a consequent loss of enzymatic activity.¹⁸ Thus, W490 likely plays an indirect role in the catalytic mechanism of the enzyme, possibly in maintaining the active site integrity. However, the K_m value of the W573F mutant was 30-fold higher than that of wild type, suggesting that W573 may play a major role in the formation of enzyme-substrate or enzyme bound intermediate complex. These results imply that W573 plays only an indirect role in cofactor binding, likely acting as an anchor that stabilizes the loop containing M569 which packs against the pyridine ring of ThDP.

In order to determine whether the mutations induced any secondary structural alterations in the enzyme, the CD spectra of the wild type and mutants were recorded in the far-UV regions. The W573F showed slight differences in the far-UV region, whereas other mutants were similar to that of




Figure 2. Far-UV CD spectra of the wild type (solid line) and W573F (dotted line) mutant enzymes. Each protein was present at a concentration of 0.20 mg/mL in 10 mM potassium phosphate buffer (pH 7.5). The secondary structural components were determined by fitting the data to an algorithm, the convex constraint analysis (CCA).²⁴

Table 2. Secondary structure of wild type and W573F AHAS calculated from CD analysis

Protein	α -Helix (%)	β -Sheet (%)	Random (%)	Turn (%)	RMS
Wild type	27.3	41.8	18.3	12.6	9.645
W573F	38.1	0.0	32.3	29.6	13.53

wild type enzyme (Figure 2 and data not shown). The secondary structural components were determined for both wild type and W573F mutant (Table 2) by fitting the data to an algorithm, the convex constraint analysis (CCA).²² However, no significant secondary structural changes were detected in the W573F except for increases in α -helix and random conformations with the disappearance of β -sheets (Table 2).

Earlier, Chong *et al.* reported that the W573F mutant of tobacco AHAS was reported as a sulfonylurea and triazolopyrimidine resistant form.¹⁸ This tryptophan residue was also responsible for herbicide resistance in yeast AHAS,²³ *Arabidopsis thaliana* AHAS²⁴ and *E. coli* AHAS II,²⁵ with a greatly decreased preference for 2-ketobutyrate as the second substrate. To date this, both wild type and mutant proteins were tested for resistance against three potent sulfonylurea herbicides (SMM, CE and PSM). The 50% inhibition concentrations (IC₅₀) were analyzed by fitting the data to Equation (2).

$$v = \frac{(V_0 - V_f) \cdot IC_{50}}{IC_{50} + [I]} + V_f \quad (2)$$

In this equation V_0 is the reaction rate without inhibitor, V_f is the rate at maximal inhibition and $[I]$ is an inhibitor concentration. These results indicate that mutants W266F, W439F and W503F showed similar inhibition patterns to wild type and the inhibition effect was not determined for W490F (Table 1). However, the W573F showed approximately 20- to 30-fold increase in IC₅₀ values with all sulfonylureas tested. The significant increase in IC₅₀ values for three potent sulfonylurea herbicides (SMM, CE and PSM) suggests that W573F has a major role in sulfonylurea herbicides binding site and in substrate recognition.

In conclusion, the present results suggest that W490 residue is likely located at or near the cofactor FAD binding site and may affect the binding affinity of FAD,¹⁸ while the W573 residue plays only an indirect role in cofactor binding but plays a role in substrate recognition and process a common binding site for the sulfonylurea classes of herbicides. The W573F mutant appears to be a good candidate for the development of herbicide resistant transgenic plants.

Experimental Section

Site-directed mutagenesis of tobacco AHAS was performed directly on the plasmid derived from pGEX-2T containing tobacco AHAS cDNA, using the PCR mega primer method. All manipulations of primers and DNA were carried

out using the techniques reported previously.¹⁸

Bacterial strains of *E. coli* BL21-DE3 cells, containing the expression vector pGEX-AHAS, were grown at 37 °C in Luria-Bertani (LB) broth medium containing 50 μ g/mL ampicillin to an OD₆₀₀ of 0.7-0.8. Expression of the pGEX-AHAS gene was induced by the addition of 1mM isopropyl-d-thiogalactoside (IPTG). Cells were grown for an additional 16 h at 18 °C and harvested by centrifugation at 5000 g for 30 min. Purification of wild type and mutant AHAS were carried out as described previously by Chang *et al.*²⁶

Enzyme activities of the wild type and mutant AHAS were measured according to the method of Westerfeld with a modification as reported previously.²⁷ The reactions were carried out in standard reaction mixture containing 100 mM potassium phosphate buffer pH 7.5, 1 mM ThDP, 10 mM MgCl₂, 20 μ M FAD, and 75 mM pyruvate. Different concentrations of herbicides were added to determine the inhibition effect. Assays were initiated by the addition of AHAS at 37 °C for 30 min and terminated by the addition of 6 N H₂SO₄. The reaction product acetohydroxyacid was allowed to decarboxylate at 60 °C for 15 min. The acetoin formed by acidification was incubated and colorized with 0.5% creatine and 5% α -naphthol at 60 °C for 15 min. The absorbance of the reaction mixture was monitored at 525 nm. One unit (U) of activity was defined as the amount of enzyme required for the production of 1 μ M of acetohydroxyacid per minute under the standard assay conditions described above.

The extent of ThDP binding was measured by monitoring the quenching of intrinsic fluorescence of the enzyme after addition of varying concentrations of ThDP. The purified enzymes (0.1 mg/mL) were pre-incubated in buffer containing 100 mM phosphate buffer pH 7.4, 20 μ M FAD, 2 mM MgCl₂ and the indicated concentrations of ThDP at 37 °C for 15 min. The wavelength of excitation was 300 nm and the emission was recorded in the range of 310-450 nm (bandwidth of 2 nm) in a computer-controlled PTI Alphascan-2 spectrofluorometer (Photon Technology Inc., South Brunswick, NJ). Each emission value was the mean of three independent experiments. The far-UV CD spectrum was obtained on a Jasco J-715 spectropolarimeter using 0.1-cm pathlength cuvettes at 25 °C. The samples were prepared in 10 mM sodium phosphate pH 7.4 and the protein concentration was 0.1-0.2 mg/mL. A minimum of four spectra were accumulated for each sample and the contribution of the buffer was always subtracted. The secondary structural components from the CD spectra were analyzed by the new algorithm, the convex constraint analysis (CCA).²²

Acknowledgments. This work was supported by the research fund of Hanyang University (HY-2007-I).

References

- LaRossa, R. A.; Schloss, J. V. *J. Biol. Chem.* 1984, 259, 8753.
- Shaner, D. L.; Anderson, P. C.; Stidham, M. A. *Plant Physiol.* 1984, 76, 545.
- Gerwick, B. C.; Subermanian, V. I.; Loney-Gallant, V. I.; Chander,

- D. P. *Pestic Sci.* **1990**, *29*, 357.
4. Shimizu, T.; Nakayama, I.; Nakao, T.; Nezu, Y.; Abe, H. *J. Pestic Sci.* **1994a**, *19*, 59.
 5. Shimizu, T.; Nakayama, I.; Wada, N.; Nakao, T.; Abe, H. *J. Pestic Sci.* **1994b**, *19*, 257.
 6. Duggleby, R. G.; Pang, S. S. *J. Biochem. Mol. Biol.* **2000**, *33*, 1.
 7. Singh, B.; Schmitt, G.; Lillis, M.; Hand, J. M.; Misra, R. *Plant Physiol.* **1991**, *97*, 657.
 8. Chang, S. I.; Kang, M. K.; Choi, J. D.; Namgoong, S. K. *Biochem. Biophys. Res. Commun.* **1997**, *234*, 549.
 9. Jung, S. M.; Le, D. T.; Yoon, S. S.; Yoon, M. Y.; Kim, Y. T.; Choi, J. D. *Biochem. J.* **2004**, *383*(1), 53.
 10. Yoon, T. Y.; Chung, S. M.; Chang, S. I.; Yoon, M. Y.; Hahn, T. R.; Choi, J. D. *Biochem. Biophys. Res. Commun.* **2002**, *293*, 433.
 11. Le, D. T.; Yoon, M. Y.; Kim, Y. T.; Choi, J. D. *J. Biochem.* **2005**, *138*, 35.
 12. Oh, K. J.; Park, E. J.; Yoon, M. Y.; Han, T. R.; Choi, J. D. *Biochem. Biophys. Res. Commun.* **2001**, *282*, 1237.
 13. Le, D. T.; Yoon, M. Y.; Kim, Y. T.; Choi, J. D. *Biochem. Biophys. Res. Commun.* **2003**, *306*, 1075.
 14. Le, D. T.; Yoon, M. Y.; Kim, Y. T.; Choi, J. D. *Biochimica et Biophysica Acta* **2005**, *1749*, 103.
 15. Choi, K. J.; Pham, C. N.; Jung, H.; Han, S. H.; Choi, J. D.; Kim, J.; Yoon, M. Y. *Bull. Korean Chem. Soc.* **2007**, *28*, 1109.
 16. Le, D. T.; Lee, H. S.; Chung, Y. J.; Yoon, M. Y.; Choi, J. D. *Bull. Korean Chem. Soc.* **2007**, *28*, 947.
 17. Choi, K. J.; Noh, K. M.; Choi, J. D.; Park, J. S.; Won, H. S.; Kim, J. R.; Kim, J. S.; Yoon, M. Y. *Bull. Korean Chem. Soc.* **2006**, *27*, 1697.
 18. Chong, C. K.; Shin, H. J.; Chang, S. I.; Choi, J. D. *Biochem. Biophys. Res. Commun.* **1999**, *259*, 136.
 19. Pang, S. S.; Duggleby, R. G.; Guddat, L. W. *J. Mol. Biol.* **2002**, *317*, 249.
 20. Kim, J.; Beak, D. G.; Kim, Y. T.; Choi, J. D.; Yoon, M. Y. *Biochem. J.* **2004**, *384*, 59.
 21. Karim, M.; Shim, M. Y.; Kim, J.; Choi, K. J.; Kim, J. R.; Choi, J. D.; Yoon, M. Y. *Bull. Korean Chem. Soc.* **2006**, *27*, 549.
 22. Perczel, A.; Hollosi, M.; Tusnady, G.; Fasman, G. D. *Protein Eng.* **1991b**, *4*, 669.
 23. Duggleby, R. G.; Pang, S. S.; Yu, H.; Guddat, L. W. *Eur. J. Biochem.* **2003**, *270*, 2895.
 24. Chang, A. K.; Duggleby, R. G. *Biochem. J.* **1998**, *333*, 765.
 25. Engel, S.; Vyamensky, M.; Vinogradov, M.; Berkovich, D.; Barilan, A.; Qimron, U.; Rosiansky, Y.; Barak, Z.; Chipman, D. J. *Biol. Chem.* **2004**, *279*, 24803.
 26. Le, D. T.; Choi, J. D. *Bull. Korean Chem. Soc.* **2005**, *26*, 916.
 27. Kim, J.; Choi, J. D.; Kim, B. H.; Yoon, M. Y. *Bull. Korean Chem. Soc.* **2005**, *26*, 260.
-