

## STRONG CONVERGENCE THEOREM FOR UNIFORMLY $L$ -LIPSCHITZIAN MAPPINGS IN BANACH SPACES

XIAOLONG QIN\*, SHIN MIN KANG\*\*, AND MEIJUAN SHANG\*\*\*

ABSTRACT. In this paper, we prove strong convergence theorems for a finite family of uniformly  $L$ -Lipschitzian mappings by a cyclic iterative algorithm in the framework of Banach spaces. Our results improve and extend the recent ones announced by many others.

### 1. Introduction and preliminaries

Throughout this paper, we assume that  $E$  is a real Banach space,  $E^*$  is the dual space of  $E$ ,  $K$  is a nonempty closed convex subset of  $E$  and  $J : E \rightarrow 2^{E^*}$  is the normalized duality mapping defined by

$$J(x) = \{f \in E^* : \langle x, f \rangle = \|x\|^2 = \|f\|^2, \|f\| = \|x\|\}, \quad \forall x \in E,$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $E$  and  $E^*$ . The single-valued normalized duality mapping is denoted by  $j$ .

DEFINITION 1.1. Let  $T : K \rightarrow K$  be a mapping.

(1)  $T$  is said to be *uniformly  $L$ -Lipschitzian* if there exists  $L > 0$  such that, for any  $x, y \in K$ ,

$$\|T^n x - T^n y\| \leq L\|x - y\|, \quad \forall n \geq 1;$$

(2)  $T$  is said to be *nonexpansive* if the following inequality holds for all  $x, y \in K$

$$\|Tx - Ty\| \leq \|x - y\|;$$

(3)  $T$  is said to be *asymptotically nonexpansive* if there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that for any given

---

Received November 18, 2007; Accepted May 20, 2008.

2000 Mathematics Subject Classification: Primary 47H09; Secondary 47H10.

Key words and phrases: asymptotically pseudo-contractive mapping, normalized duality mapping, uniformly  $L$ -Lipschitzian mapping, Banach space.

$x, y \in K$ ,

$$\|T^n x - T^n y\| \leq k_n \|x - y\|, \quad \forall n \geq 1;$$

(4)  $T$  is said to be *asymptotically pseudo-contractive* if there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that, for any  $x, y \in K$ , there exists  $j(x - y) \in J(x - y)$

$$\langle T^n x - T^n y, j(x - y) \rangle \leq k_n \|x - y\|^2, \quad \forall n \geq 1.$$

REMARK 1.1. (1) It is easy to see that if  $T$  is an asymptotically nonexpansive mapping, then  $T$  is a uniformly  $L$ -Lipschitzian mapping, where  $L = \sup_{n \geq 1} \{k_n\}$ . And every asymptotically nonexpansive mapping is asymptotically pseudo-contractive, but the inverse is not true, in general.

(2) The concept of asymptotically nonexpansive mappings was introduced by Goebel and Kirk [4]. Asymptotically nonexpansive mappings include nonexpansive mappings as a special case when  $\{k_n\} = 1$ .

Recently, Schu [7] proved the following strong convergence theorem for asymptotically pseudo-contractive mappings

THEOREM 1.1. *Let  $H$  be a Hilbert space,  $\emptyset \neq K \subset H$  closed bounded convex;  $L > 0$ ;  $T : K \rightarrow K$  completely continuous, uniformly  $L$ -Lipschitzian and asymptotically pseudo-contractive with sequence  $\{k_n\} \subset [1, \infty)$ ;  $q_n = 2k_n - 1$  for all  $n \geq 1$ ;  $\sum_{n=1}^{\infty} (q_n^2 - 1) < \infty$ ;  $\{\alpha_n\}, \{\beta_n\} \subset [0, 1]$ ;  $\epsilon \leq \alpha_n \leq \beta_n \leq b$  for all  $n \geq 1$  some  $\epsilon > 0$  and some  $b \in (0, L^{-2}[\sqrt{1 + L^2} - 1])$ ;  $x_1 \in K$ ; for all  $n \geq 1$ , define*

$$\begin{cases} y_n = (1 - \beta_n)x_n + \beta_n T^n x_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n y_n. \end{cases}$$

Then  $\{x_n\}$  converges strongly to some fixed point of  $T$ .

In [1], Chang extended Theorem 1.3 to a real uniformly smooth Banach space. To be more precise, he proved the following theorem:

THEOREM 1.2. *Let  $E$  be a uniformly smooth Banach space,  $D$  be a nonempty bounded closed convex subset of  $E$ ,  $T : D \rightarrow D$  be an asymptotically pseudo-contractive mapping with a sequence  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  and  $F(T) \neq \emptyset$ , where  $F(T)$  is the set of fixed points of  $T$  in  $D$ . Let  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$  and  $\{\delta_n\}$  be for sequence in  $[0, 1]$  satisfying the following conditions:*

- (a)  $\alpha_n + \gamma_n \leq 1, \beta_n + \delta_n \leq 1$ ;
- (b)  $\alpha_n \rightarrow 0, \beta_n \rightarrow 0$  ( $n \rightarrow \infty$ );

(c)  $\sum_{n=0}^{\infty} \alpha_n = \infty, \sum_{n=0}^{\infty} \gamma_n = \infty.$

Let  $x_0 \in D$  be any given point and let  $\{x_n\}, \{y_n\}$  be the modified Ishikawa iterative sequence errors defined by

$$\begin{cases} x_{n+1} = (1 - \alpha_n - \gamma_n)x_n + \alpha_n T^n y_n + \gamma_n u_n, \\ y_n = (1 - \beta_n - \delta_n)x_n + \beta_n T^n x_n + \delta_n v_n, \quad n \geq 0. \end{cases}$$

If there exists a strict increasing function  $\phi : [0, \infty) \rightarrow [0, \infty)$  with  $\phi(0) = 0$  such that

$$\langle T^n x_n - x^*, j(x_n - x^*) \rangle \leq k_n \|x_n - x^*\|^2 - \phi(\|x_n - x^*\|), \quad \forall n \geq 0,$$

where  $x^* \in F(T)$  is some fixed point of  $T$  in  $D$ , then  $x_n \rightarrow x^*$  as  $n \rightarrow \infty$ .

The purpose of this paper is, by using a simple and quite different method, to prove some strong convergence theorems for a finite family of  $L$ -Lipschitzian mappings in stead of the assumption that  $T$  is a uniformly  $L$ -Lipschitzian and asymptotically pseudo-contractive mapping in a Banach space. Our results extend and improve some recent results announced by Chang [1], Cho et al. [3], Ofoedu [6], Schu [7] and many others

In order to prove our main results, we also need followings lemmas:

LEMMA 1.3 ([2]). Let  $E$  be a real Banach space and  $J : E \rightarrow 2^{E^*}$  be the normalized duality mapping. Then for any  $x, y \in E$ ,

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \quad \forall j(x + y) \in J(x + y).$$

LEMMA 1.4 ([5]). Let  $\{\theta_n\}$  be a sequence of nonnegative real numbers and  $\{\lambda_n\}$  be a real sequence satisfying the following conditions:

$$0 \leq \lambda_n \leq 1, \quad \sum_{n=0}^{\infty} \lambda_n = \infty.$$

If there exists a strictly increasing function  $\phi : [0, \infty) \rightarrow [0, \infty)$  such that

$$\theta_{n+1}^2 \leq \theta_n^2 - \lambda_n \phi(\theta_{n+1}) + \sigma_n, \quad \forall n \geq n_0,$$

where  $n_0$  is some nonnegative integer and  $\{\sigma_n\}$  is a sequence of nonnegative number such that  $\sigma_n = o(\lambda_n)$ , then  $\theta_n \rightarrow 0$  as  $n \rightarrow \infty$ .

LEMMA 1.5. Let  $\{a_n\}$  and  $\{b_n\}$  be two nonnegative real sequences satisfying the following condition:

$$a_{n+1} \leq (1 + \lambda_n)a_n + b_n, \quad \forall n \geq n_0,$$

where  $\{\lambda_n\}$  is a sequence in  $(0, 1)$  with  $\sum_{n=0}^{\infty} \lambda_n < \infty$ . If  $\sum_{n=0}^{\infty} b_n < \infty$ , then  $\lim_{n \rightarrow \infty} a_n$  exists.

## 2. Main results

**THEOREM 2.1.** *Let  $E$  be a real Banach space,  $K$  a nonempty closed convex subset of  $E$  and  $\{T_i\}_{i=1}^N : K \rightarrow K$  a finite family of uniformly  $L$ -Lipschitzian mappings such that  $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$ , where  $F(T_i)$  denotes the set of fixed points of  $T_i$ . Let  $\{k_n\} \subset [1, \infty)$  be a sequence with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$ . Let  $\{\alpha_n\} \subset (0, 1)$  be a sequence satisfying the following conditions:*

- (a)  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (b)  $\sum_{n=0}^{\infty} \alpha_n^2 < \infty$ ;
- (c)  $\sum_{n=0}^{\infty} \alpha_n(k_n - 1) < \infty$ .

For any  $x_0 \in K$ , let  $\{x_n\}$  be the iterative sequence defined by

$$(2.1) \quad x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T_{s_n}^n x_n, \quad n \geq 0,$$

where  $s_n = n \bmod N$ , with the mod function taking values in the set  $\{1, 2, \dots, N\}$ . If there exists a strict increasing function  $\phi : [0, \infty) \rightarrow [0, \infty)$  with  $\phi(0) = 0$  such that

$$\langle T_i^n x - x^*, j(x - x^*) \rangle \leq k_n \|x - x^*\|^2 - \phi(\|x - x^*\|)$$

for all  $j(x - x^*) \in J(x - x^*)$  and  $x \in K$ ,  $i = 1, 2, \dots, N$ , then  $\{x_n\}$  converges strongly to a point  $x^* \in F$ .

*Proof.* First, we prove that the sequence  $\{x_n\}$  defined by (2.1) is bounded. Actually, it follows from (2.1) and Lemma 1.3 that

$$\begin{aligned} & \|x_{n+1} - x^*\|^2 \\ &= \|(1 - \alpha_n)(x_n - x^*) + \alpha_n(T_{s_n}^n x_n - x^*)\|^2 \\ (2.2) \quad &\leq \|(1 - \alpha_n)(x_n - x^*)\|^2 + 2\alpha_n \langle T_{s_n}^n x_n - x^*, j(x_{n+1} - x^*) \rangle \\ &\leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 + 2\alpha_n L \|x_n - x_{n+1}\| \|x_{n+1} - x^*\| \\ &\quad + 2\alpha_n \{k_n \|x_{n+1} - x^*\|^2 - \phi(\|x_{n+1} - x^*\|)\}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & \|x_{n+1} - x_n\| = \|(1 - \alpha_n)x_n + \alpha_n T_{s_n}^n x_n - x_n\| \\ (2.3) \quad &= \alpha_n \|x_n - T_{s_n}^n x_n\| \\ &\leq \alpha_n \|x_n - x^*\| + \alpha_n \|T_{s_n}^n x_n - x^*\| \\ &\leq \alpha_n (1 + L) \|x_n - x^*\|. \end{aligned}$$

Substituting (2.3) into (2.2), we have

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 \\
 &\quad + 2\alpha_n \{k_n \|x_{n+1} - x^*\|^2 - \phi(\|x_{n+1} - x^*\|)\} \\
 &\quad + 2L\alpha_n^2(1 + L) \|x_n - x^*\| \|x_{n+1} - x^*\| \\
 (2.4) \qquad &\leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 + 2\alpha_n \{k_n \|x_{n+1} - x^*\|^2 \\
 &\quad - \phi(\|x_{n+1} - x^*\|)\} + L\alpha_n^2(1 + L) (\|x_n - x^*\|^2 \\
 &\quad + \|x_{n+1} - x^*\|^2).
 \end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \frac{(1 - \alpha_n)^2 + L\alpha_n^2(1 + L)}{1 - 2\alpha_n k_n - \alpha_n^2 L(1 + L)} \|x_n - x^*\|^2 \\
 &\quad - \frac{2\alpha_n \phi(\|x_{n+1} - x^*\|)}{1 - 2\alpha_n k_n - \alpha_n^2 L(1 + L)} \\
 (2.5) \qquad &= \left[1 + \frac{\alpha_n^2 + 2\alpha_n(k_n - 1) + 2\alpha_n^2 L(1 + L)}{1 - 2\alpha_n k_n - \alpha_n^2 L(1 + L)}\right] \|x_n - x^*\|^2 \\
 &\quad - \frac{2\alpha_n \phi(\|x_{n+1} - x^*\|)}{1 - 2\alpha_n k_n - \alpha_n^2 L(1 + L)}.
 \end{aligned}$$

Since  $\alpha_n \rightarrow 0$  as  $n \rightarrow \infty$ , there exists a positive integer  $n_0$  such that

$$\frac{1}{2} < 1 - 2\alpha_n k_n - \alpha_n^2 L(1 + L) \leq 1, \quad \forall n \geq n_0.$$

Therefore, it follows from (2.5) that

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \left[1 + 2(\alpha_n^2 + 2\alpha_n(k_n - 1) + 2\alpha_n^2 L(1 + L))\right] \|x_n - x^*\|^2 \\
 (2.6) \qquad &\quad - 2\alpha_n \phi(\|x_{n+1} - x^*\|) \\
 &\leq \left[1 + 2(\alpha_n^2 + 2\alpha_n(k_n - 1) + 2\alpha_n^2 L(1 + L))\right] \|x_n - x^*\|^2.
 \end{aligned}$$

By the conditions (b) and (c), we have

$$2 \sum_{n=0}^{\infty} [\alpha_n^2 + 2\alpha_n(k_n - 1) + 2\alpha_n^2 L(1 + L)] < \infty.$$

It follows from Lemma 1.5 that the limit  $\lim_{n \rightarrow \infty} \|x_n - x^*\|$  exists. Therefore, the sequence  $\{\|x_n - x^*\|\}$  is bounded. Without loss of generality, we can assume that  $\|x_n - x^*\|^2 \leq M_1$ , where  $M_1$  is an appropriate positive constant. Take  $\theta_n = \|x_n - x^*\|$ ,  $\lambda_n = 2\alpha_n$  and

$$\sigma_n = 2[(\alpha_n^2 + 2\alpha_n(k_n - 1) + 2\alpha_n^2 L(1 + L))]M_1.$$

(2.6) can be written as

$$\theta_{n+1}^2 \leq \theta_n^2 - \lambda_n \phi(\theta_{n+1}) + \sigma_n, \quad \forall n \geq n_0.$$

By the conditions (a)-(c), we know that all the conditions in Lemma 1.4 are satisfied. Therefore, it follows that

$$\lim_{n \rightarrow \infty} \|x_n - x^*\| = 0,$$

that is,  $x_n \rightarrow x^*$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$

REMARK 2.1. Theorem 3.1 mainly improves the corresponding ones announced by Chang [1], Cho et al. [3], Ofoedu [6] and Schu [7]. The method given in the proof of Theorem 2.1 is also different from the method given by Ofoedu [6].

REMARK 2.2. Under suitable conditions, the sequence  $\{x_n\}$  defined by (2.1) in Theorem 2.1 can be generalized to the iterative sequences with errors. Since the proof is trivial, we omit it here.

As a corollary of Theorem 2.1, we have the following result immediately.

COROLLARY 2.2. *Let  $E$  be a real Banach space,  $K$  a nonempty closed convex subset of  $E$  and  $T : K \rightarrow K$  a uniformly  $L$ -Lipschitzian mapping with  $F(T) \neq \emptyset$ , where  $F(T)$  denotes the set of fixed points of  $T$ . Let  $\{k_n\} \subset [1, \infty)$  be a sequence with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$ . Let  $\{\alpha_n\} \subset [0, 1]$  be a sequence satisfying the following conditions:*

- (a)  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (b)  $\sum_{n=0}^{\infty} \alpha_n^2 < \infty$ ;
- (c)  $\sum_{n=0}^{\infty} \alpha_n(k_n - 1) < \infty$ .

For any  $x_0 \in K$ , let  $\{x_n\}$  be the iterative sequence defined by

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, \quad n \geq 0.$$

If there exists a strict increasing function  $\phi : [0, \infty) \rightarrow [0, \infty)$  with  $\phi(0) = 0$  such that

$$\langle T^n x - x^*, j(x - x^*) \rangle \leq k_n \|x - x^*\|^2 - \phi(\|x - x^*\|)$$

for all  $j(x - x^*) \in J(x - x^*)$  and  $x \in K$ , then  $\{x_n\}$  converges strongly to  $x^*$ .

### References

- [1] S. S. Chang, *Some results for asymptotically pseudo-contractive mappings and asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **129** (2001), 845–853.
- [2] S. S. Chang, *On Chidume's open questions and approximation solutions of multi-valued strongly accretive mapping equation in Banach spaces*, J. Math. Anal. Appl. **216** (1997), 94–111.
- [3] Y. J. Cho, J. I. Kang, and H. Y. Zhou, *Approximating common fixed points of asymptotically nonexpansive mappings*, Bull. Korean Math. Soc. **42** (2005), 661–670.
- [4] K. Gobel and W. A. Kirk, *A fixed points theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35** (1972), 171–174.
- [5] C. Moore and B.V. Nnoli, *Iterative solution of nonlinear equations involving set-valued uniformly accretive operators*, Comput. Math. Appl. **42** (2001), 131–140.
- [6] E. U. Ofoedu, *Strong convergence theorem for uniformly  $L$ -Lipschitzian asymptotically pseudocontractive mapping in a real Banach space*, J. Math. Anal. Appl. **321** (2006), 722–728.
- [7] J. Schu, *Iterative construction of fixed point of asymptotically nonexpansive mappings*, J. Math. Anal. Appl. **158** (1991), 407–413.

\*

Department of Mathematics  
Gyeongsang National University  
Chinju 660-701, Republic of Korea  
*E-mail*: qxlxajh@163.com

\*\*

Department of Mathematics and the RINS  
Gyeongsang National University  
Chinju 660-701, Republic of Korea  
*E-mail*: smkang@nongae.gsnu.ac.kr

\*\*\*

Department of Mathematics  
Shijiazhuang University  
Shijiazhuang 050035, P.R. China  
*E-mail*: meijuanshang@yahoo.com.cn