

## ALGEBRAIC STRUCTURES IN A PRINCIPAL FIBRE BUNDLE

JOON-SIK PARK\*

ABSTRACT. Let  $P(M, G, \pi) =: P$  be a principal fibre bundle with structure Lie group  $G$  over a base manifold  $M$ . In this paper we get the following facts:

1. The tangent bundle  $TG$  of the structure Lie group  $G$  in  $P(M, G, \pi) =: P$  is a Lie group.
2. The Lie algebra  $\mathfrak{g} = T_e G$  is a normal subgroup of the Lie group  $TG$ .
3.  $TP(TM, TG, \pi_*) =: TP$  is a principal fibre bundle with structure Lie group  $TG$  and projection  $\pi_*$  over base manifold  $TM$ , where  $\pi_*$  is the differential map of the projection  $\pi$  of  $P$  onto  $M$ .
4. for a Lie group  $H$ ,  $TH = H \circ T_e H = T_e H \circ H = TH$  and  $H \cap T_e H = \{e\}$ , but  $H$  is not a normal subgroup of the group  $TH$  in general.

### 1. Introduction

In this note, a general survey on the principal fibre bundle  $TP(TM, TG, \pi_*) =: TP$  induced from a principal fibre bundle  $P(M, G, \pi) =: P$  which is appeared in [7, p. 55] is explained in details. As by-products, we obtain the following facts:

- the Lie algebra  $\mathfrak{g} = T_e G$  is a normal subgroup of the Lie group  $TG$ .
- for a Lie group  $H$ ,  $H \circ T_e H = T_e H \circ H = TH$  and  $H \cap T_e H = \{e\}$ , but the subgroup  $H$  of the group  $TH$  is not a normal subgroup of  $TH$  in general, and so the group  $TH$  and the product group  $H \times T_e H (= T_e H \times H)$  of  $H$  and  $T_e H$  are not group-isomorphic in general.

---

Received June 08, 2008; Accepted August 14, 2008.

2000 Mathematics Subject Classification: Primary 53C07; Secondary 53A15.

Key words and phrases: principal fibre bundles, free actions, splitting exact sequences.

\*This research was supported by Pusan University of Foreign Studies Research Grant in 2006.

## 2. The proof of main results

Let  $M$  be a  $C^\infty$ -manifold and  $G$  a Lie group. A principal fibre bundle over  $M$  with group  $G$  consists of a manifold  $P$  and an action of  $G$  on  $P$  satisfying the following conditions:

(1)  $G$  acts freely on  $P$  on the right:

$$P \times G \ni (u, a) \mapsto ua = R_a(u) \in P;$$

(2)  $M$  is the quotient space of  $P$  by the equivalence relation by  $G$ ,  $M = P/G$ , and the canonical projection  $\pi : P \rightarrow M$  is differentiable;

(3)  $P$  is locally trivial, that is, every point  $x$  of  $M$  has a neighbourhood  $U$  such that  $\pi^{-1}(U)$  is isomorphic with  $U \times G$  in the sense that there is a diffeomorphism  $\Psi : \pi^{-1}(U) \rightarrow U \times G$  such that  $\Psi(u) = (\pi(u), \psi(u))$  where  $\psi$  is a mapping of  $\pi^{-1}(U)$  into  $G$  satisfying  $\psi(ua) = \psi(u) \cdot a$  for all  $u \in \pi^{-1}(U)$  and  $a \in G$ .

A principal fibre bundle will be denoted by  $P(M, G, \pi)$ ,  $P(M, G)$  or simply  $P$ . In general, we call  $P$  the total space or the bundle space,  $M$  the base space,  $G$  the structure group and  $\pi$  the projection.

Given a mapping  $f$  of a manifold  $M$  into another manifold  $M'$ , the differential at a point  $p \in M$  of  $f$  is the linear mapping  $f_*$  of  $T_p M$  into  $T_{f(p)} M'$  which is defined as follows. For each  $X \in T_p M$ , choose an integral curve  $x(t)$  of the vector  $X$  in  $M$  such that  $X$  is the vector tangent to  $x(t)$  at  $p = x(t_0)$ . Then  $f_*(X)$  is the vector tangent to the curve  $f(x(t))$  at  $f(p) = f(x(t_0))$ . It follows immediately that if  $g$  is a function differentiable in a neighbourhood of  $f(p)$ , then  $(f_*(X))(g) = X(g \circ f)$ . We may also consider  $f_*$  as the map of  $TM := \bigcup_{p \in M} T_p M$  into

$$TM' = \bigcup_{q \in M'} T_q M'.$$

A binary operation  $\circ$  on the tangent bundle  $TG$  of a Lie group  $G$  can be defined as follows:

For  $X \in T_g G$  and  $Y \in T_{g'} G$  ( $g, g' \in G$ ), choose curves  $x(t)$  and  $y(t)$  in  $G$  such that  $X$  and  $Y$  are the vectors tangent to  $x(t)$  and  $y(t)$  at  $g = x(t_0)$  and  $g' = y(t_0)$ , respectively. Then  $X \circ Y := Xg' + gY := dR_{g'}(X) + dL_g(Y) \in T_{gg'} G \subset TG$  is the vector tangent to the curve  $x(t) \cdot y(t)$  at  $g \cdot g' = x(t_0) \cdot y(t_0) \in G$ .

Then  $TG$  is a group with respect to the operation  $\circ$  on  $TG$  which is just defined. In fact, the zero vector  $O_e$  belonging to  $T_e G$  is the identity element of  $TG$  with respect to the operation  $\circ$ , where  $e$  is the identity element of the Lie group  $G$ . For  $X \in T_g G, Y \in T_h G$  and  $Z \in T_k G$

$(g, h, k \in G), (X \circ Y) \circ Z = X \circ (Y \circ Z)$ . Moreover, with respect to the operation  $\circ$  on  $TG$ , the inverse element of  $X(\in T_gG)$  which is tangent to a curve  $x(t)$  in  $G$  at  $g = x(t_0)$  is the vector tangent to the curve  $x(t)^{-1}$  at  $g^{-1} = x(t_0)^{-1} \in G$ .

We may regard the Lie group  $G$  and its Lie algebra  $\mathfrak{g}$  also as subgroups  $\{O_g|O_g$  is the zero vector in  $T_gG, g \in G\}$  and  $T_eG$  of the group  $TG$  with the operation  $\circ$ . Moreover, for an arbitrary given  $g \in G$  and an arbitrary given  $X$  belonging to the space  $T_gG, X \circ T_eG = T_eG \circ X$ . So,  $T_eG = \mathfrak{g}$  is a normal subgroup of the Lie group  $TG$ .

For  $X \in T_gG, X \circ G = \{X \circ k = Xk|k \in G\}$  and  $G \circ X = \{h \circ X = hX|h \in G\}$ . So, in general  $X \circ G \neq G \circ X$ . Hence,  $G$  is not a normal subgroup of the group  $TG$  in general. Evidently,  $G \cap T_eG = \{e\}$ . And, by the definition of the operation  $\circ$  on  $TG, G \circ T_eG = T_eG \circ G = TG$ .

Thus we have

**THEOREM 2.1.** *Let  $G$  be a Lie group. Then,*

- (1) *the differential of the group operation of  $G \times G$  into  $G$  is a group operation on  $TG$ .*
- (2) *the Lie group  $G$  is a subgroup of the Lie group  $TG$ .*
- (3) *its Lie algebra  $\mathfrak{g} = T_eG$  are a normal subgroup of the group  $TG$ .*
- (4)  *$G \circ T_eG = T_eG \circ G = TG$  and  $G \cap T_eG = \{e\}$ , but in general  $G$  is not a normal subgroup of  $TG$ .*

Let  $H$  and  $K$  be two normal subgroups of a group  $X$ . Then,  $X$  is said to be the *direct product* of the normal subgroups  $H$  and  $K$  iff  $HK(= KH) = X$  and  $H \cap K = \{e\}$  ([5, p. 62]).

By virtue of the fact (4) of Theorem 2.1, we obtain

**COROLLARY 2.2.** *For a Lie group  $G, TG = G \circ T_eG = T_eG \circ G = TG$  and  $G \cap T_eG = \{e\}$ , but the group  $TG$  and the product group  $G \times T_eG(= T_eG \times G)$  of  $G$  and  $T_eG$  are not group-isomorphic in general.*

A sequence of group homomorphisms

$$G_1 \xrightarrow{f_1} G_2 \xrightarrow{f_2} G_3 \rightarrow \dots \xrightarrow{f_{n-1}} G_n$$

is said to be *exact* if it is exact at each joint, i.e., if  $\text{Im}f_i = \text{Ker}f_{i+1}$  for each  $i = 1, 2, \dots, n - 2$ . And, we say that an exact sequence

$$G_1 \xrightarrow{f_1} G_2 \xrightarrow{f_2} G_3 \rightarrow \dots \xrightarrow{f_{n-1}} G_n$$

*splits* at the group  $G_i, (i = 2, 3, \dots, n - 1)$ , iff the group  $G_i$  is the direct product of  $\text{Im}f_{i-1} = \text{Ker}f_i$  and another normal subgroup of  $G_i$ . Moreover,

if an exact sequence splits at each of its non-end groups, we say that it *splits* (or, it is a *split* exact sequence) ([5, p. 71]).

There is the following problem in [7, Problem 4.1 in p. 55]:

*Prove the fact that the following exact sequence splits;*

$$0 \hookrightarrow \mathfrak{g} = T_e G \hookrightarrow TG \xrightarrow{\pi} G \rightarrow 0.$$

Referring to the fact (4) of Theorem 2.1, it may be difficult for us to prove the problem on *split* of the short exact sequence above. Probably, I do not think that the exact sequence above splits in general.

The differential of the action of the group  $G$  on  $P(M, G, \pi)$  induces the right action of the group  $TG$  on  $TP$ , (i.e.,  $TP \times TG \rightarrow TP$ ). Similarly we may regard  $P$  as a subset of  $TP$ . Moreover, for  $A \in \mathfrak{g} \subset TG$  and  $u \in P \subset TP, uA (\in TP)$  makes sense. In fact,  $uA = A_u^* \in T_u P$ , where  $A^*$  is the fundamental vector field corresponding to  $A$  (cf. [4, 7]). The projection  $\pi : P \rightarrow M$  and the group operation  $u : G \times G \rightarrow G$  induce differentials  $\pi_* : TP \rightarrow TM$  and  $\mu_* : TG \times TG = T(G \times G) \rightarrow TG$ . Here,  $\mu_*(= \circ)$  is the group operation on the Lie group  $TG$ .

Since  $P$  is local trivial, for each point  $x \in M$  there exists a proper open neighbourhood  $U$  of the point  $x$  such that  $\pi^{-1}(U)$  is diffeomorphic with  $U \times G$ . Then there exists a cross section  $\sigma_U$  of  $U$  into  $\pi^{-1}(U) (\subset P)$ . Then  $\Psi|_{\pi^{-1}(U)} : \pi^{-1}(U) \ni u = \sigma_U(\pi(u)) \cdot g_U(\pi(u)) \rightarrow (\pi(u), \psi_U(u)) = (\pi(u), g_U(\pi(u))) \in U \times G$  is  $C^\infty$ -diffeomorphic.

Now, assume that  $U$  and  $V$  are open neighbourhoods in  $M$  with  $U \cap V \neq \emptyset$  such that  $\pi^{-1}(U)$  and  $\pi^{-1}(V)$  are diffeomorphic with  $U \times G$ , and  $V \times G$ , respectively. Then  $\sigma_V = \sigma_U \varphi_{UV}$  on  $U \times V$ , where  $\varphi_{UV} : U \cap V \rightarrow G$  is a transition function.

Moreover,  $TP \supset \pi_*^{-1}(TU) \supset T_u P \ni B \xrightarrow{\Psi_*} (\pi_*(B), dg_U(c(t))/dt|_{t=0}) \in TU \times TG$  is  $C^\infty$ -diffeomorphic, where  $B$  is the vector tangent to a curve  $\alpha(t)$  at point  $u = \alpha(0)$  in  $P$  such that  $\alpha(t) = \sigma_U(c(t)) \cdot g_U(c(t))$  and  $c(t) := \pi(\alpha(t))$ . Then,  $\sigma_U(c(t))g_U(c(t)) = \sigma_V(c(t))\varphi_{VU}(c(t))g_U(c(t)) = \sigma_V(c(t))g_V(c(t))$  and  $dg_V/dt = (d\varphi_{VU}/dt) \cdot g_U + \varphi_{VU} \cdot (dg_U/dt) \equiv (d\varphi_{VU}/dt) \circ (dg_U/dt)$ , where the operation  $\circ$  is the group operation on  $TG$ . Hence the differential  $d\varphi_{UV}$  of  $\varphi_{UV}$  is a transition function which is defined on  $TU \cap TV$ . Thus we obtain

**THEOREM 2.3.** *Let  $P(M, G, \pi)$  be a principal fibre bundle. Then  $TP(TM, TG, \pi_*)$  is a principal fibre bundle with group  $TG$  over the base manifold  $TM$ .*

## References

- [1] F. Dillen, K. Nomizu and L. Vrancken, *Conjugate connections and Radon's theorem in affine differential geometry*, Monat. Math. **109** (1990), 221-235.
- [2] S. Dragomir, T. Ichiyama and H. Urakawa, *Yang-Mills theory and conjugate connections*, Diff. Geom. Appl. **18** (2003), 229-238.
- [3] M. A. Guest, *Geometry of maps between generalized flag manifolds*, J. Diff. Geom. **25** (1987), 223-247.
- [4] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York (1978).
- [5] S. T. Hu, *Elements of Modern Algebra*, Holden-Day series in mathematics, San Francisco (1965).
- [6] M. Itoh, *Compact Einstein-Weyl manifolds and the associated constant*, Osaka J. Math. **35** (1998), 567-578.
- [7] S. Kobayashi, *Differential Geometry of Connections and Gauge Theory*, Shokabo (in Japanese), Tokyo, 1990.
- [8] A.B. Madsen, H. Pedersen, Y.S. Poon and A. Swann, *Compact Einstein-Weyl manifolds with large symmetry group*, Duke Math. J. **88** (1997), 407-434.
- [9] K. Nomizu and T. Sasaki, *Affine Differential Geometry-Geometry of Affine Immersions*, Cambridge, 1994.
- [10] J.-S. Park, *Stability of the identity maps of  $SU(3)/T(k,l)$* , Tokyo J. Math. **17** (1994), 281-289.
- [11] J.-S. Park, *New stable normal homogeneous Riemannian manifolds*, Kyushu J. Math. **53** (1999), 407-419.
- [12] J.-S. Park, *Yang-Mills connections in the orthonormal frame bundles over Einstein normal homogeneous manifolds*, Int. J. of Pure & Appli. Math. **5** (2003), 213-223.
- [13] J.-S. Park, *Critical homogeneous metrics on the Heisenberg manifold*, Interdisci. Inform. Sci. **11** (2005), 31-34.
- [14] J.-S. Park, *The conjugate connection of a Yang-Mills connection*, Kyushu J. Math. **62** (2008), 217-220.
- [15] J.-S. Park, *Yang-Mills connections with Weyl structure*, to appear.
- [16] J.-S. Park, *Projectively flat Yang-Mills connections*, to appear.
- [17] H. Pedersen, Y.S. Poon and A. Swann, *The Hitchin-Thorpe inequality for Einstein-Weyl manifolds*, Bull. London Math. Soc. **26** (1994), 191-194.
- [18] H. Pedersen, Y.S. Poon and A. Swann, *Einstein-Weyl deformations and submanifolds*, Internat. J. Math. **7** (1996), 705-719.
- [19] H. Pedersen and A. Swann, *Riemannian submersions, four-manifolds, and Einstein-Weyl geometry*, Proc. London Math. Soc. **66** (1993), 381-399.
- [20] H. Pedersen and A. Swann, *Einstein-Weyl geometry, the Bach tensor and conformal scalar curvature*, J. reine angew. Math. **441** (1993), 99-113.
- [21] H. Pedersen and K.P. Tod, *Three dimensional Einstein geometry*, Adv. Math. **97** (1993), 74-109.
- [22] K. P. Tod, *Compact 3-dimensional Einstein-Weyl structures*, J. London Math. Soc. **45** (1992), 341-351.
- [23] H. Urakawa, *Yang-Mills theory in Einstein-Weyl geometry and affine differential geometry*, Rev. Bull. Calcutta Math. Soc. **10** (2002), 7-18.

\*

Department of Mathematics  
Pusan University of Foreign Studies  
Pusan 55-1, Republic of Korea  
*E-mail:* [iohpark@pufs.ac.kr](mailto:iohpark@pufs.ac.kr)