



True Digestibility of Phosphorus in Different Resources of Feed Ingredients in Growing Pigs

X. Wu¹, Z. Ruan^{2,3}, Y. G. Zhang^{1,a}, Y. Q. Hou³, Y. L. Yin^{1,2,3,*}, T. J. Li¹, R. L. Huang¹
W. Y. Chu¹, X. F. Kong¹, B. Gao⁴ and L. X. Chen⁵

¹Laboratory of Animal Nutrition and Health and Key Laboratory of Subtropical Agro-ecology, Institute of Subtropical Agriculture, The Chinese Academy of Sciences, Changsha, Hunan Province, 410125, China

ABSTRACT : To determine the true digestible phosphorus (TDP) requirement of growing pigs, two experiments were designed with the experimental diets containing five true digestible P levels (0.16%, 0.20%, 0.23%, 0.26% and 0.39%) and the ratio of total calcium to true digestible P (TDP) kept at 2:1. In Experiment 1, five barrows (Duroc×Landrace×Yorkshire) with an average initial body weight of 27.9 kg were used in a 5×5 Latin-square design to evaluate the effect of different dietary P levels on the digestibility and output of P and nitrogen. In Experiment 2, sixty healthy growing pigs (Duroc×Landrace×Yorkshire) with an average body weight (BW) of 21.4 kg were assigned randomly to one of the five dietary treatments (12 pigs/diet), and were used to determine the true digestible phosphorus (TDP) requirement of growing pigs on the basis of growth performance and serum biochemical indices. The results indicated that the true digestibility of P increased ($p < 0.05$) linearly with increasing dietary TDP level below 0.26%. The true P digestibility was highest (56.6%) when dietary TDP was 0.34%. Expressed as g/kg dry matter intake (DMI), fecal P output increased ($p < 0.05$) linearly with increasing P input. On the basis of g/kg fecal dry matter (DM), fecal P output was lowest for Diet 4 and highest ($p < 0.05$) for Diet 5. The apparent digestibility of crude protein (CP) did not differ ($p > 0.05$) among the five diets, with the average nitrogen output of 12.14 g/d and nitrogen retention of 66% to 74% ($p > 0.05$), which suggested that there was no interaction between dietary P and CP protein levels. During the 28-d experimental period of Experiment 2, the average daily gain (ADG) of pigs was affected by dietary TDP levels as described by Eq. (1): $y = -809.532x^4 + 788.079x^3 - 276.250x^2 + 42.114x - 1.759$; ($R^2 = 0.99$; $p < 0.01$; $y = \text{ADG, g/d}$; $x = \text{dietary TDP, \%}$), F/G for pigs by Eq. (2): $y = 3.651.1x^4 - 3.480.4x^3 + 1.183.8x^2 - 172.5x + 10.9$ ($R^2 = 0.99$; $p < 0.01$; $y = \text{F/G}$; $x = \text{dietary TDP, \%}$), and Total P concentrations in serum by Eq. (3): $y = -3.311.7x^4 + 3.342.7x^3 - 1.224.6x^2 + 195.6x - 8.7$ ($R^2 = 0.99$; $p < 0.01$; $y = \text{total serum P concentration}$ and $x = \text{dietary TDP, \%}$). The highest ADG (782 g/d), the lowest F/G (1.07) and the highest total serum P concentration (3.1 mmol/L) were obtained when dietary TDP level was 0.34%. Collectively, these results indicate that the optimal TDP requirement of growing pigs is 0.34% of the diet at a total Ca to TDP ratio of 2:1. (**Key Words :** True Digestible Phosphorus, Growing Pigs, Growth Performance, Biochemical Indices, Calcium)

INTRODUCTION

As the second most abundant mineral in the animal

body, phosphorus (P) has many metabolic and physiological functions, including energy utilization and transfer, cell differentiation and growth, cell signaling and nutrient metabolism, development and maintenance of skeletal tissue, intestinal integrity and function, protein synthesis and maintenance of osmotic pressure and acid-base balance (Anderson, 1991; Wu, 1998). Symptoms of P deficiency vary but, in all cases, they are associated with reduced feed intake, impaired growth, and compromised health in animals. P deficiency will ultimately result in a reduction in plasma phosphate concentrations, followed by resorption of Ca and P from bones, thereby minimizing weight gain and growth performance of livestock.

Phytate P is the major form, however, it is poorly

* Corresponding Author: Y. L. Yin. Tel: +86-731 4619703, Fax: +86-731-4612685, E-mail: yinyulong@isa.ac.cn

²The key Laboratory of Food Science of Ministry of Education of China, Department of Food Science & Engineering, Nanchang University, Nanchang, China.

³Department of Feedstuff Science, Wuhan Polytechnic University, Wuhan, Hubei 430023, China.

⁴Guangzhou Tanke Industry Co. Ltd., Guangzhou, 510627, China.

⁵Guang An Biological Technique Company, Changsha, Hunan Province, 410129, China.

^aMade the same contribution as the first author.

Received April 15, 2007; Accepted May 4, 2007

Table 1. Fecal apparent and true digestibility of phosphorus in some feedstuffs (%)

Feedstuffs	Fecal apparent digestibility	Fecal true digestibility
Corn (Yellow)	24.31±5.16	49.15±4.16
Corn (White)	17.16±5.78	40.33±5.24
Paddy meal	32.20±4.16	43.65±3.88
Barley meal	35.61±4.09	56.68±3.56
Wheat meal	32.70±4.10	49.74±4.92
Oat meal	35.98±5.90	27.45±3.83
Buckwheat meal	30.36±6.23	49.72±5.43
Sorghum meal	31.02±5.89	42.34±5.35
Rough rice meal	28.61±5.67	41.89±5.84
Broken rice meal	31.91±4.88	44.62±3.22
Rough rice bran	11.82±4.09	23.41±3.44
Defatted rice bran	14.54±4.09	26.08±4.06
Wheat middlings meal	40.19±4.89	63.93±4.42
Wheat bran	41.25±3.99	56.82±1.59
Soybean meal	27.59±3.09	51.30±2.61
Cottonseed meal	15.02±5.90	30.22±4.61
Rapeseed meal	27.67±4.00	39.30±2.50
Field bean meal	31.60±5.89	50.76±5.86
Peas meal	26.33±7.44	36.56±6.50

Data are means±SEM, n = 5.

digested by pigs due to a lack of phytase (Jongbloed and Lenis, 1998). So producers always add excessive phosphate in feed in order to fill the requirement of pigs, which can

result in the following problems: (1) excessive P output in swine manure enters our waterways where it can stimulate algal blooms and eutrophication. (Mallin, 2000; Pouslen, 2000; Sacakli et al., 2006; Fang et al., 2007a); (2) excessive P in feed adds to the cost of swine production; (3) as an anti-nutrient factor, inorganic P combines with Ca and fat in diets into an insoluble compound -saponin- which decreases the digestibility of other nutrients; and (4) as a crucial component of all forms of life on earth, inorganic phosphate is a limited and non-renewable natural resource and the conservation of P has become a global issue (Abelson, 1999). Thus, research on determining P requirement accurately and improving the efficiency of its utilization by pigs has become an important issue in swine nutrition (Bhanja et al., 2005; Shen et al., 2005; Ruan et al., 2007).

Considerable research has been conducted to assess the requirements of total P by swine (Miller et al., 1964; Cromwell et al., 1972; Lin et al. 2002; Ruan et al., 2007). Because P bioavailabilities vary greatly among feed ingredients and endogenous P and nitrogen (N) were included in the feces, so the apparent digestibility underestimated the biological values of amino acid and P, only true digestibility reflects the process of digestion and absorption. So P requirements should be based on true digestible P (TDP) rather than total P or available P in feeds. Despite considerable progress in many areas of swine nutrition, TDP has not been previously established for pigs

Table 2. Composition of experimental diets and nutrient indexes for growing pigs

Ingredients (%)	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5
Corn	57.80	57.80	57.80	57.80	57.80
Soybean meal	22.10	22.10	22.10	22.10	22.10
Comstarch	2.04	1.80	1.55	1.30	0.17
Rice bran meal	5.00	5.00	5.00	5.00	5.00
Cottonseed meal	3.00	3.00	3.00	3.00	3.00
Wheat middling meal	5.00	5.00	5.00	5.00	5.00
Soybean oil	3.00	3.00	3.00	3.00	3.00
Lys	0.13	0.13	0.13	0.13	0.13
NaCl	0.30	0.30	0.30	0.30	0.30
CaCO ₃	0.63	0.67	0.72	0.77	1.00
CaHPO ₄	0.00	0.20	0.40	0.60	1.50
Premix ¹	1.00	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00	100.00
Calculated nutritive values:					
Crude protein (%)	17.34	17.34	17.34	17.34	17.34
Digestible energy (MCal/kg)	3.43	3.42	3.41	3.40	3.35
Total Ca (%)	0.32	0.40	0.46	0.52	0.78
Total phosphorus (%)	0.44	0.47	0.51	0.54	0.69
True digestible phosphorus (%)	0.16	0.20	0.23	0.26	0.39
Total Ca:true digestible P	2:1	2:1	2:1	2:1	2:1
True digestible Lys (%)	0.83	0.83	0.83	0.83	0.83
True digestible (Met+cys) (%)	0.51	0.51	0.51	0.51	0.51
True digestible Thr (%)	0.58	0.58	0.58	0.58	0.58
True digestible Trp ¹² (%)	0.19	0.19	0.19	0.19	0.19

¹ Supplied the following vitamins and minerals (mg/kg diet): retinyl palmitate, 5.2; cholecalciferol, 0.38; all-rac- α -tocopherol acetate, 44.0; menadione, 3.0; riboflavin, 2.2; niacin, 12.0; d-pantothenic acid, 11.0; vitamin B-12, 0.012; thiamine, 1.1; choline chloride, 550.0; pyridoxine, 1.1; d-biotin, 0.1; and folic acid, 0.6; FeSO₄·H₂O, 152; ZnCO₃, 95.9; MnSO₄·H₂O, 6.2; CuSO₄·5H₂O, 11.8; KI, 0.6; Na₂SeO₃, 0.3.

at any stage of the life cycle. Thus the objective of these studies was to determine the TDP requirement of growing pigs and to study the effect of dietary P levels on the digestibility and output of P and N. The studies offer a scientific basis for making diets with low pollution characteristics.

MATERIALS AND METHODS

Digestibility trial and experimental diets

Knowledge about the true digestibility of P in feed ingredients is necessary for designing a sound experiment to determine the TDP requirements for swine growth. Thus, the fecal apparent and true digestibility of P in 19 major feedstuffs of China were determined over a 2-year period in our laboratory. These studies were conducted using a 5×5 Latin square design, according to the methods of Yin et al. (Yin et al., 1995; Yin and McCracken, 1996; Yin et al., 2000a, b, c; 2002a, b) Fan et al. (2005) and Fang et al. (2007a, b). The results are summarized in Table 1 as a basis for preparing diets used in the trials.

Five corn- and soybean meal-based diets were formulated on the basis of the TDP values of the ingredients mentioned above to provide five levels of TDP: 0.16%, 0.20%, 0.23%, 0.26% and 0.39% and the ratio of total calcium to TDP was 2:1 (Table 2). Except for Ca and P, other nutrients were adequate for growing pigs (NRC, 1998) and were similar among the experimental diets. Chromic oxide (0.4%) was added as an inert marker for estimation of the coefficient of total tract apparent digestibility (Huang et al., 2005; Yang et al., 2005).

Animals, experimental design and management

In Experiment 1, five Duroc×Landrace×Yorkshire barrows, with an average initial body weight (BW) of 27.86 kg, were fitted with a simple T-cannula at the distal ileum. The pigs were housed individually in stainless steel metabolism crates in a temperature-controlled room (20–25°C) and fed the five experimental diets (Table 2) according to a 5×5 Latin square design. The experiment had 5 periods and during a 7-d recovery period, the pigs were fed a regular grower diet. During the experimental period the pigs were fed twice daily, equal amounts for each meal at 08:00 and 20:00 h (Yin et al., 1993). The dietary allowance was 4% of body weight (Yin et al., 2001a, b, c, d; Yin et al., 2004).

In Experiment 2, sixty healthy cross-bred pigs (Duroc×Landrace×Large White) with an average BW of 21.4 kg were assigned randomly to one of the five dietary treatments (Table 2). Each treatment consisted of 3 replications, with 4 pigs (2 barrows and 2 gilts) in each replication. Each pen housed 2 pigs. Feeds were offered to pigs at 5% BW for 28 d (Yin et al., 1991a, b; Yin et al.,

1993a, b; Yin et al., 1994; Yin 1994; Yang et al., 2007).

Drinking water was freely available to all the pigs via low-pressure drinking nipples.

Sample collection and treatment

In Experiment 1, each experimental period comprised 8 d. After a 4-d adaptation, all possible fecal samples were collected on d 5, 6, 7 and 8. The ileal digesta samples were collected in soft plastic boxes which were attached to the barrel of the cannula with Velcro tape. The box contained 10ml of a solution of formic acid (2.86 ml/L) to minimize further bacterial activity and was replaced as soon as it was partially filled with sample. Samples were immediately frozen at -20°C. Then samples were freeze-dried, pooled within the same barrow and period of the same diet, ground through a 1-mm mesh screen and mixed before analysis. Samples of the diets were ground similarly. Analyses were performed in duplicate (Huang et al., 2003).

In Experiment 2, the BW of pigs was measured after a 12-h fast in the morning of the first and last days of the growth trial to calculate average daily gain (ADG). Feed intake was recorded daily during the 28-d period to determine average daily feed intake (ADFI) and the feed:gain ratio(F/G). On the 1st and 28th day of the experiment, the fasted pigs were weighed, and jugular venous blood samples (10 ml) were obtained from 1 pig of each pen into heparin-free tubes, as previously described (Tang et al., 2005). Blood samples were centrifuged at 4,000×g for 15 min. The supernatant fluid (serum) was collected and immediately frozen at -20°C for biochemical analyses (Deng et al., 2007a, b; Kong et al., 2007a, b, c).

Chemical analysis

Analysis for dry matter (DM) was carried out according to AOAC methods. Chromic oxide was determined (Saha and Gilbreath, 1991) by using an atomic absorption spectrometer (SpectrAA-10/20, Varian, Mulgrave, Australia). Diet and fecal samples were weighed into 60-ml Pyrex beakers and ashed overnight at 550°C. Cr₂O₃, as part of the ash, was then oxidized to dichromate by digestion in 6 ml of phosphoric acid (16.7 mol/L)-manganese sulfate (13.5 m mol/L) solution mixed with 8 ml of potassium bromate (0.27 mol/L) solution on a hot plate. Potassium dichromate was used as standard (Deng et al., 2007a, b; Huang et al., 2007).

Analyses of total inorganic P in samples were carried out by spectrophotometric analysis at 355 nm (Heinoen and Lahti, 1981). Potassium monobasic phosphate was used as standard.

An Automated Biochemistry Analyzer (Synchron CX Pro, Beckman Coulter, Fullerton, CA, USA) was used to determine the concentrations of total P, total Ca and alkaline phosphatase in serum (Kong et al., 2007a, b; Li et al., 2007). All the kits were purchased from Beijing Chemlin Biotech

Co., Ltd (Beijing, China).

Estimation principles

Determination of the gastrointestinal endogenous P outputs by the regression analysis technique relies on establishing linear relationships between P outputs in feces and their dietary inputs (Fan and Sauer, 1999).

The total outputs of P in feces, expressed as g/kg dry matter diet intake (DMI), are calculated from Eq. (1) according to the previous studies.

$$P_O = P_I \times (I_D / I_f) \quad (1)$$

where P_O represents the outputs of P in feces (g/kg DMI), P_I is the content of P in feces (g/kg DM feces), I_D is the Cr_2O_3 concentration in the diets (g/kg DMI) and I_f is the Cr_2O_3 concentration in feces (g/kg DM feces).

The outputs of P in feces have both dietary and endogenous origins. If there are linear relationships between P outputs in feces and the graded levels of P inputs from diets, when expressed as g/kg DMI, their relationships can be expressed according to Eq. (2).

$$P_{O_i} = P_E + D_i \times P_{D_i} \quad (2)$$

where P_{O_i} represents the output of P in feces collected from animals fed the *i*th assay diet, determined using Eq. (1) (g/kg DMI). P_E represents the level of the gastrointestinal endogenous P in feces (g/kg DMI), D_i is the percentage of dietary P that is truly indigestible going through the gastrointestinal tract (%) and P_{D_i} is the P content in the *i*th assay diet (g/kg DMI). D_T is the true fecal P digestibility values (%) in the P-containing assay ingredient and can be calculated according to Eq. (3). once D_i is estimated from regression analysis according to Eq. (2).

$$D_T = 100\% - D_i \quad (3)$$

Eq. (2) represents a simple linear regression model in which P_{O_i} and P_{D_i} are the dependent and independent variables, respectively. P_E and D_i are the regression coefficients and are estimated by fitting the simple linear regression model. If there are linear relationships between P outputs in feces and graded levels of P inputs from the diets with significant intercepts, then the endogenous P level in feces can be determined directly by extrapolating the dietary inputs of P to zero by obtaining the intercepts of the linear regression Eq. (P_E).

To determine true fecal P digestibility values in a P-containing ingredient, a series of assay diets are formulated to contain graded dietary levels of P, but only from the

assay ingredient. The contents of other dietary factors, such as anti-nutritive factors that likely affect P digestion and endogenous P outputs, should be controlled between the assay diets.

The apparent fecal digestibility values of DM and P in the experimental diets were calculated according to Eq. (4).

$$D_{A_i} = 100\% - ((I_D \times P_f) / (I_f \times P_D)) \times 100\% \quad (4)$$

where D_{A_i} represents the apparent fecal P digestibility values in the assay diets (% on as-fed basis), I_D is the digestibility marker concentration in the *i*th assay diet (% on as-fed basis), P_f is the P concentration in feces (% on as-fed basis), P_D is the P concentration in the *i*th assay diet (% on as-fed basis) and I_f is the digestibility marker concentration in feces (% on as-fed basis).

On the basis of the apparent ileal and fecal P digestibility values and the levels of endogenous P extrapolated with regression analysis, the true P digestibility values in the assay diets and also in the test ingredient, SBM, can be determined according to Eq. (5).

$$D_{T_i} = D_{A_i} + (P_E / P_{D_i}) \times 100\% \quad (5)$$

Alternatively, the endogenous P outputs corresponding to individual diets can also be calculated according to Eq. (6), if corresponding true ileal and fecal P digestibility values are determined.

$$P_E = ((D_{T_i} - D_{A_i}) \times P_{D_i}) / 100\% \quad (6)$$


where D_{T_i} represents the true fecal P digestibility values in the assay diets (%). D_{A_i} represents the apparent fecal P digestibility values in the assay diets (%), P_E represents the levels of endogenous P in feces (g/kg DM diet intake) and P_{D_i} is the P concentration in the assay diets (g/kg DM diet).

Statistical analyses

The digestibility values were first subjected to ANOVA for a 5×5 Latin square design. The treatment effect was therefore partitioned and tested according to equally spaced orthogonal polynomial analyses (Steel and Torrie, 1980). The ANOVA and the orthogonal polynomial analyses were carried out using the General Linear Procedures of SAS (SAS, 2000). Related linear and curvilinear regression analyses were conducted using the Fig.P program (FigP Software Corporation, 1992). The comparison of true P digestibility values and the endogenous P outputs between feces was conducted according to the pooled t test (Byrkit, 1987). Pen was the experimental unit in the statistical analysis. For the orthogonal contrast among the treatment groups, the linear and quadratic effects were tested using

Table 3. Dry matter, crude protein, apparent and true phosphorus digestibility values in experimental diets (%)

Items	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P
Dietary true digestible phosphorus concentration	0.16	0.20	0.23	0.26	0.39	-	-
Dry matter digestibility values	91.74	91.03	90.88	90.04	91.00	0.75	0.6315
Apparent crude protein digestibility values	80.86	81.55	82.60	81.76	81.23	0.85	0.6794
Apparent phosphorus digestibility values	25.60	31.05	32.57	34.49	34.59	2.16	0.0676
True phosphorus digestibility values	48.40	51.03	51.68	52.69	49.56	2.16	0.6567

**Figure 1.** The effects of dietary true digestible phosphorus concentration on the apparent and true phosphorus digestibility.


SAS (2000). Probability values of <0.05 were taken to indicate statistical significance.

RESULTS

All the pigs remained healthy and symptoms of Ca and P deficiency were not found. There was no effect of individual animal and experimental period on the result according to the statistical analysis.

The digestibility of P and endogenous output

The apparent P digestibility could be calculated according to Eq. (4) and increases with increasing dietary TDP concentration in growing pigs. The apparent P digestibility values increased from 25.60% to 34.59%

**Figure 2.** Linear relationship between phosphorus output in feces and dietary phosphorus input.

(Table 3 and Figure 1).

When expressed as g/kg DMI, the fecal P output could be calculated according to Eq. (1). Fecal P output increased significantly ($p = 0.0002$) linearly with increasing dietary TDP input (Table 4). So it can be concluded that fecal P output was influenced by the dietary TDP concentration. Endogenous P output was calculated according to Eq. (5). Endogenous P output of growing pigs was calculated as 1.14 g/kg DMI in Experiment 1.

True P digestibility was calculated according to Eq. (5) or Eq. (6). From Table 3 and Figure 1, true P digestibility increased with increasing dietary TDP concentration when dietary TDP concentration was below 0.26%. When dietary TDP concentration increased from 0.26% to 0.39%, the true P digestibility decreased from 52.69% to 49.56% and the true P digestibility was highest when dietary TDP

Table 4. Dietary phosphorus input and the partitioning of phosphorus flow in feces

Items	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P
Total dietary phosphorus intake (g/kg DMI)	5.00	5.71	5.97	6.27	7.26	-	-
Apparent digestible phosphorus in feces (g/kg DMI)	1.28 ^a	1.77 ^b	1.94 ^b	2.16 ^b	2.64 ^c	0.13	0.0001
True digestible phosphorus in feces (g/kg DMI)	2.42 ^a	2.91 ^b	3.01 ^b	3.30 ^b	3.78 ^c	0.13	0.0001
Total fecal phosphorus output							
g/kg feces DM	21.86 ^{ab}	23.77 ^c	23.21 ^{bc}	21.28 ^a	25.82 ^d	0.49	0.0002
g/kg DMI	3.72 ^a	3.94 ^a	4.02 ^a	4.11 ^a	4.99 ^b	0.13	0.0002
Endogenous fecal phosphorus output							
g/kg feces DM	7.35	6.70	6.20	5.15	6.27	0.58	0.1772
g/kg DMI	1.25	1.12	1.08	1.01	1.22	0.13	0.6710
Fecal phosphorus output of dietary origin							
g/kg feces DM	15.15 ^a	16.86 ^a	16.62 ^a	15.33 ^a	19.67 ^b	0.44	0.0001
g/kg DMI	2.58 ^a	2.80 ^a	2.88 ^a	2.96 ^a	3.84 ^b	0.13	0.0002

Values in the same row with different superscript letters differ significantly ($p < 0.05$).

Table 5. The balance of phosphorus and nitrogen for growing pigs

Items	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P
Dietary true digestible phosphorus concentration (%)	0.16	0.20	0.23	0.26	0.39	-	-
Dietary phosphorus input (g/d)	6.45 ^a	7.36 ^b	7.84 ^b	8.02 ^b	9.62 ^c	0.18	0.0001
Fecal phosphorus output (g/d)	3.38 ^a	3.55 ^a	3.86 ^a	3.98 ^a	5.39 ^b	0.23	0.0004
Urinary phosphorus output (g/d)	0.025 ^a	0.034 ^b	0.035 ^b	0.042 ^c	0.072 ^d	0.0020	0.0001
Total phosphorus output (g/d)	3.40 ^a	3.58 ^a	3.90 ^a	4.02 ^a	5.47 ^b	0.23	0.0003
Detention phosphorus (g/d)	3.05 ^a	3.18 ^b	3.94 ^b	4.00 ^b	4.26 ^b	0.17	0.0043
Phosphorus retention (%)	47.67	51.26	50.29	49.90	43.30	2.77	0.3715
Dietary nitrogen input (g/d)	40.85	40.53	41.73	40.41	39.85	0.91	0.6829
Fecal nitrogen output (g/d)	6.17	6.92	6.47	6.01	4.78	0.79	0.4402
Urinary nitrogen output (g/d)	6.07	6.57	6.28	5.76	5.69	0.48	0.6857
Total nitrogen output (g/d)	12.24	13.49	12.74	11.77	10.47	0.92	0.2596
Retentive nitrogen (g/d)	28.61	27.04	28.99	28.64	29.38	1.03	0.5790
Nitrogen retention (%)	70.25	66.60	69.46	70.92	73.64	2.13	0.2834

Values in the same row with different superscript letters differ significantly ($p < 0.05$).

concentration was 0.34% (Figure 1). There were no significant differences among the 5 diets.

Fecal P flow

As shown in Table 4, apparent fecal digestible P output increased with increasing dietary TDP concentration. Apparent fecal digestible P output increased from 1.28 g/kg DMI to 2.64 g/kg DMI. The apparent fecal digestible P output of diet 1 was significantly lower than diet 2 ($p = 0.022$) and highly significantly lower than diet 3 ($p = 0.007$), diet 4 ($p = 0.003$) and diet 5 ($p = 0.001$). The apparent fecal digestible P output of diets 2, 3 and 4 were highly significantly lower than diet 5 ($p < 0.001$).

True fecal digestible P output increased from 2.42 g/kg DMI to 3.78 g/kg DMI with increasing dietary TDP concentration. The true fecal digestible P output of diet 1 was significantly lower than diet 2 ($p = 0.022$) and highly significantly lower than diets 3 ($p = 0.007$), 4 ($p = 0.003$) and 5 ($p = 0.001$). The true fecal digestible P output of diets 2, 3 and 4 were highly significantly lower than diet 5 ($p < 0.001$).

Expressed as g/kg DMI, fecal P output increased linearly with increasing dietary P input. The linear equation was $y = 0.493x + 0.1409$, $R^2 = 0.9574$, where x represents dietary P input (g/kg DMI), y represents fecal P output (g/kg DMI). When x was extrapolated to zero, the y value obtained from the intercept of the linear regression equation was endogenous P output (Figure 2). The average endogenous P output of growing pigs in the experiment was 1.14 g/kg DMI. The fecal P output of diet 5 was very significantly higher than diets 1, 2, 3 and 4 ($p < 0.001$). The differences among other diets were not significant (Table 4). The ratio of fecal P output to dietary P input decreased from 74.4% to 65.5%, then increased to 68.7% with increasing dietary TDP concentration.

Expressed as g/kg feces, there was no clear relationship in fecal P output with increasing dietary TDP concentration

and diet 5 was higher than diets 1 ($p < 0.001$), 2 ($p = 0.021$), 3 ($p = 0.005$) and 4 ($p < 0.001$). The fecal P output of diet 4 was very significantly ($p < 0.001$) lower than diets 2 and 5 and significantly ($p = 0.029$) lower than diet 3. The fecal P output of diet 1 was significantly ($p = 0.03$) lower than diet 2.

After calculating the true P digestibility, endogenous P output could be calculated according to Eq. (5) or (6). Endogenous P output decreased with increasing dietary P input when dietary P input was lower than 6.27 g/kg DMI. When dietary P input increased from 6.27 g/kg DMI to 7.26 g/kg DMI, endogenous P output increased from 1.01 g/kg DMI to 1.22 g/kg DMI. There were no significant differences among experimental diets in endogenous P output and fecal P output of dietary origin of diet 5 was very significantly ($p < 0.001$) higher than the other 4 diets. Expressed as g/kg DM feces, fecal P output of dietary origin of diet 5 was very significantly ($p < 0.001$) higher than the other 4 diets, while expressed as g/kg DMI, faecal P output of dietary origin increased with increasing dietary P input.

P balance

Fecal P output of pigs fed diet 5 was very significantly ($p < 0.01$) higher than with the other 4 diets. Fecal and urinary P output increased with increasing daily dietary P input. Urinary P output of diet 5 was very significantly ($p < 0.001$) higher than diet 4. Urinary P output of diet 4 was significantly higher than diets 2 ($p = 0.034$) and 3 ($p = 0.046$) and very significantly higher than diet 1 ($p < 0.001$). There were differences of 2.02 g/d between the highest (diet 5) and lowest diets (diet 4) in fecal P output and 0.047 g/d in urinary P output (Table 5). Total P output was influenced by fecal P output and urinary P output, so it increased with increasing daily dietary P input, and the highest value was 2.07 g/d higher than the lowest ($p < 0.01$).

Retention of P in the body increased with increasing daily dietary P input. Retained P of diet 1 was very significantly ($p < 0.01$) lower than the other 4 diets. There




Figure 3. The effects of dietary true digestible phosphorus concentration on the apparent crude protein digestibility values.

was no clear relationship for P retention in the body with values from high to low being diet 2, 3, 4, 1 and 5. The highest value was 7.96% higher than the lowest, but there were no significant differences among the diets.

Apparent N digestibility and N balance

The average apparent N digestibility was 81.6%. There was no significant effect of dietary TDP concentration on apparent N digestibility among diets (Table 3). The effect of dietary TDP concentration on apparent N digestibility showed a parabolic trend (Figure 3). The highest point of apparent N digestibility was evident when dietary TDP concentration was 0.23%. The apparent N digestibility ranged from high to low on diets 3, 4, 2, 5 and 1 with respective values of 82.60%, 81.76%, 81.55%, 81.23% and 80.86%.

There was no relationship in fecal N, urine N and total N output with increasing dietary N input and there were no significant differences among the diets (Table 5). The amount of retained N decreased initially and then increased with increasing dietary N input (Figure 4). The lowest point of retained N in the body was evident when dietary TDP




Figure 4. The relationship between dietary nitrogen input and nitrogen retention.

concentration was 0.20%, and its value was 27.04 g/d. There were no significant differences in retained N among the 5 diets. The highest point was evident at 29.38 g/d when dietary TDP concentration was 0.39%.

The retention of N in the body decreased with increasing dietary P input from 7.36 to 9.62 g/d, but the differences among diets were not significant. The lowest value was observed at a dietary TDP concentration of 0.20% and the highest value was present at a dietary TDP concentration of 0.39% (Table 5).

Growth performance

Feeding different levels of TDP had no linear effects ($p > 0.05$) on ADG, ADFI, or the F/G in growing pigs during the 28-d trial (Table 6). However, dietary TDP levels exhibited a quadratic effect ($p < 0.05$) on ADG and the F/G (Table 6). An equation was developed to relate the ADG of pigs to dietary TDP level: $y = -809,532x^4 + 788,079x^3 - 276,250x^2 + 42,114x - 1,759$; $R^2 = 0.99$; $p < 0.01$; where x = dietary TDP level (%) and y = ADG of pigs (g/d). The ADG of pigs was the greatest (750 g/d) when the dietary TDP

Table 6. The effects of dietary true digestible phosphorus levels on growth performance in growing pigs

Items	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P (linear)	P (quadratic)
Total phosphorus (%)	0.44	0.47	0.51	0.54	0.69			
Apparent phosphorus (%)	0.14	0.17	0.20	0.23	0.37			
True digestible phosphorus (%)	0.16	0.20	0.23	0.26	0.39			
Total calcium (%)	0.33	0.38	0.45	0.51	0.79			
Initial weight (kg)	21.38	21.58	21.34	21.36	21.29	0.35	0.973	0.995
Final weight (kg)	38.31	39.04	39.18	40.08	40.00	1.06	0.753	0.458
Average daily feed intake (g/d)	1,029	1,059	1,069	1,060	1,062	88	0.998	0.700
Total phosphorus intake (g/d)	4.5 ^a	4.9 ^a	5.5 ^{ab}	5.7 ^{ab}	7.3 ^b	0.49	0.019	0.219
Apparent phosphorus intake (g/d)	1.4 ^a	1.8 ^{ab}	2.1 ^{ab}	2.4 ^b	3.9 ^c	0.22	0.001	0.308
True digestible phosphorus intake (g/d)	1.6 ^a	2.1 ^{ab}	2.4 ^{ab}	2.7 ^b	4.1 ^c	0.24	0.001	0.209
Average daily gain (g/d)	604 ^b	623 ^b	637 ^{ab}	668 ^a	668 ^a	34	0.640	0.045
Feed:gain ratio	1.71 ^u	1.72 ^u	1.69 ^a	1.56 ^b	1.67 ^a	0.14	0.984	0.0344

Values in the same row with different superscript letters differ significantly ($p < 0.05$).

Data are means with pooled SEM for 12 pigs per treatment group.

Table 7. The effects of dietary true digestible phosphorus levels on serum biochemical indices

Items	Diet1	Diet 2	Diet3	Diet4	Diet5	SEM	P (linear)	P (quadratic)
Total phosphorus (%)	0.44	0.47	0.51	0.54	0.69	-	-	-
Apparent phosphorus (%)	0.14	0.17	0.20	0.23	0.37	-	-	-
True digestible phosphorus (%)	0.16	0.20	0.23	0.26	0.39	-	-	-
Total calcium (%)	0.33	0.38	0.45	0.51	0.79	-	-	-
Serum alkaline phosphatase activity (U/L)	186	183	168	166	159	18	0.808	0.782
Total serum phosphorus concentration (mmol/L)	2.72 ^b	2.83 ^b	2.86 ^b	2.94 ^a	2.94 ^a	0.11	0.565	0.049
Total serum Ca concentration (mmol/L)	2.57	2.63	2.66	2.75	2.67	0.05	0.147	0.521

^{a,b} Values in the same row with different superscript letters differ ($p < 0.05$).

Data are means with pooled SEM for 6 pigs per treatment group.

level was 0.34%. Notably, the ADG of pigs decreased ($p < 0.05$) when dietary TDP levels exceeded 0.34%.

The F/G of growing pigs decreased ($p < 0.05$) from 1.71 to 1.56 when dietary TDP levels increased from 0.16% to 0.26%, but increased ($p < 0.05$) from 1.56 to 1.67 when dietary TDP level was elevated from 0.26% to 0.39%. The relationship between the F/G for pigs and dietary TDP levels could be described by the following equation: $y = 3,651.1x^4 - 3,480.4x^3 + 1,183.8x^2 - 172.5x + 10.9$, $R^2 = 0.99$, $p < 0.01$; where x = dietary TDP level (%) and y = the F/G for growing pigs. It was calculated that the F/G was the lowest (1.07) when the dietary TDP level was 0.34%.

Serum biochemical indices

Neither alkaline phosphatase activities nor total Ca concentrations in serum differed ($p > 0.05$) when dietary TDP levels increased from 0.16% to 0.39% (Table 7). However, feeding different levels of TDP had a quadratic effect ($p < 0.05$) on total P concentrations in serum (Table 3). A regression equation relating total serum P concentrations to dietary TDP levels was established: $y = -3,312x^4 + 3,343x^3 - 1,225x^2 + 196x - 9$; $R^2 = 0.99$, $p < 0.05$; where x = dietary TDP level (%) and y = total serum P concentration (mmol/L). Total P concentration in serum was the highest (3.1 mmol/L; $p < 0.05$) when dietary TDP was 0.34% (or 5.1 g/d for a 30-kg pig that consumed 1.5 kg feed/d). Total P concentration in serum (0.22 mmol/L or 1.36 mg/100 ml) was lower ($p < 0.05$) in pigs fed a diet containing 0.16% TDP than on diets providing 0.26% and 0.39% TDP.

DISCUSSION

Determination of true P digestibility and endogenous P output

There are several disadvantages in using apparent P digestibility to estimate P efficiency, and only true P digestibility value can reflect the actual digestion and absorption of P in the diet. Apparent P digestibility underestimates the biological values of P because endogenous P contribution to intestinal or fecal P is not taken into consideration, so apparent fecal P digestibilities

were 20% to 25% lower than true P digestibility for feed ingredients (Table 1). Apparent P availability of the same feed ingredient may vary considerably (as much as 15% to 35%) with swine diets (Weremko et al., 1997; Fan et al., 2001 and 2002; Shen et al., 2002; Fang et al., 2007). Further, apparent P digestibility values are not always additive in single feed ingredients for growing pigs (Fan et al., 2002; Fang et al., 2007). In contrast, available evidence shows that true P digestibility values are additive in ingredients containing low levels of phytate phosphorus and antinutritional factors (Fang et al., 2007). Compared with the traditional total dietary P content and apparent P digestibility systems, the use of TDP in formulating swine diets offers a distinct advantage of accuracy in meeting P requirements. Results from the present study indicate that changes in the ADG of growing pigs and total serum P concentrations were positively correlated with dietary TDP levels but not with total dietary P or fecal digestible P (Tables 6 and 7).

Exact determination of endogenous P output in animals is the key to determining the true P digestibility. However, it is difficult to determine true P digestibility because P recirculated in the body. The traditional method of determining endogenous P output is to feed animals a diet with no P, then the fecal P output is only endogenous P, but animals fed diets without P will suffer perturbations of digestion and metabolism caused by the shortage of P. The endogenous P output determined under these conditions is far different from that under normal conditions. Fan et al. (2001) found that piglets suffered diarrhea and shiver after feeding a no-P diet for 5 to 7 days. ³²P tracer has been used to determine endogenous P output, but fast recycling of P in the gastrointestinal tract creates difficulty in handling radioactive material in routine whole animal studies, so this area of research has been relatively unexplored (Fan et al., 1999). Fan et al. (1999) have successfully determined endogenous P output and true amino acid digestibility by using a Regression Analysis Technique. Fan et al. (2001) and Shen et al. (2002) used this method to determine endogenous P output and true P digestibility. Zhang (2004) used this method to determine true P and Ca digestibility.

These experiments proved the feasibility of this method which is cheaper, safer and more convenient.

Endogenous P determined in the present study was 1.14 g/kg DMI which is similar to the experimental result (1.08 g/kg DMI) of Zhang (2004) and higher than that of Shen (2002). The endogenous percentage in total P and TDP requirement was 17% and 30.54%, respectively, in this experiment, compared to were 18% and 39.13% recorded by Zhang (2004). Fan et al. (2001) determined that the endogenous P output of pigs (body weights were 5 to 20 kg) was 0.31 g/kg DMI, which was 5.8% to 12.8% of total P requirement and 9.5% to 24.1% of TDP requirement. Later, Shen et al. (2002) determined the value in growing pigs (BW 20-45 kg) was 0.67 g/kg DMI which was 12.3% of total P requirement and 26.6% of TDP requirement. There are gender differences and at different periods even in the same gender and period, because endogenous P output can be influenced by factors such as the ratio of Ca to P, P levels and feed management.

According to the present results, it can be deduced that true P digestibility reaches the highest point when dietary TDP concentration is 0.34%. When dietary TDP concentration exceeds the requirement of pigs, the excess would be excreted from the body so that the percentage of absorbed P in total dietary P input is reduced. Thus, true P digestibility decreased when dietary TDP concentration exceeded the pig's need. Except for diet 1 the experimental diets contained dicalcium phosphate. The biological availability of P in dicalcium phosphate is higher than in plant ingredients so the higher true P digestibility of diets 2, 3, 4 and 5 compared to diet 1 was attributable to dicalcium phosphate. True P digestibility increased progressively to the highest point, and then decreased when dietary true digestible TDP concentration increased from 0.26% to 0.39%. However, the range of dietary TDP concentration between diets 4 and 5 was too wide to determine accurately the change of true P digestibility and the deduced result needs confirmation.

The effect of dietary TDP concentration on P output and P balance

Fecal and urinary P output increased with increasing dietary P input which indicated that there was a positive correlation between fecal and urinary P output and dietary P input. The result was in accord with the results of Vipperman et al. (1974) and Miller et al. (1964). Vipperman et al. (1974) found urinary P output increased with increased dietary P input when dietary Ca concentration was higher (7.5%) and increased linearly with increasing dietary P input when dietary Ca concentration was lower (2.5%). They also found that urine P output decreased with increasing dietary Ca input when dietary P level was higher

and urinary P output was nearly zero when dietary Ca level exceeded dietary P level. Miller et al. (1962) found that the retention of P increased with an increasing dietary P level and the amount of retentive P was influenced primarily by urinary P output.

In this study, when expressed as g/kg DM feces, P output was lowest when dietary TDP concentration was 0.26% which indicated that diet 5 may be better than the other experimental diets in terms of P availability and P output. When analysed further, there was a lower point of P output between dietary TDP concentration of 0.26% and 0.39%. The lower point was apparent when dietary TDP concentration was approximately 0.34% according to the results. When expressed as g/kg DMI, fecal P output increased with increasing dietary TDP concentration, but the ratio of fecal P output to dietary P input decreased from 74.4% to 65.6%, then increased to 68.7%. When the ratio of fecal P output to dietary P input was analysed, the lower point was evident when dietary TDP concentration was near to 0.34% according to the results. The above two conclusions were in accord with the result of another part of the study, which proved that the performance of growing pigs was best when dietary TDP concentration was 0.34%. It is concluded that the optimum level of addition of TDP was 0.34%.

Ca to P ratio

A major factor that affects the determination of P requirements by animals is the Ca to P ratio (Combs et al., 1991). This ratio greatly influences the availability of dietary Ca and P that can enter the portal circulation (Anderson, 1991). NRC (1998) recommended a ratio of total Ca to total P in a typical corn- and soybean meal-based diet of between 1:1 and 1:1.2 and a ratio of total Ca to available P of between 2:1 and 3:1. ARC (1981) suggested that the ratio of total Ca to total P should not exceed 2:1 for growing pigs and that a ratio of total Ca to total P ratio between 1:1 and 1.2:1 for diets containing phytic acid is beneficial for growth performance and bone function in pigs. The phytate and phosphatase concentrations differ among dietary ingredients, and thus there were marked disparities in P availability in the swine alimentary tract. Therefore, formulating diets on the basis of a total Ca to total P ratio cannot accurately reflect the actual requirements for these two minerals by pigs. To meet the metabolic requirements of both Ca and P by swine precisely, using a ratio of total Ca to TDP is a better choice. The results from our extensive research have established that a total Ca to TDP ratio of 2:1 is optimal for growth performance and the efficiency of utilization of dietary P in growing pigs (Yin, 2005).

Mudd et al. (1969) found retentive Ca and P increased with increasing dietary Ca and P level while the retention

decreased gradually, which was proved by this study, but the retention was lower when dietary TDP concentration was lower (0.16%). The retentive Ca in this study differed to that of Mudd et al. (1969), and the reason may be that the ratio of Ca to P were not equal in the two experiments. Vipperman et al. (1974) found that the retentive P decreased when dietary Ca level was added without additional dietary P level, and retentive P increased with increasing dietary Ca level when dietary P level was higher. The results of the present study indicated that serum alkaline phosphatase activity was not affected by dietary Ca content when the ratio of total Ca to TDP was kept at 2:1. Also, varying dietary Ca intakes at a constant ratio (2:1) of total Ca to TDP had no effect on total Ca concentration in the serum of growing pigs (Table 7). Similar findings were reported by Lin et al. (2002) and Wang et al. (2002). However, total P concentration in serum increased with increasing dietary TDP levels (Table 7). These results are comparable to those reported by Lin et al. (2002) for weanling pigs as well as growing-finishing pigs. Taken together, the findings from the present study and the work of Lin et al. (2002) demonstrate that total P concentrations in serum rise as dietary P levels increase to an optimal level. Moreover, the research shows that an optimal TDP level in the diet is 0.34% (on an as-fed basis) for 20- to 40-kg pigs. Thus, a further increase in dietary TDP levels beyond 0.34% did not increase total P concentration in serum (Table 7). Interestingly, Wang et al. (2002) reported that serum P concentration increased progressively with increasing dietary P level in Chinese Xiangzhu pigs. It is possible that there are significant differences in Ca or P digestion and metabolism between different breeds. Further study is needed to test this important hypothesis.

Besides the data on serum P concentrations, growing pigs exhibited the highest ADG and highest feed efficiency when fed a diet containing 0.34% TDP and 0.68% Ca (Table 6). This diet provided the 20- to 40-kg pigs with daily TDP, available P and total P intakes of 3.76, 3.65, and 6.81 g/d, respectively (Table 6). Note that the recommended requirements (NRC, 1998) of the available P in the diet was 0.23% with daily intakes of available P and total P intake being 4.27 and 9.28 g/d respectively, for pigs with BW of 20-50 kg. Thus, the NRC-recommended dietary intake of total P by growing pigs (NRC, 1998) is 36% more than the optimal value (6.81 g/d) obtained in the present study. Clearly, NRC (1998) overestimated substantial P requirement by growing pigs fed corn- and soybean meal-based diets.

The effect of dietary TDP concentration on apparent N digestibility and N balance

The influence of dietary TDP concentration on apparent

N digestibility might be attributable to saponin formed by the connecting of P with Ca and fat which restrained the digestibility of amino acids. Vipperman et al. (1974) found that retentive N decreased when dietary Ca level was added without increasing dietary P level. Mudd et al. (1969) found that the effect of dietary P level on the amount of retentive N was greater than the effect of dietary Ca level. In this study, dietary CP level was 17.34% and the average total fecal N output was 12.14 g/d which was lower than the value of 16.6 g/d obtained by Cromwell (1996) when dietary CP level was 16%. The retention of N was over 66% in this study which is higher than the result of Fan et al. (1999). The reason for N output being higher in the latter experiment may be that the phytate phosphorus level was higher and available phosphorus was lower than in this experiment and, as a result, more saponin was formed.

Fecal and urinary N output decreased all along in Experiment 1 and may have been related to the decrease of dietary N input. The results indicated that dietary TDP concentration could influence the apparent N digestibility, but not significantly. These studies also validated the experimental result of Cromwell (1996) which concluded that there was no interaction between dietary P and CP levels.

CONCLUSIONS AND IMPLACATIONS

The results showed that true P digestibility and total P output increased with increasing dietary TDP concentration. When dietary TDP concentration was 0.34%, the ratio of total P output to dietary P input was the lowest, fecal P output was not the lowest but the most appropriate. Apparent N digestibilities were similar at different levels of dietary TDP, which indicated that dietary TDP concentration may not influence dietary apparent N digestibility and N output. N output was influenced mainly by dietary CP level.

In summary, results of the present studies indicate that the appropriate dietary TDP level is 0.34% of the diet (on an as-fed basis) at a total Ca to TDP ratio of 2:1. This corresponds to 5.1 g TDP/d for a 30-kg pig that consumes 1.5 kg feed daily. The findings suggest that dietary P requirement by growing pigs is substantially overestimated in the current version (10th edition) of NRC-recommended nutrient requirements of swine.

ACKNOWLEDGEMENTS

This research was jointly supported by grants from National Basic Research Program of China (contract No. 2004CB117502), the National Natural Science Foundation of China (contract No. 30528006, 30671517, 30700581,

30771558 and 30371038), National Scientific and Technological Supporting Project (2006BAD12B07) The Chinese Academy of Sciences and Knowledge Innovation Project (contract No. KZCX3-SW-441, YW-N-022, and KSCX2-SW323), Program for Hubei Cu Tiang Scholars. Fund of Agricultural Science and Technology outcome application (contract No. 2006GB24910468). National Scientific and Technological Supporting Project (2006BAD12B07 and 2006BAD12B02-5-2) and Guang Dong Province Project (contract No.2006B200330005 and Program for Changjiang Scholars and Innovative University Research Team (contract No. 65292 and IRT0540).

REFERENCES

- Abelson, P. H. 1999. A potential phosphate crisis. Sci. Washington, DC. 283:2015.
- Agricultural Research Council. 1981. The Nutrient Requirements of Pigs. Slough, England: Commonwealth Agricultural Bureaux.
- Anderson, J. B. 1991. Nutritional biochemistry of calcium and phosphorus. J. Nutr. Biochem. 6:200-307.
- Bhanja, S. K., V. R. Reddy, A. K. Panda, S. V. Rama Rao and R. P. Sharma. 2005. Effect of supplementing microbial phytase on performance of broiler breeders fed low non-phytate phosphorus diet. Asian-Aust. J. Anim. Sci. 18:1299-1304.
- Byrkit, D. R. 1987. Statistics Today-A Comprehensive Introduction. Benjamin/Cummings Publishing Company, Menlo Park, CA.
- Chen, Lixiang, Yu-Long Yin, Wenjuan S. Jobgen, Scott C. Jobgen, Darrell A. Knabe, Wei-Xin Hu and Guoyao Wu. 2007. *In vitro* oxidation of essential amino acids by jejunal mucosal cells of growing pigs. Livest. Sci. 109: 19-23.
- Deng, Ze-yuan, Jin-wu Zhang, Jing Li, Ya-wei Fan, Shu-wen Cao, Rui-lin Huang, Yu-long Yin, Hua-yi Zhong and Tie-jun Li. 2007a. Effect of polysaccharides of cassiae seeds on the intestinal microflora of piglets. Asia Pac. J. Clin. Nutr. 16(Suppl 1):143-147.
- Deng, Ze-Yuan, Jin-Wu Zhang, Guo-Yao Wu, Yulong Yin, Zheng Ruan, Tie-Jun Li, Wu-Ying Chu, Xiang-Feng Kong, You-Ming Zhang, Ya-Wei Fan, Rong Liu and Rui-Lin Huang. 2007b. Dietary supplementation with polysaccharides from Semen cassiae enhances immunoglobulin production and interleukin gene expression in early-weaned piglets. J. Sci. Food Agric. 87:1868-1873.
- Combs, N. R., E. T. Kornegay, M. D. Lindemann, D. R. Notter, J. H. Wilson and J. P. Mason. 1991. Calcium and phosphorus requirement of swine from weaning to market weight: 2. Development of response curves for bone criteria and comparison of bending and shear bone testing. J. Anim. Sci. 69: 682-693.
- Cromwell, G. L. 1996. Reduce the excretion of phosphorus and nitrogen by regulating diets. Proceedings of the 14th IPV S Congress, Bologna, Italy, 7(10):418.
- Cromwell, G. L., V. W. Hays, C. W. Scheer and J. R. Overfield. 1972. Effects of dietary phosphorus and calcium level on performance and carcass, metacarpal and turbinat characteristics of swine. J. Anim. Sci. 34:746-750.
- Deng, D., L. Huang, T. J. Li, G. Y. Wu, M. Y. Xie, Z. R. Tang, P. Kang, Y. M. Zhang, M. Z. Fan, X. F. Kong, Z. Ruan, H. Xiong, Z. Y. Deng and Y.-L. Yin. 2007a. Nitrogen balance in barrows fed low-protein diets supplemented with essential amino acids. Livest. Sci. 109:220-223.
- Deng, D., Ai-Ke Li, W. Y. Chu, R. L. Huang, T. J. Li, X. F. Kong, Z. J. Liu, G. Y. Wu, Y. M. Zhang and Y. L. Yin. 2007b. Growth performance and metabolic responses in barrows fed low-protein diets supplemented with essential amino acids. Livest. Sci. 109:224-227.
- Fan, M. Z. and W. C. Sauer. 1999. Variability of apparent ileal amino acid digestibility in peas for pigs with the direct, difference, and regression methods. Can. J. Anim. Sci. 79:467-475.
- Fan, M. Z., T. Archbold, W. C. Sauer and D. Lackeyram. 2001. Novel methodology allows simultaneous measurement of true phosphorus digestibility and the gastrointestinal endogenous phosphorus outputs in studies with pigs. J. Nutr. 131:2388-2396.
- Fan, M. Z., T. J. Li, Y. L. Yin, R. J. Fang, Z. Y. Tang, Z. P. Hou, R. L. Huang, Z. Y. Deng, H. Y. Zhong, R. G. Zhang, J. Zhang, B. Wang and H. Schulze. 2005. Effect of phytase supplementation with two levels of phosphorus diets on ileal and faecal digestibilities of nutrients and phosphorus, calcium, nitrogen and energy balances in growing pigs. Anim. Sci. 81:67-75.
- Fan, M. Z. and W.C. Sauer. 2002. Additivity of apparent ileal and fecal phosphorus digestibility values measured in single feed ingredients for growing-finishing pigs. Can. J. Anim. Sci. 82:183-191.
- Fang, R. J., Y. L. Yin, K. N. Wang, J. H. He, Q. H. Chen, T. J. Li, M. Z. Fan and G. Wu. 2007a. Comparison of the regression analysis technique and the substitution method for the determination of true phosphorus digestibility and faecal endogenous phosphorus losses associated with feed ingredients for growing pigs. Livest. Sci. 109:251-254.
- Fang, R.-J., T.-J. Li, K.-N. Wang, G.-Y. Wu, D. Qi, J. H. He, Y.-L. Yin and M.- Z. Fan. 2007b. Additives of apparent and true phosphorus digestibilities in feed ingredients for growing pigs. Asian-Aust. J. Anim. Sci. 20:1092-1099.
- Fig P Software Corporation. 1992. Fig.P, version 6.0 Biosoft. Cambridge, UK.
- Heinoen, J. K. and R. J. Lahti. 1981. A new and convenient colorimetric determination of inorganic orthophosphate and its application to the assay of inorganic pyrophosphate. Anal. Biochem. 133:313-317.
- Huang, R. -L., Y. -L. Yin, K.-P. Wang, T.-J. Li and J.-X. Liu. 2003. Nutritional value of fermented and not fermented material of distiller's grains in pig nutrition. J. Anim. Feed Sci. 12:261-269.
- Huang, R.-L., Y.-L. Yin, G.-Y. Wu, Y.-G. Zhang, T.-J. Li, L.-L. Li, M.-X. Li, Z.-R. Tang, J. Zhang, B. Wang, J.-H. He, and X.-Z. Nie. 2005. Effect of dietary oligochitosan supplementation on ileal digestibility of nutrients and performance in broilers. Poult. Sci. 84:1383-1388.
- Huang, R. L., Y. L. Yin, M. X. Li, G. Y. Wu, T. J. Li, L. L. Li, C. B. Yang, J. Zhang, B. Wang, Z. Y. Deng, Y. G. Zhang, Z. R. Tang, P. Kang and Y. M. Guo. 2007. Dietary oligochitosan supplementation enhances immune status of broilers. J. Sci.

- Food Agric. 87:153-159.
- Jongbloed, A. W. and N. P. Lenis. 1998. Environmental concerns of animal manure. *J. Anim. Sci.* 76:2641-2648.
- Kong, X.-F., G.-Y. Wu, Y.-L. Yin, H.-J. Liu, F.-G. Yin, T.-J. Li, R.-L. Huang, P. Kang, F.-F. Xing, M.-Z. Fan, C.-B. Yang and Q.-H. He. 2007a. Dietary supplementation with Chinese herbal ultra-fine 3 powder enhances cellular and humoral immunity in early weaned piglets. *Livest. Sci.* 108:94-98.
- Kong, X. F., G. Y. Wu, Y. P. Liao, Z. P. Hou, H. J. Liu, F. G. Yin, T. J. Li, R. L. Huang, Y. M. Zhang, D. Deng, P. Kang, R. X. Wang, Z. Y. Tang, C. B. Yang, Z. Y. Deng, H. Xiong, W.-Y. Chu, Z. Yuan, M. Y. Xie and Y. L. Yin. 2007b. Effects of Chinese herbal ultra-fine powder as a dietary additive on growth performance, serum metabolites and intestinal health in early-weaned piglets. *Livest. Sci.* 108:272-275.
- Kong, Xiangfeng, Yulong Yin, Guoyao Wu, Hejun Liu, Fugui Yin, Tiejun Li, Ruilin Huang, Zheng Ruan, Hua Xiong, Zeyuan Deng, Mingyong Xie, Yiping Liao and Sungwoo Kim. 2007c. Dietary supplementation with *acanthopanax senticosus* extract. *Asian-Aust. J. Anim. Sci.* 20:1453-1461.
- Modulates Cellular and Humoral Immunity in Weaned Piglets
- Lin, Y. C. and Z. Y. Jiang. 2002. Study on dietary available phosphorus requirement of growing-finishing pigs. *Swine Production*, 4:1-7 (in Chinese).
- Li, T.-J., Y.-L. Yin, S.-Y. Bin, R.-L. Huang, Q.-Z. Dai, P. Zhang, Z.-P. Hou and H. Yang. 2007a. Growth performance and nitrogen metabolism in weaned pigs fed diets containing different sources of starch. *Livest. Sci.* 109:73-76.
- Li, L.-L., Y. -L. Yin, Y.-H. Liu, D.-X. Hou, Z.-P. Hou, C.-B. Yang and X.-J. Yang. 2007b. Effects of metallothionein on antioxidative function and related enzyme activities in blood and liver, and pork meat quality. *Asian-Aust. J. Anim. Sci.* 20:761-767.
- Mallin, M. A. 2000. Impacts of industrial animal production on river and estuaries. *American scientist* Jan-Feb, 26-37.
- Miller, E. R., D. E. Ullery, C. L. Zutaut, B. V. Baltzer, D. A. Schmidt, J. A. Hofer and R. W. Lueche. 1962. Calcium requirement of baby pig. *J. Nutr.* 77:7.
- Miller, E. R., D. E. Ullery, C. L. Zutaut, J. A. Hofer and R. W. Luecke. 1964. Mineral balance studies with the baby pig: effects of dietary phosphorus level upon calcium and phosphorus balance. *J. Nutr.* 82:111.
- Mudd, A. J., W. C. Smith and D. G. Armstrong. 1969. The influence of dietary concentration of calcium and phosphorus on their retention in the body of the growing pig. *J. Agric. Sci.* 73:189-196.
- National Research Council. 1998. *Nutrient Requirements of Swine*, 10th ed. National Academy Press, Washington, DC., USA.
- Pouslen, H. 2000. Phosphorus utilization and excretion in pig production. *J. Environ. Qualif.* 29:24-27.
- Ruan, Z., Y.-G. Zhang, Y.-L. Yin, T.-J. Li, R.-L. Huang, S. W. Kim, G. Y. Wu and Z. Y. Deng. 2007. Dietary requirement of true digestible phosphorus and total calcium for growing pigs. *Asian-Aust. J. Anim. Sci.* 20:1236-1242.
- Sacakli, P., A. Sehu, A. Ergün, B. Genç and Z. Selçuk. 2006. The effect of phytase and organic acid on growth performance, carcass yield and tibia ash in quails fed diets with low levels of non-phytate phosphorus. *Asian-Aust. J. Anim. Sci.* 19:198-202.
- SAS Institute Inc. 2000. *The SAS System*. SAS Institute, Cary, NC, USA.
- Saha, D. C. and R. L. Gilbreath. 1991. Analytical recovery of chromium from diet and feces determined by colorimetry and atomic absorption spectrophotometry. *J. Sci. Food Agric.* 55:433-446.
- Shen, Y. R., Z. F. Ming, A. Ajakaiye and T. Archbold. 2002. Use of the regression analysis technique to determine the true phosphorus digestibility and the endogenous phosphorus output associated with cron in growing pigs. *J. Nutr.* 132:1199-1206.
- Shen, Yingran, Yulong Yin, Eduardo R. Chavez and Ming Z. Fan. 2005. Methodological aspects of measuring phytase activity and phytate phosphorus content in selected cereal grains and intestinal samples of pigs. *J. Agric. Food Chem.* 53:853-859.
- Steel, R. G. D. and J. H. Torrie. 1980. *Principles and Procedures of Statistics: A Biometrical Approach*, 2nd ed. McGraw-Hill, New York, NY.
- Tang, Z. R., Y. L. Yin, C. M. Nyachoti, R. L. Huang, T. J. Li, C. B. Yang, X. J. Yang, J. S. Gong, J. Peng, D. S. Qi, J. J. Xing, Z. H. Sun and M. Z. Fan. 2005. Effect of dietary supplementation of chitosan and galacto-mannan- oligosaccharide on serum parameters and the insulin-like growth factor-I mRNA expression in early-weaned piglets. *Dom. Anim. Endocrin.* 28:430-441.
- Vipperman, P. E., E. R. Peo and P. J. Cunningham. 1974. Effect of dietary calcium and phosphorus level upon calcium, phosphorus and nitrogen balance in swine. *J. Anim. Sci.* 38:758-765.
- Wang, F. L., M. F. Zhang, Q. M. Chen and M. Q. Xu. 2002. The effects of dietary phosphorus and calcium to phosphorus ratio on the activity of alkaline phosphatase and serum calcium and phosphorus in miniature-pigs (Xiang pig). *Acta Zoonutrimenta Sinica.* 13:36-42.
- Weremko, D., H. Fandrejowski, T. Zebrowska, I. K. Han, J. H. Kim and W. T. Cho. 1997. Bioavailability of phosphorus in feeds of plant origin for pigs. *Asian-Aust. J. Anim. Sci.* 10:551-566.
- Wu, G. 1998. Intestinal mucosal amino acid catabolism. *J. Nutr.* 128:1249-1252.
- Yang, H., A. K. Li, Y. L. Yin, T. J. Li, Z. R. Wang, G. Wu, R. L. Huang, X. F. Kong, C. B. Yang, P. Kang, J. Deng, S. X. Wang, B. E. Tan, Q. Hu, F. F. Xing, X. Wu, Q. H. He, K. Yao, Z. J. Liu, Z. R. Tang, F. G. Yin, Z. Y. Deng, M. Y. Xie and M. Z. Fan. 2007. True phosphorus digestibility and the endogenous phosphorus outputs associated with brown rice for weanling pigs measured by the simple linear regression analysis technique weanling pigs. *Anim.* 1:213-220.
- Yang, Cheng-bo, Ai-ke Li, Yu-long Yin, Rui-Lin Huang, Tie-jun Li, Li-li Li, Yi-ping Liao, Ze-yuan Deng, Hua-yi Zhong, Xiao-jian Yang and Ming-Z. Fan. 2005. Effects of dietary supplementation of cysteamine on growth performance, carcass quality, serum hormones and gastric ulcer in finishing pigs. *J. Sci. Food and Agric.* 85:1947-1952.
- Yin, Y. -L., Huang, R.-L., Zhong, H.-Y., Chen, C.-M., and Dai, H. 1991a. Influence of different cannulation techniques on the pre-cecal digestibility of protein, amino acids and cell wall constituents from diets, containing different protein meal, in pigs. *Animal Feed Science and Technology* 35:271-281.

- Yin, Y. -L., H.-Y. Zhong, R.-L. Huang and C.-M. Chen. 1991b. Digestibility of energy, crude protein and cell wall constituents in different parts of the intestinal tract in growing pigs. *Agribiol. Res.* 44:14-22.
- Yin, Y.-L., R.-L. Huang and H.-Y. Zhong. 1993a. Comparison of the ileorectal anastomosis and conventional method for the measurement of ileal digestibility of protein sources and mixed diets in growing pigs. *Anim. Feed Sci. Technol.* 42:297-308.
- Yin, Y. -L., H.-Y. Zhong, R.-L. Huang, C.-M. Chen, T.-J. Li and Y.-F. Pai. 1993b. Nutritive value of feedstuffs and diets for pigs. I. Chemical composition, apparent ileal and fecal digestibility. *Anim. Feed Sci. Technol.* 44:1-27.
- Yin, Y. -L., R.-L. Huang and H.- Y. Zhong. 1993c. Effect of autoclaving on ileal digestibility of crude protein, amino acid, urease activity of beans. *Zeitschrift für Tierphysiologie, Tierernährung und Futtermittelkunde* 71:65-74.
- Yin, Y. -L. 1994. Nutritive value of feedstuffs and diets for pigs, II. Apparent post-ileal digestibility and interrelationship between dietary constituents and fecal and ileal digestibility. *Anim. Feed Sci. Technol.* 45:243-255.
- Yin, Y. -L., C.-M. Chen, H.-Y. Zhong and R.-L. Huang. 1994. Digestibility of energy, cell wall constituents, crude protein and amino acids of the Chinese oil seed meals for pigs. *Anim. Feed Sci. Technol.* 45:283-298.
- Yin, Y. -L., H.-Y. Zhong and R.-L. Huang. 1995. Determination of the apparent ileal digestibility of protein and amino acid in feedstuffs and mixed diets for growing- finishing pigs with the mobile nylon bag technique. *Asian-Aust. J. Anim. Sci.* 8:433-441.
- Yin, Y.-L. and K. J. McCracken. 1996. Methodological aspects of the *in vivo* measurement of ileal amino acid digestibility in pigs- a review. *Asian-Aust. J. Anim. Sci.* 9:495-502.
- Yin, Y.-L., J. McEvoy, W.-B. Souffrant, H. Schulze and K. J. McCracken. 2000a. Apparent digestibility (ileal and overall) of nutrients as evaluated with PVTC-cannulated or ileo-rectal anastomised pigs fed diets containing two indigestible markers. *Livest. Prod. Sci.* 62:133-141.
- Yin, Y.-L., J. McEvoy, W.-B. Souffrant, H. Schulze and K. J. McCracken. 2000b. Apparent digestibility (ileal and overall) of nutrients and endogenous nitrogen losses in growing pigs fed wheat or wheat by-products without or with xylanase supplementation. *Livest. Prod. Sci.* 62:119-132.
- Yin, Y.-L., J. McEvoy, H. Schulze and K. J. McCracken. 2000c. Studies on cannulation method and alternative indigestible markers and the effects of food enzyme supplementation in barley-based diets on ileal and overall apparent digestibility in growing pigs. *Anim. Sci.* 70:63-72.
- Yin, Y. -L., S. K. Baidoo, J. J. J. Boychul and H. Simmins. 2001a. Performance and carcass characteristics of growing pigs and broiler fed diets containing micronized barley, ground barley, wheat and maize. *J. Sci. Food Agric.* 81:1487-1497.
- Yin, Y.-L., J. McEvoy, H. Schulze and K. J. McCracken. 2001b. Effects of xylanase and antibiotic addition on ileal and overall apparent digestibility and evaluating HCL-insoluble ash as an indigestible marker in growing pigs. *Anim. Sci.* 72:95-103.
- Yin, Y.-L., S. K. Baidoo, H. Schulze and P. H. Simmins. 2001C. Effect of supplementing diets containing hullless barley varieties having different levels of non-starch polysaccharides with -glucanase and xylanase on the physiological status of gastrointestinal tract and nutrient digestibility of weaned pigs. *Livest. Prod. Sci.* 71:97-107.
- Yin, Y.-L., S. K. Baidoo, Z. L. Jin, Y. G. Liu, H. Schulze and P. H. Simmins. 2001d. The effect of different carbohydrase and protease supplementation on apparent (ileal and overall) digestibility of nutrient of five hullless barley varieties in young pigs. *Livest. Prod. Sci.* 71:109-120.
- Yin, Y.-L., R.-L. Huang, H.-Y. Zhong, T. J. Li, W. B. Souffrant and C. F. M. de Lange. 2002a. Evaluation of mobile bag technique for determining apparent ileal digestibilities of crude protein and amino acids in growing pigs. *J. Anim. Sci.* 80:409-420.
- Yin, Y.-L., Nar K. Gurung, E. A. Jeaurond, P. H. Sharpe and C. F. M. de Lange. 2002b. Nutrient digestibility of Canadian-developed sorghum and pearl millet grains fed to growing pigs compared to traditional cereal grains. *Canadian J. Anim. Sci.* 82:385-391.
- Yin, Y.-L., Z.-Y. Deng, H.-L. Huang, T.-J. Li and H. Y. Zhong. 2004. The effect of arabinoxylanase and protease supplementation on nutritional value of diets containing wheat bran or rice bran in growing pigs. *J. Anim. Feed Sci.* 13:445-461.
- Zhang, Y. L. 2004. Determination of true calcium and phosphorus digestibility values and the endogenous calcium and phosphorus losses associated with soybean meal for gilt and barrow by linear regression analysis technique. Master thesis, Neimenggu Agricultural University, China.