



## Investigation into the mechanism of action of *Moringa oleifera* for its anti-asthmatic activity

Anita Mehta\* and Babita Agrawal

Department of Pharmacology, L.M. College of Pharmacy, Ahmedabad - 3800009, India

### SUMMARY

In the present investigation, we studied the effect of alcoholic extract of *Moringa oleifera* (*M. oleifera*) seed kernels on various experimental models of bronchial asthma. Significant ( $P < 0.05$ ) increase in preconvulsion time was observed due to pretreatment with *M. oleifera* when the guinea pigs were exposed to either acetylcholine (Ach) or histamine aerosol. This bronchodilating effect of *M. oleifera* was comparable to ketotifen fumarate. Spasmolytic effect of *M. oleifera* was also observed by dose dependent inhibition of ideal contractions induced by Ach, 5HT, histamine and  $BaCl_2$ . Alcoholic extract of *M. oleifera* produced significant dose dependent protection by egg albumin and compound 48/80 induced mast cell degranulation. Pretreatment with alcoholic extract of *M. oleifera* also decreased carrageenan induced rat paw edema, which was comparable to that of standard diclofenac sodium. Minimum inhibitory concentration for alcoholic extract of *M. oleifera* was low as compared to cold-water extract and hot water extract when antimicrobial activity was tested against various respiratory pathogens like *Escherichia coli* (*E. coli*), *Staphylococcus aureus* (*S. aureus*) and *Pseudomonas aeruginosa* (*P. aeruginosa*). Our data suggest that antiasthmatic activity of *M. oleifera* seed kernels may be due to its bronchodilator, anti-inflammatory, mast cell stabilization and antimicrobial activity.

**Key words:** Bronchial asthma; *Moringa oleifera*; Bronchodilator; Anti-inflammatory; Mast cell stabilization

### INTRODUCTION

Bronchial asthma is a chronic respiratory disorder affecting a large proportion of population throughout the world. The currently used drugs for the treatment of this disease in modern medicine are far from satisfactory as they provide only symptomatic relief, produce several adverse effects and may lose effectiveness on continued use. Muscle tremor and hypokalemia are major adverse effects of  $\beta_2$  agonists (Haalboom *et al.*, 1985; Nelson, 1986). Theophylline has narrow therapeutic

index and requires monitoring of drug levels (Nasser and Rees, 1993; Stoloff, 1994). Adverse effects of corticosteroids include fluid retention, increased cell mass, increased appetite, weight gain, osteoporosis, capillary fragility, hypertension, peptic ulceration, diabetes, cataract, and psychosis (Dajani *et al.*, 1981). Hence Ayurveda has recommended number of drugs from indigenous plants sources for the treatment of bronchial asthma and other allergic disorders and have been successful in controlling the disease as well. Large numbers of medicinal plant preparations have been reported to possess anti-asthmatic effects.

*Moringa oleifera* (*M. oleifera*) is a small or medium sized tree, cultivated throughout India. The tender

\*Correspondence: Anita Mehta, Department of Pharmacology, L.M. College of Pharmacy, Ahmedabad - 3800009, India. E-mail: dranitalmcp@rediffmail.com

Pods are esteemed as a vegetable. Seeds are used as purgative, antipyretic and anti-inflammatory (Warrier *et al.*, 1997). The plant possesses antimicrobial activity (Caceres *et al.*, 1991), while the seeds reportedly have antispasmodic, anti-inflammatory and diuretic activity (Caceres *et al.*, 1992). Its leaves and fruits are edible, rich in ascorbic acid. The plant is also reported to elicit good clinical response in children suffering from upper respiratory tract infection and skin infection. It has been reported that alkaloid from the plant closely resembles ephedrine in action and useful in treatment of asthma. Alkaloid *Moringine* relaxes bronchioles (Kirtikar and Basu, 1975). In the present study, we have investigated the anti-asthmatic potential of *M. oleifera* on various experimental models like Bronchodilating, mast cell stabilizing, anti-inflammatory and antimicrobial activity.

## MATERIALS AND METHODS

### Plant material

Seed kernels of *M. oleifera* were purchased from the local market of Ahmedabad and were identified and authenticated by Dept. of pharmacognosy, L.M. College of Pharmacy, Ahmedabad, India. A voucher specimen was deposited at the Dept. of pharmacognosy, Ahmedabad. The coarse powder (500 g) of the dried seed kernels was defatted using petrol ether and then it was exhaustively extracted using 95% ethanol (2,000 ml) in a Soxhlet extractor. Cold aqueous extract of *M. oleifera* was prepared by extracting 1 part of seed kernels with 10 parts of water for 2 h without heating, while hot aqueous extract was prepared by heating the seed kernels with water. The extracts were concentrated under reduced pressure to yield a syrupy mass and stored in air tight container in cool place and used throughout the project.

### Animals

All animals were housed at ambient temperature ( $22 \pm 1^\circ\text{C}$ ), relative humidity ( $55 \pm 5\%$ ) and 12/12 h

light/dark cycle. Animals had access to standard pellet diet and water given ad libitum. The protocol of the experiment was approved by the institutional animal ethical committee as per the guidance of the Committee for the purpose of Control and Supervision of Experiments on Animals (CPCSEA), Ministry of Social Justice and Empowerment, Government of India.

### Studies on Acetylcholine and Histamine induced bronchospasm in guinea pigs

Guinea pigs of either sex weighing 350 - 500 g were selected and randomly divided into six groups each containing six animals. The drugs were administered orally in 0.5% sodium carboxymethyl cellulose (CMC). The single dose treatments were given one and half an hour before the study. The following schedule of treatment was administered:

Group I: 0.5% CMC (control)

Group II: Ketotifen (1 mg/kg) (standard)

Group III: Alcoholic extract of *M. oleifera* (100 mg/kg)

Group IV: Alcoholic extract of *M. oleifera* (200 mg/kg)

Later the animals were exposed to an aerosol of 0.25% histamine and time for preconvulsion state was noted for each animal (Sheth *et al.*, 1972). After about 15 days of wash out period, the same animals were given the above treatments and time for preconvulsion state was noted for 0.5% acetylcholine bromide aerosol spray.

### Studies on isolated Guinea pig ileum

Overnight fasted guinea pigs of either sex weighing 400 - 600 g were sacrificed using cervical dislocation method. Ileum was quickly dissected out and mounted in an organ bath maintained at  $37 \pm 1^\circ\text{C}$  and containing 20 ml Tyrode's solution under basal tension of 500 mg. The composition of solution in mM was NaCl, 137; CaCl<sub>2</sub>, 1.8; KCl, 2.7; glucose, 5.55; NaHCO<sub>3</sub>, 11.9; MgCl<sub>2</sub>, 1; NaH<sub>2</sub>PO<sub>4</sub>, 0.4. The solution was continuously bubbled with air. The responses to drug were recorded on a Student

physiograph (BioDevices) using isotonic transducer, which exerted a basal tension equivalent to 500 mg load on tissue. The tissue was allowed to equilibrate for 30 min, during which, the bathing solution was changed at every 10 min. The contractile responses of ileum to various agonists (Acetylcholine, histamine, 5-HT and BaCl<sub>2</sub>) were recorded in presence and absence of alcoholic extract of *M. oleifera*.

#### **Studies on compound 48/80 and egg albumin induced rat peritoneal mast cell degranulation**

Normal saline containing 5 units/ml of heparin was injected in the peritoneal cavity of male rats lightly anaesthetized with ether. After a gentle abdominal massage, the peritoneal fluid containing mast cells was collected in centrifuge tubes placed over ice. Peritoneal fluid of 4 - 5 rats was collected and pooled and centrifuged at 2,000 rpm for 5 min. Supernatant solution was discarded and the cells were washed twice with saline and resuspended in 1 ml of saline.

0.1 ml of the peritoneal cell suspension was transferred to 6 test tubes and was treated as follows.

Test tube no. 1 & 2 - Saline

Test tube no. 3 - 0.1 ml of 0.5 mg/ml alcoholic extract of *M. oleifera* in Saline

Test tube no. 4 - 0.1ml of 1.0 mg/ml alcoholic extract of *M. oleifera* in Saline

Test tube no. 5 - 0.1 ml of 2.0 mg/ml alcoholic extract of *M. oleifera* in Saline

Test tube no. 6 - 0.1 ml of 10 µg/ml of Ketotifen fumarate

Each test tube was incubated for 15 min at 37°C and then Compound 48/80 (0.1 ml, 10 µg/ml) was added to each test tube except test tube no. 1. After further incubation for 10 min. at 37°C, the cells were stained with 0.1% toluidine blue solution made in distilled water and examined under the high power of light microscope. Percent protection of the mast cells in the control group and the treated groups were calculated by counting the

number of degranulated mast cells from total of atleast 100 mast cells counted.

In another study, rats were sensitized by administering three doses of 350 µg of egg albumin adsorbed on 60 mg of aluminum hydroxide gel, the doses being given on the first, third and fifth day subcutaneously. The mast cells were collected on the tenth day of sensitization. The study was conducted in the same manner as above and the sensitized cells were degranulated using egg albumin (1 mg/ml). Percent protection of the mast cells in the control group and the treated groups were calculated by counting the number of degranulated mast cells from total of atleast 100 mast cells counted. Control group is consisted of positive control group in which egg albumin was added without addition of test agent and a negative control group in which neither egg albumin nor the test agent was added to correct for spontaneous degranulation of mast cells without any degranulating agent.

#### **Anti inflammatory study**

##### **Carrageenan induced rat paw edema**

Albino rats of either sex weighing 200 - 250 g were divided in 4 groups of 6 animals each. The following schedule of treatment was administered:

Group I: 0.5% CMC

Group II: Diclofenac sodium (20 mg/kg p.o.)

Group III: Alcoholic extract of *M. oleifera* (200 mg/kg p.o.) in 0.5% CMC

Group IV: Alcoholic extract of *M. oleifera* (400 mg/kg p.o.) in 0.5% CMC

Animals were treated with drugs and subsequently 1 h after treatment; 0.1 ml of 1% carrageenan was injected subcutaneously into the planter region of right hind paw to induce edema. The paw volume was measured initially and at 1, 3 and 5 h after carrageenan injection using plethysmographic method of Harris and Spencer (1962). Percentage increase in paw volume from baseline was calculated and compared with control.

### Anti microbial studies

The *in vitro* antimicrobial activity of the *M. oleifera* was studied by broth dilution method and minimum inhibitory concentration was found out. Cold aqueous extract, hot aqueous extract and the alcoholic extract were prepared from the seeds of *M. oleifera*. These extracts at different concentrations (5 - 100 mg/ml) were tested against the organisms *Escherichia coli*, *Staphylococcus aureus* and *pseudomonas aeruginosa*.

**Statistical analysis:** All the results were tested for significance using Student's *t*-test at the probability level of 95%.

## RESULTS

### Effect of *M. oleifera* on Ach and Histamine induced bronchospasm in guinea pigs

Significant increase in preconvulsion time was

observed due to pretreatment with *M. oleifera* (100 mg/kg and 200 mg/kg) when the guinea pigs were exposed to either acetylcholine (0.5%) or histamine (0.25%) aerosol. The increase in preconvulsion time was comparable to that of Ketotifen (1 mg/kg) (Table 1).

### Effect of *M. oleifera* on agonists induced contractions of guinea pig ileum

Alcoholic extract of *M. oleifera* (50 - 150 µg/ml) dose dependently inhibited ileal contractions induced by histamine ( $3.84 \times 10^{-4}$  mM), Ach ( $4.12 \times 10^{-5}$  mM), 5HT ( $5.67 \times 10^{-5}$  mM) and BaCl<sub>2</sub> ( $2.4 \times 10^{-3}$  mM) (Table 2).

### Compound 48/80 induced rat mast cell degranulation

Compound 48/80 (10 µg/ml) produced significant disruption of mast cells which was significantly inhibited in a dose-dependent manner by pretreatment with the alcoholic extract of *M. oleifera*

**Table 1.** Effect of *M. oleifera* on Ach and Histamine induced bronchospasm in guinea pigs

Sr. no.	Treatment	% Increase in preconvulsion time	
		Acetylcholine	Histamine
1	<i>M. oleifera</i> (100 mg/kg p.o.)	41.90 ± 7.94 <sup>*</sup>	27.85 ± 3.96 <sup>***</sup>
2	<i>M. oleifera</i> (200 mg/kg p.o.)	56.31 ± 3.11 <sup>***</sup>	36.13 ± 3.68 <sup>***</sup>
3	Ketotifen fumarate (1 mg/kg p.o.)	28.80 ± 2.77	33.92 ± 3.12

\*Significantly different from baseline (Student's paired *t* test) (n = 6). \**P* < 0.05, \*\*\**P* < 0.001.

**Table 2.** Effect of *M. oleifera* on agonists induced contractions of guinea pig ileum

Conc. of <i>M.oleifera</i> extract (mg/ml)	% Inhibition of Histamine contractions	% Inhibition of Ach contractions	% Inhibition of 5HT contractions	% Inhibition of BaCl <sub>2</sub> contractions
50	48.718 ± 1.16	16.412 ± 0.89	28.306 ± 0.65	48.72 ± 2.05
100	61.88 ± 1.19	27.27 ± 1.18	58.692 ± 0.44	63.385 ± 2.25
150	65.4821 ± 1.24	38.448 ± 1.09	72.216 ± 0.47	80.74 ± 0.99

**Table 3.** Effect of *M. oleifera* extract on compound 48/80 induced rat peritoneal mast cell degranulation

Treatment	Concentration (mg/ml)	% Mast cells degranulation ± S.E.M.	% Inhibition of degranulation
Negative Control	-	1.19 ± 0.96	-
Positive Control	-	74.82 ± 0.467	-
<i>M. oleifera</i> extract	0.5	33.37 ± 1.338 <sup>**</sup>	56.29
<i>M. oleifera</i> extract	1.0	28.71 ± 0.131 <sup>**</sup>	62.62
<i>M. oleifera</i> extract	2.0	24.15 ± 0.432 <sup>**</sup>	68.82
Ketotifen fumarate	10 µg/ml	18.02 ± 0.343 <sup>**</sup>	77.14

N=6 in each group. \*Significantly different from control. \*\**P* < 0.01 (Student's *t* test).

**Table 4.** Effect of *M. oleifera* extract on Egg Albumin induced rat peritoneal mast cell degranulation

Treatment	Concentration (mg/ml)	% Mast cells degranulation $\pm$ S.E.M.	% Inhibition of Degranulation
Negative Control	-	1.19 $\pm$ 0.96	-
Positive Control	-	79.56 $\pm$ 1.126	-
<i>M. oleifera</i> Extract	0.5	29.95 $\pm$ 0.334**	63.30
<i>M. oleifera</i> Extract	1.0	27.77 $\pm$ 0.447**	66.09
<i>M. oleifera</i> Extract	2.0	22.68 $\pm$ 0.317**	72.58
Ketotifen fumarate	0.01	17.01 $\pm$ 0.463**	79.81

N=6 in each group. \*Significantly different from control. \*\* $P < 0.01$  (Student's *t* test).

**Table 5.** Effect of *M. oleifera* on Carageenan induced rat paw edema

Treatment	Dose (mg/kg)	% Increase volume of paw (% anti-inflammatory effect)		
		1 h	3 h	5 h
Control		7.607 $\pm$ 1.03	20.522 $\pm$ 1.579	27.66 $\pm$ 1.685
<i>M. oleifera</i>	200	6.978 $\pm$ 0.097(8.27)	17.96 $\pm$ 0.787*** (12.48)	16.87 $\pm$ 0.96** (39.01)
<i>M. oleifera</i>	400	5.195 $\pm$ 0.453(31.7)	9.214 $\pm$ 1.328* (55.1)	6.40 $\pm$ 0.429** (76.87)
Diclofenac sodium	20	2.577 $\pm$ 0.461(66.11)	8.175 $\pm$ 1.17* (60.16)	4.58 $\pm$ 0.819** (83.43)

Significantly different from control. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  (Student's *t* test).

**Table 6.** Anti microbial study of *M. oleifera*

Sr. No.	Organism	Minimum Inhibitory concentration (mg/ml)		
		Cold water extract	Hot water extract	Alcoholic extract
1	<i>E-coli</i> ATCC-25922	20	100	10
2	<i>P. aeruginosa</i> ATCC-27853	20	-	-
3	<i>S. aureus</i> ATCC-25923	10	50	50

in concentrations of 0.5 - 2.0 mg/ml (Table 3). Egg Albumin (1 mg/ml) induced rat mast cell degranulation was significantly inhibited by pretreatment of the animals with the alcoholic extract of *M. oleifera* in concentrations of 0.5 - 2.0 mg/ml. The protection was comparable to the reference standard Ketotifen (10  $\mu$ g/ml) (Table 4).

#### Effect of *M. oleifera* on Carrageenan induced rat paw edema

Alcoholic extract of *M. oleifera* at the dose of 200 mg/kg and 400 mg/kg decreased rat paw edema (76.87% reduction in edema volume), which was comparable to that of standard Diclofenac Sodium 20 mg/kg (83.43% reduction in edema volume) (Table 5).

#### Anti microbial study

Minimum Inhibitory Concentration for alcoholic extract was low as compared to cold-water extract and hot water extract of *M. oleifera*. Hot water extract and alcoholic extract were ineffective against *P. aeruginosa*. Cold-water extract of *M. oleifera* was found to be more active against Gram-positive bacteria, while alcoholic extract was found to be active against Gram-negative bacteria (Table 6).

## DISCUSSION

The results from our earlier clinical study on *M. oleifera* suggest that, there was appreciable decrease in severity of symptoms of asthma and also simultaneously improvement in lung function

parameters. Also, none of the patients showed change in any general parameters or any adverse effect suggest safety of drug in dose used. Considering the availability along with convenience and efficacy in oral administration, the drug offers a good future in treatment of asthma. Since bronchodilators, mediator release inhibitors, anti-inflammatory drugs and anti-microbials are the different classes of drugs used conventionally in the treatment of bronchial asthma; various animal models and experimental protocols were used in the present study to determine the mechanisms of anti-asthmatic activity of *M. oleifera*.

Bronchial asthma is characterized by increased airway reactivity to spasmogens (Cockcroft, 1983). An initial event in asthma appears to be the release of inflammatory mediators (e.g. Histamine, Tryptase, Leukotrienes and prostaglandins). Some of these mediators directly cause acute bronchoconstriction, airway hyperresponsiveness and bronchial airway inflammation. Spasmolytic drugs like beta adrenergic agonists, xanthine derivatives and anticholinergics relax the airway smooth muscles and are used as quick relief medications in acute asthmatic attacks. Beta adrenergic agonists promote bronchodilation by direct stimulation of beta adrenergic receptors in the airway smooth muscle, that lead to relaxation of bronchial smooth muscle by rapid decrease in airway resistance in vivo. Specific  $\beta_2$  agonists like salbutamol, salmeterol etc. are used since long for symptomatic relief in asthma. In present study, significant increase in preconvulsion time was observed due to pretreatment with *M. oleifera*, when the guinea pigs were exposed to either Ach or histamine aerosol. This bronchodilating effect of *M. oleifera* was comparable to ketotifen. It has been reported that *Albizzia lebeck* (Tripathi and Das, 1977) and *Ocimum sanctum* (Singh and Agarwal, 1991), which are well known anti-asthmatic herbal drugs have similar mechanism of action. Spasmolytic effect of *M. oleifera* was also evaluated by observing the effect of its alcoholic extract on histamine, Ach, 5HT and  $\text{BaCl}_2$  induced ileal contractions. *M.*

*oleifera* produced dose dependent inhibition of ileal contractions induced by histamine, Ach, 5-HT and  $\text{BaCl}_2$ . These indicate that *M. oleifera* has a non-specific spasmolytic activity on smooth muscle. *Tylophora asthmatica* has also been shown to possess non-specific spasmolytic activity (Harnath and Shyamalakumari, 1975). These effects of *M. oleifera* correlate with our earlier results of improvement in the symptoms and lung function parameters of asthmatic subjects.

In addition to bronchodilating activity, a significant number of therapeutic approaches for bronchial asthma have been designed based on the antagonism of specific mediators released from mast cells. Mast cell degranulation is important in the initiation of immediate responses following exposure to allergens. Degranulated cells liberate mediators of inflammation such as histamine, leukotrienes, platelet activating factors and chemotactic factors for eosinophils, neutrophils etc. from mast cells (Cushing, 1957; Bellanti, 1971). They play a significant role in airway inflammatory response such as airway eosinophilia, late asthmatic response and airway hyperresponsiveness as well as in immediate hypersensitivity reaction like bronchial contraction. Degranulation of mast cells has been taken as the criteria of positive anaphylaxis. Ketotifen fumarate, a well-known mast cell stabilizer, reduces synthesis of prostaglandins  $\text{E}_2$ , thromboxane  $\text{A}_2$ , leukotriene  $\text{C}_4$  and  $\text{B}_4$ . It also inhibits release of histamine, serotonin and other inflammatory mediators from mast cells. Simultaneously it blocks  $\text{H}_1$  receptors. Khellin is a compound isolated from *Ammi visnaga* and its structural analogue furanochromone khellin. Cromolyn sodium, which is developed from the structural modification of Khellin (Cox *et al.*, 1970) is the mast cell stabilizer used in the treatment of mild to moderate asthma. *Adhatoda vasica*, *Albizzia lebeck*, *Coleus forskohlii*, *Tylophora asthmatica* etc. are several well known drugs from indigenous plant sources used in asthma and have been reported to have mast cell stabilizing activity (Tripathi *et al.*, 1979; Atal, 1980; Geetha *et al.*, 1981;

Marone *et al.*, 1987). A significant protection of rat peritoneal mast cells from disruption by antigen and compound 48/80 by alcoholic extract of *M. oleifera* points towards its ability to interfere the release and/or synthesis of mediators of inflammation, indicating its mast cell stabilizing activity.

Further, airway inflammation has been demonstrated in all forms of asthma. Even in mild asthma, there is an inflammatory response involving infiltration, particularly with activated eosinophils and lymphocytes, with neutrophils and mast cells. The degree of bronchial hyperresponsiveness and airway obstruction is closely linked to the extent of inflammation (Bousquet *et al.*, 2000). Anti-inflammatory drugs suppress the inflammatory response by inhibiting infiltration and activation of inflammatory cells as well as their synthesis, or release of mediators and the effects of inflammatory mediators. The carrageenan induced paw edema model in rats is known to be sensitive to cyclooxygenase inhibitors. Alcoholic extract of *M. oleifera* possess potent anti-inflammatory activity, which was comparable to that of standard Diclofenac Sodium. Since, serotonin, histamine and prostaglandins are the common mediators of both bronchial asthma and inflammation, the beneficial effect of alcoholic extract of *M. oleifera* could be due to inhibition of their release possibly due to inhibition of the enzyme cyclooxygenase leading to inhibition of prostaglandin synthesis.

In India, the patients with bronchial asthma are commonly prescribed with antibiotics. It has been reported that 78.4% of asthmatic patients receive different antibiotics (Goyal and Patel, 2003). On further investigation, it has been reported that these patients are resistant to many antibiotics prescribed (Goyal and Patel, 2003). It is possible that these patients are suffering from bronchial infection but have been diagnosed as asthmatic patients because of their symptoms like breathlessness. In allopathy, multidrug approach is there where patients receive bronchodilators, corticosteroids along with antibiotics. Sometimes,

nonpathogenic bacteria accumulate due to the bronchial obstruction and plugging, causing serious infection. Plants produce a range of chemical substance to protect themselves from the attack of various pathogenic microorganisms. The substances that can either inhibit the growth of microorganisms or kill them are considered for developing new drugs for various infectious diseases. Use of these medicinal plants can substitute antibiotics to treat associated infection. In the present study, *M. oleifera* possess good Antimicrobial activity when tested against various respiratory pathogens that can be used to control respiratory complications.

In conclusion our data suggests that *M. oleifera* seed kernels have potential anti-asthmatic activity that may be due to its bronchodilator, mast cell stabilization, anti-inflammatory and anti microbial property. Further study is ongoing to characterize the active principles of the ethanolic extract which are responsible for antiasthmatic activity.

## REFERENCES

- Atal CK. (1980) Chemistry and Pharmacology of Vasicine: A new oxyoxic and abortifacient, RRL, Jammu, India. 1980.
- Bellant JA. (1971) Mechanism of Tissue Injury produced by Immunologic Reactions In Immunology, Asian Edn: p. 184, W.B. Saunder Co. Tokyo.
- Bousquet J, Jeffery PK, Busse WW. (2000) Asthma: From bronchoconstriction to airways inflammation and remodeling. *Am. J. Respir. Crit. Care Med.* **161**, 1749-1745.
- Caceres A, Cebreva O, Morales O, Miollinedo P, Mendia P. (1991) Pharmacological properties of *M. oleifera* 1: Preliminary screening for antimicrobial activity. *J. Ethnopharmacol.* **33**, 213-216.
- Caceres A, Saravia A, Rizzo S, Zabala L, De-Leon E, Nave F. (1992) Pharmacological properties of *M. oleifera* 2: Screening for antispasmodic, antiinflammatory and diuretic activity. *J. Ethnopharmacol.* **36**, 233-237.
- Cockcroft DW. (1983) Mechanism of perennial allergic asthma. *Lancet* **2**, 253-256.
- Cox JSG, Beach JE, Blair AM, Clarke AJ, King J, Lee TB. (1970) Disodium chromoglycate (Intal). *Adv.*

- Drug Res.* **5**, 115-196.
- Cushing JE, Campbell DH. (1957) Manifestations of Antigen Antibody Reactions In Principals of Immunology. p. 278, *McGraw-Hill Book Co. Inc. NY.*
- Dajani BM, Sliman NA, Shubair KS, Hamzeh YS. (1981) Bronchospasm caused by intravenous hydrocortisone sodium succinate (Solu-Cortef<sup>®</sup>) in aspirin-sensitive asthmatics. *J. Allergy Clin. Immunol.* **68**, 201-206.
- Geetha VS, Viswanathan S, Kameswaran L. (1981) Comparison of total alkaloids of *Tylophora indica* and disodium cromoglycate on mast cell stabilization. *Indian J. Pharm.* **13**, 199-201.
- Goyal RK, Patel NJ. (2003) Pharmacovigilance for respiratory disorders in N. Gujarat. In: *Pharmacovigilance: an update.* p. 190-201.
- Haalboom JRE, Deenstra A, Stuyvenberg A. (1985) Hypokalaemia induced by inhalation of Fenoterol. *Lancet* **1**, 1125-1127.
- Haranath PSRK, Shyamalakumari S. (1975) Experimental study on the mode of action of *Tylophora asthmatica* in bronchial asthma. *Indian J. Med. Res.* **63**, 661-670.
- Harris JM, Spencer PSJ. (1962) A modified plethysmographic apparatus for recording volume changes in rat paw. *J. Pharm. Pharmacol.* **14**, 464-466.
- Kirtikar KR, Basu BD. (1975) Indian medicinal plants. (M/s Bishen Singh, Mahendra Pal Singh, New Cannaught Place, Dehradun). Vol. 1, Ed. 2, 676-683.
- Marone G, Columbo M, Triggiani M, Cirillo R, Genovese A, Formisano S. (1987) Inhibition of IgE mediated release of histamine and peptide leukotriene from human basophils and mast cells by forskolin. *Biochem. Pharmacol.* **36**, 13-20.
- Nasser SS, Rees PJ. (1993) Theophylline. Current thoughts on the risk and benefits of its use in asthma. *Drug Saf.* **8**, 12-18.
- Nelson HS. (1986) Adrenergic therapy of bronchial asthma. *J. Allergy Clin. Immunol.* **77**, 771-785.
- Sheth UK, Dadkar NK, Kamat NG. (1972) Selected topics in experimental pharmacology. *Kohari book depot. Bombay.* **5**, 63.
- Singh S, Agrawal SS. (1991) Anti asthmatic and anti-inflammatory activity of *Ocimum sanctum*. *Indian J. Pharmacol.* **29**, 306-310.
- Stoloff SW. (1994) The changing role of theophylline in pediatric asthma. *Am. Fam. Physician* **49**, 839-844.
- Tripathi RM, Das PK. (1977) Studies on anti-asthmatic and anti-anaphylactic activity of *Albizia lebeck*. *Indian J. Pharmacol.* **9**, 189-194.
- Tripathi RM, Sen PC, Das PK. (1979) Studies on the mechanism of action of *Albizia lebeck*, an Indian indigenous drug used in the treatment of atopic allergy. *J. Ethnopharmacol.* **1**, 385-396.
- Warrier PK, Nambiar VPK, Ramankutty C. (1997) *Compendium of Indian medicinal plants.* **4**, 59.