

A New Spongilipid from the Freshwater Sponge *Spongilla lacustris*

Jiang-Miao Hu, You-Xing Zhao, Ji-Jun Chen, Ze-Hong Miao,[‡] and Jun Zhou^{*}

State Key Laboratory of Phytochemistry and Plant Resource in West China, Kunming Institute of Botany, the Chinese Academy of Science, Kunming 650204, P. R. China. *E-mail: jzhou@mail.kib.ac.cn

[†]Division of Anti-Tumor Pharmacology, Shanghai Institute of Materia Medica, The Chinese Academy of Sciences, Shanghai 201203, P. R. China

Received November 2, 2008, Accepted March 18, 2009

Key Words: Spongillidae, Freshwater sponge, *Spongilla lacustris*, Spongilipid, Sterol

Sponge is an immemorial species appeared on the earth in the early Cambrian period.¹ It outspread very widely in the Jurassic and the Cretaceous period from marine to freshwater.² Many novel and biologically active compounds have frequently been isolated from the marine sponges these years, while research papers about freshwater sponges were lesser. Freshwater sponge *Spongilla lacustris* has been used as a traditional Chinese medicine^{3,4} for near upon 500 years in China for reinforcing the kidney and supporting yang (aphrodisiac), nevertheless the pharmacy research of this species is fewer.⁵ To make clear the chemical components of *Spongilla lacustris* further, the cosmopolitan species⁶ were selected as our material and leading to the isolation of one new spongilipid, tetracosan-1-ol-1-*O*- β -D-glucopyranoside (**1**), together with ten known compounds, 1-palmitoyl-3- β -D-galactosyl-sn-glycerol (**2**),⁷ cholesterol (**3**),⁸ 5 α -cholest-7-ene-3 β , 6 α -diol (**4**),⁸⁻¹⁰ cholest-5-ene-3 β , 7 β -diol (**5**),¹¹ cholest-5-ene-3 β , 7 α -diol (**6**),¹¹ (22*E*)-cholest-5, 22-diene-3 β , 7 α -diol (**7**),¹¹ 24 ξ -ethylcholest-5-ene-3 β , 7 α -diol (**8**),¹² cholest-7-ene-3 β , 5 α , 6 β -

triol (**9**),¹³ (24*S*)-24-ethyl-cholest-7, 22-ene-3 β , 5 α , 6 β -triol (**10**)¹³ and (24*S*)-24-ethyl-cholest-7-ene-3 β , 5 α , 6 β -triol (**11**).¹³⁻¹⁵ The cytotoxic activity of compound **1** against two human tumor cell lines (A549 and HL-60) was also assessed and showed no activities at concentrations up to 10⁻⁴ mol/L.

Compound **1** was obtained as a white powder, its molecular was assigned as C₃₀H₅₉O₆ from the negative HR-FABMS (*m/z* 515.4298 [M]⁻) and the NMR data, indicating one degree of unsaturation. IR spectrum exhibited absorption bands (3406 cm⁻¹) for hydroxyl. From the ¹³C-NMR (DEPT) and MS spectrum of **1**, thirty carbons were observed as a *D*-glucopyranose moiety (δ 104.8, 78.7, 78.6, 75.3, 71.8, 62.9), and twenty three methylenes (δ 69.9, 32.2, 30.4-29.7, 26.5, 23.0) and one methyl (δ 14.3). The proton signal (δ 4.85, d, *J* = 7.7 Hz, H-1') crossed with carbon (δ 104.8, C-1') in the HSQC spectrum confirm the moiety to be a β -*D*-glucopyranose. The residual twenty four carbons with no unsaturation hint a tetracosanol group, which was proved by GC-MS spectrum of hydrolysate of **1** and ion peak (*m/z* 336 [1 - Glc - H₂O]⁺) in the




Figure 1. The structure of compounds 1-11.




Figure 2. Key HMBC correlations of compound 1.

EIMS spectrum of acetylating of hydrolysate of 1. The ion peak (m/z 397 [$1 - \text{Glc} + \text{AcO}$] $^-$) in the FABMS spectrum after hydrolyzing and acetylating of 1 proved it further. The crosspeak of atom at δ 4.85 (H-1') with carbon at δ 69.9 (C-1) in the HMBC spectrum of 1 (Fig. 2) indicated that the β -D-glucopyranose was linked to C-1 of the tetracosanol group. Compound 1 with four hydroxyls was further confirmed by ion peak (m/z 707 [$1 + 4\text{AcO} - 4\text{H} + \text{Na}$] $^+$) in ESIMS after its acetylation. Thus, compound 1 was determined as tetraacosan-1-yl-1-O- β -D-glucopyranoside.

The ten known compounds were identified on the basis of spectroscopic analysis and comparing spectra data with literature or R_f values with authentic samples.

Experimental

General procedures. Melting points were measured on a XRC-1 micro-melting point apparatus and are uncorrected. MS spectra were obtained on a VG Auto Spec-3000 mass spectrometer. 1D and 2D NMR spectra were recorded on Bruker AM-400 MHz and DRX-500 spectrometers, with chemical shifts (δ) in ppm relative to TMS as internal standard and coupling constants in hertz (Hz). IR spectra were measured with a Bio-Rad FTS-135 IR spectrometer with KBr pellets. UV spectra were measured on a Hitachi UV-3210 spectrophotometer. Silica gel (200-300 mesh) for column chromatography and preparative thin-layer chromatography were the products of the Qindao Marine Chemical Ltd., Qingdao, P. R. China. Sephadex LH-20 for chromatography was purchased from Amersham Biosciences. Reversed-phase chromatography was with RP-18 (LiChrorep, 40-63 μm , Merck, Darmstadt, Germany).

Animal material. The dried green sponge *Spongilla lacustris* (4.5 kg) was collected from Tenchong, Yunnan Province, P. R. China, in November 2004 and identified by Prof. Li-zhen Wang, Yunnan University, Kunming, Yunnan, P. R. China. A voucher specimen (Zhai-1) is deposited at the Kunming Institute of Botany, Chinese Academy of Science, Kunming, Yunnan, P. R. China.

Extraction and isolation. The air-dried sponge (4.0 kg) was powdered and extracted three times with 80 % ethanol aq. (30 L \times 3) at room temperature for 24 hours each time and filtered. The filtrate was evaporated under reduced pressure to give a residue, which was suspended in water and partitioned successively with petroleum ether, EtOAc and *n*-BuOH. The EtOAc-soluble fraction was concentrated to give a deep green gum (180.0 g) and subjected to column chromatography (CC) (silica gel, CHCl_3 - CH_3OH , 10 : 0 \rightarrow 8 : 2) to give 5 fractions A-E. Fraction B (10 g) was further purified by CC (silica gel, CHCl_3 - CH_3OH , 92 : 8) and CC (Sephadex LH-20, CHCl_3 -

CH_3OH , 7 : 3) to afford compound 3 (1.0 g) and 4 (15 mg). Fraction C was treated same as fraction B to produce compound 5 (8 mg) and a white crystal (40 mg), the crystal was further separate by semi preparative HPLC (Agilent ODS-C18, CH_3OH - H_2O , 91 : 9) to produce compound 6 (5 mg), 7 (6 mg) and 8 (4 mg). Fraction D was treated same as fraction C and afford compound 9 (4 mg), 10 (5 mg) and 11 (4 mg). The *n*-BuOH soluble fraction was concentrated to give a deep brown gum (100.0 g), part of which (95.0 g) was subjected to CC (silica gel, CHCl_3 - CH_3OH , 9 : 1 \rightarrow 7 : 3) to give eight fractions (I-VIII). Fraction III (9.2 g) was further chromatographed on silica gel by CHCl_3 - CH_3OH - H_2O (80 : 20 : 5) to afford nine fractions III-1-III-9. Fraction III-5 (230 mg) was purified by CC (Sephadex LH-20, MeOH) and CC (RP-18, MeOH- H_2O , 4 : 6) to produce compound 1 (20 mg). Fraction III-7 (150 mg) was treated same as III-5 to yield compound 2 (6 mg).

Lacustrisglycoside A (1): White powder, m.p. 82-84 $^\circ\text{C}$; $[\alpha]_D^{25} = -4.3$ (c 1.04, pyridine); UV (MeOH) λ_{max} (log ϵ): 202.0 (6.11) nm; IR (KBr) ν_{max} : 3406 (OH), 2919, 2850, 1471, 1374, 1255, 1170, 1098, 1074, 1038, 719, 652 cm^{-1} ; ^1H NMR (400 MHz, $\text{C}_5\text{D}_5\text{N}$) δ 4.85 (d, $J = 7.7$ Hz, 1H, H-1'), 4.57 (brd, $J = 10.4$ Hz, 1H, H-6' α), 4.40 (dd, $J = 10.4, 5.3$ Hz, 1H, H-6' β), 4.26 (m, 1H, H-5'), 4.25 (m, 1H, H-4'), 4.08 (dd, $J = 9.0, 7.2$ Hz, 1H, H-1 α), 4.05 (dd, $J = 7.7, 8.0$ Hz, 1H, H-2'), 3.97 (m, 1H, H-3'), 3.66 (dd, $J = 9.0, 7.2$ Hz, 1H, H-1 β), 1.65 (m, 2H, H-2), 1.30-1.19 (m, 42H, H-3-23), 0.85 (t, $J = 6.0$ Hz, 3H, H-24); ^{13}C NMR (100 MHz, $\text{C}_5\text{D}_5\text{N}$) δ : 104.8 (C-1'), 78.7 (C-5'), 78.6 (C-3'), 75.3 (C-2'), 71.8 (C-4'), 69.9 (C-1), 62.9 (C-6'), 32.2 (C-22), 30.4-29.7 (C-4-21), 29.9 (C-2), 26.5 (C-3), 23.0 (C-23), 14.3 (C-24); FABMS (negative ion) m/z 515 (59) $[\text{M}-\text{H}]^-$, 473 (100); HR-FABMS (negative ion) m/z 515.4298 $[\text{M}]^-$ (calcd. for $\text{C}_{30}\text{H}_{59}\text{O}_6$, 515.4311).

Acetylate and hydrolysis. The sample (compound 1, 2 mg) was dissolved in MeOH/ H_2O (8 : 2) in a 50 mL rockered flask and 5 drops of HCl (36%) were added. After refluxed at 80 $^\circ\text{C}$ for 4 hours, the hydrolysate was allowed to cool and separated successfully between petroleum ether and MeOH/ H_2O (8 : 2). The petroleum ether layer was concentrated and sent for EIMS and GC-MS. The residual petroleum ether layer dissolved in Ac $_2$ O/pyridine (6 : 1) in a sealed micro-flask and reacted at 60 $^\circ\text{C}$ for 4 hours, and then the acetic reactant was subjected to positive FABMS analysis.

The sample (compound 1, 1 mg) was dissolved in Ac $_2$ O/pyridine (6 : 1) in a sealed micro-flask and reacting at 60 $^\circ\text{C}$ for 4 hours, then the reactant was subjected to positive ESIMS analysis.

Cytotoxic assay. Compound 1 was tested for its cytotoxic effects against human lung carcinoma A549 and human leukemia HL-60 cell lines using the sulforhodamine B (SRB)

assay and the methyl-thiazol-tetrazolium (MTT) assay, respectively.

Acknowledgments. The authors are grateful to the helps from Dai-Yun Li and to the members of the analytical group in the State Key Laboratory of Phytochemistry and Plant Resource in West China, Kunming Institute of Botany, for the spectral measurements. The authors are also grateful to Prof. Li-Zhen Wang for identification of animal samples.

References

1. Djerassi, C.; Lam, W. K. *Acc. Chem. Res.* **1991**, *24*, 69.
 2. He, X. Y.; Xu, G. R. *Paleontology tutorial*; Geologic Press: 1993; p 49.
 3. Li, S. Z. *Bencao Gangmu*; People's sanitation express: 1975; p 2380.
 4. Zhonghua Bencao editing committee of State Administration of Traditional Chinese Medicine of the People's Republic of China, *Zhonghua Bencao (Volume 9)*; Shanghai Scientific and Technical Publishers: 1999; p 7930.
 5. Manconi, R.; Piccialli, V.; Pronzato, R.; Sica, D. *Comparative Biochemistry and Physiology* **1988**, *91B*, 237.
 6. Cheng, L.; Wang, K. L.; Yu, Z. M.; Yang, Y. *Zoological Research* **1991**, *12*, 235.
 7. Pettit, G. R.; Bond, T. J.; Herald, D. L.; Penny, M.; Doubek, D. L.; Williams, M. D.; Pettit, R. K.; Hooper, J. N. A. *Can. J. Chem.* **1997**, *75*, 920.
 8. Piccialli, V.; Sica, D. *J. Nat. Prod.* **1987**, *50*, 915.
 9. Piccialli, V.; Sica, D. *J. Nat. Prod.* **1986**, *49*, 779.
 10. Simone, F. D.; Dini, A.; Finamore, E.; Minale, L.; Pizza, C.; Riccio, R. *J. Chem. Soc (Perkin I)* **1981**, 1855.
 11. Notaro, G.; Piccialli, V.; Sica, D. *J. Nat. Prod.* **1992**, *55*, 1588.
 12. Guerriero, A.; D'Ambrosio, M.; Pietra, F. *J. Nat. Prod.* **1993**, *56*, 1962.
 13. Madaio, A.; Piccialli, V.; Sica, D. *J. Nat. Prod.* **1989**, *52*, 952.
 14. Das, B.; Srinivas, K. V. N. S. *J. Nat. Prod.* **1993**, *56*, 2210.
 15. Das, B.; Srinivas, K. V. N. S. *J. Nat. Prod.* **1992**, *55*, 1310.
-