

Novel *N*-Hydroxymethylation of Phthalimide by Titanium Dioxide Photocatalyst in Methanol

Hye Jong Lee, Young Shik Kong, and Sung Sik Kim*

Department of Chemistry and Research Institute of Physics and Chemistry (RINPAC), Chonbuk National University, Chonju 561-756, Korea. *E-mail: hvssk@chonbuk.ac.kr
Received January 8, 2009, Accepted February 6, 2009

Key Words: Phthalimide, Titanium dioxide, Photoreduction, Hydroxymethylation

Photochemical reactions of some nitrogen-containing compounds catalyzed by semiconductors such as titanium dioxide (TiO₂) in solution phase have been extensively investigated.¹⁻⁷ It was reported that various type of compounds are produced depending on the structures of reactants. For example, secondary amines were generated from primary amines by an aqueous suspension of TiO₂/Pt.³ Photocatalytic oxidation of primary amines on irradiated TiO₂ powders suspended in anhydrous acetonitrile led to the corresponding symmetrical *N*-alkylidene amines in good yields.⁴ UV irradiation of methylamine in the presence of TiO₂ and oxygen gave ammonia, carbon dioxide, etc.⁵ Trimethylamine can be bound to TiO₂ surface. With photoirradiation in the presence of oxygen, the adsorbed trimethylamine decomposed and formed CO₂, NCO, etc.⁶ Imides were also generated from the reaction of *N*-acetylamine in the presence of oxygenated aqueous suspensions of TiO₂.⁷ We have been interested in heterogeneous photocatalysis with dispersed titanium dioxide in the viewpoint of synthetic photochemistry. Chemical transformations of organic compounds by TiO₂ photocatalyst are of great interest, because of the diversity in its role. Several attempts have been made to get some useful organic molecules in organic solvent, instead of aqueous phase. It was found that anhydrides were cleaved to give the monoesters of dicarboxylic acids in good yields, when they were irradiated on TiO₂ in methanol with UV light.⁸

Herein we report that 350 nm UV irradiation of phthalimide **1** in methanol in the presence of TiO₂ photocatalyst afforded *N*-hydroxymethylphthalimide **5** and *N*-hydroxymethyl-3-hydroxyphthalimidine **6** as the major product. A literature survey of existing photochemical reactions of phthalimides without TiO₂ indicates that irradiation of phthalimide **1** with UV light gave several kinds of photoreduction products such as 3-hydroxyphthalimidine **2**, its dimer **3**, and *C*-hydroxymethyl-3-hydroxyphthalimidine **4** depending on the reaction conditions.⁹⁻¹³ Similarly, upon irradiation of **1** in methanol gave **2** in low yield. Meanwhile, irradiation of phthalimide **1** in methanol (40 mL) in the presence of NaOH (1.5 M) reported the production of **3** and **4** in moderate yields, 35% and 40% yields, respectively.

As shown in Table 1, when phthalimide **1** was just irradiated in methanol without TiO₂ photocatalyst with 350 nm UV light for 24 h, only 3-hydroxyphthalimidine **2** was produced as a simple reduction product (< 5% yield), in which *ca.* 95% of **1** was recovered. In contrast, irradiation of **1** in methanol in the



presence of 1 equivalent of TiO₂ afforded *N*-hydroxymethylphthalimide **5** in 74% yield, as well as **2** (14% yield) with a quantitative conversion of the starting material. When **1** was irradiated in methanol in the presence of TiO₂ containing 3 × 10⁻¹ vol% of water, only **5** was produced in 21% yield, as determined by TLC and NMR of the reaction mixture. It is well understood that formaldehyde is produced from the oxidation of methanol by TiO₂ photocatalyst.¹⁴⁻¹⁶ Thus, the hydroxymethylation of phthalimide **1** could be rationalized by the formation of formaldehyde molecules from methanol by TiO₂ photocatalyst. Interestingly, when this reaction was carried in methanol containing 3 × 10⁻² vol% of water, *N*-hydroxymethyl-3-hydroxyphthalimidine **6** was observed in 85% yield as the major product along with **2** (15% yield). ¹H-NMR spectrum of **6** taken in chloroform-*d* shows a singlet peak of methyne proton (CH) at δ 6.04. Methylene protons (CH₂) of *N*-hydroxymethyl group appeared at δ 5.22 (*J* = 11.2 Hz) and δ 5.00 (*J* = 11.2 Hz) as doublet peaks. GC/MS spectrum of **6** shows a parent ion peak at *m/e* 179 (C₉H₉O₃N; Calculated, 179.175).

Because 3-hydroxyphthalimidine **2** is one of heterocyclic compounds of biological relevance that have attracted considerable interest in recent years,¹⁷ it is also interesting to note that while *N*-hydroxymethylation was not observed in the photoreactions of *N*-substituted phthalimides such as *N*-methylphthalimide and *N*-phenylphthalimide, only 3-hydroxyphthalimidines were produced in 79% and 75% yield, respectively. Especially, *N*-bromophthalimide was completely inert, when irradiated in methanol in the presence of TiO₂.

Table 1. TiO₂-mediated photoreactions of phthalimide **1** in methanol.

TiO ₂ (mmol)	H ₂ O added	Conversion of 1 (%)	Photoproducts and Yields ^a		
			% 2	% 5	% 6
0	-	5	< 5	-	-
1	-	100	14	74	-
1	3 × 10 ^{-1b}	21	-	21	-
1	3 × 10 ^{-2b}	100	15	-	85

^aIsolated chemical yields from 1 mmol of **1** in methanol (40 mL).
^bvolume% based on methanol solvent.



In worth to note, some other semiconductors such as SnO, SnO₂, ZrO₂, WO₃, and ZnO had no effect on the photoreactions of phthalimides in methanol. Scheme 1 shows that when the reductive dimerization product **3** and C-hydroxymethylation product **4**, which were yielded as reduction products by the irradiation of phthalimide **1** in methanol containing sodium hydroxide, were irradiated in the methanol in the presence of TiO₂, the two types of 1,2-diols were converted into N-hydroxymethylation product **6** in quantitative yields as the final product.

In the same manner, Scheme 2 shows that the primary photoreduction products (**7** and **8**) were produced with >90% conversion of **1** by the direct irradiation of **1** in ethanol and *i*-propyl alcohol, respectively, and that **7** and **8** were also converted into **6** in quantitative yields, when they were irradiated in methanol in the presence of TiO₂.

In summary, we observed for the first time that irradiation of phthalimide **1** in methanol in the presence of TiO₂ affords N-hydroxymethyl-3-hydroxyphthalimidine **6** as the final product in good yield. The chemical yield of **6** depends on the amount of water in methanol solvent. Maximum yield was

achieved when **1** was irradiated in methanol containing 3 × 10⁻² vol% of water. The primary photoreduction products, *i.e.*, reductive dimer **3** and C-hydroxymethyl-3-hydroxyphthalimidine **4**, were converted into the N-hydroxymethyl derivative **6** in quantitative yields by the TiO₂-mediated photoreactions. In addition, C-hydroxyethyl-3-hydroxyphthalimidine **7** and C-hydroxyisopropyl-3-hydroxyphthalimidine **8**, produced in ethanol and *i*-propyl alcohol, respectively, were also converted into **6** by the TiO₂-mediated photoreactions. Mechanistic study and synthetic application of this photoreaction are under progress.

Acknowledgments. This work was supported by the Korea Science and Engineering Foundation (Grant No. R01-2002-000-00272-0). This paper was (partially) supported by CBNU funds for overseas research, 2006.

References

- Kasturirangan, H.; Ramakrishnan, V.; Kuriacose, J. C. *J. Catal.* **1981**, *69*, 216.
- Fox, M. A.; Chen, M.-J. *J. Am. Chem. Soc.* **1983**, *105*, 4497.
- Nishimoto, S.; Ohtani, B.; Yoshikawa, T.; Kagiya, T. *J. Am. Chem. Soc.* **1983**, *105*, 7180.
- Fox, M. A.; Younathan, J. N. *Tetrahedron* **1986**, *42*, 6285.
- Liao, L.-F.; Wu, W.-C.; Chuang, C.-C.; Lin, J.-L. *J. Phys. Chem. B* **2001**, *105*(25), 5928.
- Lien, C.-F.; Lin, Y.-F.; Lin, Y.-S.; Chen, M.-T.; Lin, J.-L. *J. Phys. Chem. B* **2005**, *109*, 10962.
- Pavlik, J. W.; Tantayanon, S. *J. Am. Chem. Soc.* **1981**, *103*, 6755.
- Kim, S. S.; Mah, Y. J.; Lee, H. J.; Park, S. K. *J. Photosci.* **2003**, *10*(3), 241.
- Kanaoka, Y. *Acc. Chem. Res.* **1978**, *11*, 407.
- Yoon, U. C.; Mariano, P. S. *Acc. Chem. Res.* **2001**, *34*(7), 523.
- Griesbeck, A. G.; Kramer, W.; Schmickler, H. *Org. Lett.* **2001**, *3*(4), 537.
- Yoon, U. C.; Kwon, H. C.; Hyung, T. G.; Choi, K. H.; Oh, S. W.; Yang, S.; Zhao, Z.; Mariano, P. S. *J. Am. Chem. Soc.* **2004**, *126*(4), 1110.
- Najera, F.; Garcia-Segura, R.; Perez-Inestrosa, E.; Sanchez-Sanchez, C.; Suau, R. *Photochem. Photobiol.* **2006**, *82*, 248.
- Effects of water addition on the methanol oxidation on Pt/TiO₂ photocatalyst was studied by time-resolved infrared absorption spectroscopy: Yamakata, A.; Ishibashi, T.-A.; Onishi, H. *J. Phys. Chem. B* **2003**, *107*, 9820.
- Bronkema, J. L.; Leo, D. C.; Bell, A. T. *J. Phys. Chem. C* **2007**, *111*, 14530.
- Gondal, M. A.; Hameed, A.; Yamani, Z. H.; Arfaj, A. *Chem. Phys. Lett.* **2004**, *392*, 372.
- Alajarin, M.; Sanchez-Andrada, P.; Lopez-Leonardo, C.; Alvarez, A. *J. Org. Chem.* **2005**, *70*, 7617.