

## DIRECT SUM ON WFI-ALGEBRAS

EUN HWAN ROH\*

ABSTRACT. The notion of subdirect sum and direct sum in WFI-algebras is introduced, and several properties are investigated.

### 1. Introduction

In 1990, W. M. Wu [8] introduced the notion of fuzzy implication algebras (FI-algebra, for short), and investigated several properties. In [7], Z. Li and C. Zheng introduced the notion of distributive (resp. regular, commutative) FI-algebras, and investigated the relations between such FI-algebras and MV-algebras. In [1], Y. B. Jun discussed several aspects of WFI-algebras, and gave a characterization of a WFI-algebra. He introduced the notion of associative (resp. normal, medial) WFI-algebras, and investigated several properties. He gave conditions for a WFI-algebra to be associative/medial, and provided characterizations of associative/medial WFI-algebras, and showed that every associative WFI-algebra is a group in which every element is an involution. He also verified that the class of all medial WFI-algebras is a variety. Y. B. Jun and S. Z. Song [6] introduced the notions of simulative and/or mutant WFI-algebras and investigated some properties. They established characterizations of a simulative WFI-algebra, and gave a relation between an associative WFI-algebra and a simulative WFI-algebra. They also found some types for a simulative WFI-algebra to be mutant. Jun, Park and Roh [5] introduced the concept of ideals of WFI-algebras. They gave relations between a filter and an ideal, and provided characterizations of an ideal. Also they established an extension property for an ideal. In [2] and [3], Y. B. Jun introduced the concept of perfect filters, concrete filters, mote, beam and osculatory in WFI-algebra. He gave relations

---

Received July 01, 2009; Accepted August 14, 2009.

2000 Mathematics Subject Classification: Primary 03G25, 03C05, 08A05.

Key words and phrases: WFI-algebras, (closed) filter, subdirect sum, direct sum.

\*This paper was supported by the research fund in Chinju National University of Education, 2009.

between various these notions. Y. B. Jun and C. H. Park [4] discussed uncanny filters, and investigated related properties. In this paper, we introduce the notion of subdirect sum and direct sum on WFI-algebra. Also we provide related properties.

## 2. Preliminaries

Let  $K(\tau)$  be the class of all algebras of type  $\tau = (2, 0)$ . By a *WFI-algebra* we mean a system  $\mathfrak{X} = (X, \ominus, 1) \in K(\tau)$  such that for all  $x, y, z \in X$ :

- (a1)  $x \ominus (y \ominus z) = y \ominus (x \ominus z)$ ,
- (a2)  $(x \ominus y) \ominus ((y \ominus z) \ominus (x \ominus z)) = 1$ ,
- (a3)  $x \ominus x = 1$ ,
- (a4)  $x \ominus y = y \ominus x = 1 \Rightarrow x = y$ .

For the convenience of notation, we shall write  $[x, y_1, y_2, \dots, y_n]$  for

$$(\dots((x \ominus y_1) \ominus y_2) \ominus \dots) \ominus y_n.$$

We define  $[x, y]^0 = x$ , and for  $n > 0$ ,  $[x, y]^n = [x, y, y, \dots, y]$ , where  $y$  occurs  $n$ -times.

LEMMA 2.1. [1] *In a WFI-algebra  $\mathfrak{X}$ , the following are true:*

- (b1)  $x \ominus [x, y]^2 = 1$ ,
- (b2)  $1 \ominus x = 1 \Rightarrow x = 1$ ,
- (b3)  $1 \ominus x = x$ ,
- (b4)  $x \ominus y = 1 \Rightarrow (y \ominus z) \ominus (x \ominus z) = 1, (z \ominus x) \ominus (z \ominus y) = 1$ ,
- (b5)  $(x \ominus y) \ominus 1 = (x \ominus 1) \ominus (y \ominus 1)$ ,
- (b6)  $[x, y]^3 = x \ominus y$ .

A nonempty subset  $S$  of a WFI-algebra  $\mathfrak{X}$  is called a *subalgebra* of  $\mathfrak{X}$  if  $x \ominus y \in S$  whenever  $x, y \in S$ . A nonempty subset  $F$  of a WFI-algebra  $\mathfrak{X}$  is called a *filter* of  $\mathfrak{X}$  if it satisfies:

- (c1)  $1 \in F$ ,
- (c2)  $x \ominus y \in F$  and  $x \in F$  imply  $y \in F$  for all  $x, y \in X$ .

A filter  $F$  of a WFI-algebra  $\mathfrak{X}$  is said to be *closed* [1] if  $F$  is also a subalgebra of  $\mathfrak{X}$ .

LEMMA 2.2. [1] *Let  $F$  be a filter of a WFI-algebra  $\mathfrak{X}$ . Then  $F$  is closed if and only if  $x \ominus 1 \in F$  for all  $x \in F$ .*

LEMMA 2.3. [1] *In a finite WFI-algebra, every filter is closed.*

We now define a relation “ $\preceq$ ” on  $\mathfrak{X}$  by  $x \preceq y$  if and only if  $x \ominus y = 1$ . It is easy to verify that a WFI-algebra is a partially ordered set with respect to  $\preceq$ . For a WFI-algebra  $\mathfrak{X}$ , the set

$$\mathcal{S}(\mathfrak{X}) := \{x \in X \mid x \preceq 1\}$$

is called the *simulative part* of  $\mathfrak{X}$  ([6]). Note that  $\mathcal{S}(\mathfrak{X})$  is a subalgebra of  $\mathfrak{X}$ .

LEMMA 2.4. [6] *Let  $\mathfrak{X}$  be a WFI-algebra. Then  $\mathcal{S}(\mathfrak{X})$  is a filter of  $X$ .*

The *doubly simulative part* of  $\mathfrak{X}$  [5] is defined to be the set

$$\mathcal{DS}(\mathfrak{X}) := \{x \in X \mid [x, 1]^2 = x\}.$$

Obviously,  $1 \in \mathcal{DS}(\mathfrak{X})$  and  $\mathcal{DS}(\mathfrak{X}) \cap \mathcal{S}(\mathfrak{X}) = \{1\}$ .

### 3. Main results

In what follows let  $\mathfrak{X}$  denote a WFI-algebra  $(X; \ominus, 1)$  unless otherwise specified.

LEMMA 3.1. *For any  $\mathfrak{X}$ , if  $a \in X$ , then the following conditions are equivalent:*

- (1)  $a \leq x \Rightarrow a = x$  for any  $x \in X$ .
- (2)  $[a, 1]^2 = a$ .
- (3) there is  $x \in X$  such that  $a = x \ominus 1$ .

*Proof.* (1)  $\Rightarrow$  (2). By (b1), we have  $[a, 1]^2 = a$ .

(2)  $\Rightarrow$  (1). Let  $a \preceq x$  for any  $x \in X$ . Then we have

$$x \ominus a = x \ominus [a, 1]^2 = (a \ominus 1) \ominus (x \ominus 1) = 1.$$

and so  $a = x$  by (a4).

(2)  $\Rightarrow$  (3). We have  $a = [a, 1]^2 = x \ominus 1$ , where  $x := a \ominus 1$ .

(3)  $\Rightarrow$  (2). Suppose that  $a = x \ominus 1$  for some  $x \in X$ . Then we have

$$[a, 1]^2 \ominus a = [x \ominus 1, 1]^2 \ominus (x \ominus 1) = (x \ominus 1) \ominus (x \ominus 1) = 1,$$

and so  $[a, 1]^2 = a$  by (a4). □

LEMMA 3.2. [1] *Let  $\mathfrak{X}$  be a WFI-algebra. Then the following conditions are equivalent:*

- (1)  $[a, 1]^2 = a$  for any  $a \in X$ .
- (2)  $[a, x]^2 = a$  for any  $a, x \in X$ .

We now consider the generated filters in WFI-algebras.

DEFINITION 3.3. Let  $S$  be a subset of  $\mathfrak{X}$ . We call the least filter of  $\mathfrak{X}$  containing  $S$ , the *generated filter* of  $\mathfrak{X}$  by  $S$ , denoted by  $\langle S \rangle$ .

It is obvious that the intersection of any filter family of  $\mathfrak{X}$  is a filter. So the generated filter is well-defined and we have an obvious assertion as follows.

LEMMA 3.4. Let  $\mathfrak{X}$  be a WFI-algebra and  $S, T \subseteq X$ . If  $S \subseteq T$ , then  $\langle S \rangle \subseteq \langle T \rangle$ . In particular, if  $S = \emptyset$ , then  $\langle S \rangle = \{1\}$ .

We denote  $\langle \{a_1, a_2, \dots, a_n\} \rangle$  by  $\langle a_1, a_2, \dots, a_n \rangle$  in brevity. Sometimes, the filter  $\langle a \rangle$  generated by one element  $a$  is also called a *principal filter* of  $\mathfrak{X}$ . The next theorem gives the description of elements in  $\langle S \rangle$ .

THEOREM 3.5. Let  $S$  be a nonempty subset of  $\mathfrak{X}$  and let

$$G := \{x \in X \mid a_1 \ominus (a_2 \ominus (\dots \ominus (a_{n-1} \ominus (a_n \ominus x)) \dots)) = 1 \text{ for some } a_1, a_2, \dots, a_n \in S\}.$$

Then  $\langle S \rangle = G \cup \{1\}$ .

*Proof.* The proof is straightforward. □

THEOREM 3.6. Let  $F$  be a filter of  $\mathfrak{X}$ . Define a binary operation  $\equiv$  on  $X$  as follow:

$$x \equiv y(\text{mod } F) \Leftrightarrow x \ominus y \in F \text{ and } y \ominus x \in F$$

for any  $x, y \in X$ . Then  $\equiv$  is a congruence on  $\mathfrak{X}$ .

*Proof.* The proof is standard. □

THEOREM 3.7. Let  $F$  be a filter of  $\mathfrak{X}$  and  $\equiv$  be a congruence relation on  $\mathfrak{X}$  defined by Theorem 3.6. We denote

$$\equiv_x := \{y \in X \mid x \equiv y(\text{mod } F)\} \text{ and } X/\equiv := \{\equiv_x \mid x \in X\}.$$

Then the quotient algebra  $\mathfrak{X}/\equiv := (X/\equiv; \odot, \equiv_1)$  is a WFI-algebra, where the operation  $\odot$  on  $X/\equiv$  given by  $\equiv_x \odot \equiv_y := \equiv_{x \odot y}$ .

*Proof.* The proof is immediate. □

DEFINITION 3.8. Let  $F_1$  and  $F_2$  be filters of  $\mathfrak{X}$  such that  $X = \langle F_1 \cup F_2 \rangle$  and  $F_1 \cap F_2 = \{1\}$ , then  $\mathfrak{X}$  is called the *subdirect sum* of  $F_1$  and  $F_2$ , denoted by  $\mathfrak{X} = F_1 \oplus F_2$ .

EXAMPLE 3.9. Let  $X := \{1, a, b\}$  be a set with the following Cayley table.

$$\begin{array}{c|ccc}
 \ominus & 1 & a & b \\
 \hline
 1 & 1 & a & b \\
 a & 1 & 1 & b \\
 b & 1 & a & 1
 \end{array}$$

Then  $\mathfrak{X} = (X; \ominus, 1)$  is a WFI-algebra. It is easy to verify that  $F_1 = \{1, a\}$  and  $F_2 = \{1, b\}$  are filters of  $X$ , and  $\mathfrak{X} = F_1 \oplus F_2$ .

THEOREM 3.10. Let  $F_1$  and  $F_2$  be closed filters of  $\mathfrak{X}$ . If  $\mathfrak{X} = F_1 \bar{\oplus} F_2$ . Then there are unique  $a \in F_1$  and  $b \in F_2$  such that  $x \equiv a \pmod{F_2}$  and  $y \equiv b \pmod{F_1}$ .

*Proof.* Let us first prove that there is unique  $a \in F_1$  such that  $x \equiv a \pmod{F_2}$ . Let  $x \in X$ . Since  $X = \langle F_1 \cup F_2 \rangle$ , by Theorem 3.5, we know that there exist  $b_1, b_2, \dots, b_n \in F_2$  such that  $b_1 \ominus (b_2 \ominus (\dots \ominus (b_n \ominus x) \dots)) = 1$ . Put  $a := b_1 \ominus (b_2 \ominus (\dots \ominus (b_n \ominus x) \dots))$ , then  $a \in F_1$ . Thus we have

$$b_1 \ominus (b_2 \ominus (\dots \ominus (b_n \ominus (a \ominus x)) \dots)) = a \ominus a = 1,$$

and so  $a \ominus x \in F_2$ . Moreover, since  $x \ominus a = b_1 \ominus (b_2 \ominus (\dots \ominus (b_n \ominus 1) \dots))$ , by  $F_2$  being a closed filter of  $\mathfrak{X}$ , it follows  $x \ominus a \in F_2$ . Therefore we have  $x \equiv a \pmod{F_2}$ . Let  $a, a' \in F_1$  such that  $x \equiv a \pmod{F_2}$  and  $x \equiv a' \pmod{F_2}$ . By the symmetry and transitivity of congruence, we have  $a \equiv a' \pmod{F_2}$ , and so  $a \ominus a' \in F_2$  and  $a' \ominus a \in F_2$ . Also, since  $F_1$  is a closed filter of  $\mathfrak{X}$  and  $a, a' \in F_1$ , we obtain  $a \ominus a' \in F_1$  and  $a' \ominus a \in F_1$ . Hence we get  $a \ominus a' \in F_1 \cap F_2$  and  $a' \ominus a \in F_1 \cap F_2$ . Since  $F_1 \cap F_2 = \{1\}$ , we have  $a \ominus a' = 1 = a' \ominus a$ . Therefore we get  $a = a'$ .

Similarly, there is unique  $b \in F_2$  such that  $x \equiv b \pmod{F_1}$ . This completes the proof.  $\square$

THEOREM 3.11. Let  $F$  be a closed filter of  $\mathfrak{X}$ . If  $\mathcal{S}(\mathfrak{X}) \cap F = \{1\}$ , then  $F \subseteq \mathcal{S}(\mathfrak{X})^*$ . Further, if  $\mathfrak{X} = \mathcal{S}(\mathfrak{X}) \bar{\oplus} F$ , then  $F = \mathcal{S}(\mathfrak{X})^*$ , where  $\mathcal{S}(\mathfrak{X})^* := \{x \in X \mid a \ominus x = x \text{ for any } a \in \mathcal{S}(\mathfrak{X})\}$ .

*Proof.* Let  $\mathcal{S}(\mathfrak{X}) \cap F = \{1\}$  and  $x \in F$ . Then by (b1) and Lemma 2.4, we have  $[a, x]^2 \in A$  for any  $a \in \mathcal{S}(\mathfrak{X})$ . Since  $x \ominus [a, x]^2 = (a \ominus x) \ominus 1 = (a \ominus 1) \ominus (x \ominus 1) = x \ominus (1 \ominus 1) = x \ominus 1$  and  $F$  is a closed filter of  $\mathfrak{X}$ , we have  $[a, x]^2 \in F$ . Hence we get  $[a, x]^2 = 1$ . Also,  $x \ominus (a \ominus x) = a \ominus (x \ominus x) = 1$ . therefore  $a \ominus x = x$  and so  $x \in \mathcal{S}(\mathfrak{X})^*$ , i.e.  $F \subseteq \mathcal{S}(\mathfrak{X})^*$ .

On the other hand, if  $\mathfrak{X} = \mathcal{S}(\mathfrak{X}) \bar{\oplus} F$ , then  $\mathcal{S}(\mathfrak{X}) \cap F = \{1\}$ , and so  $F \subseteq \mathcal{S}(\mathfrak{X})^*$ . For any  $x \in \mathcal{S}(\mathfrak{X})^*$ , there is  $a \in \mathcal{S}(\mathfrak{X})$  such that  $x \equiv a \pmod{F}$  by

Theorem 3.10. Thus we get  $a \ominus x \in F$ . Note that  $a \ominus x = x$ . Hence we have  $\mathcal{S}(\mathfrak{X})^* \subseteq F$ . Therefore  $F = \mathcal{S}(\mathfrak{X})^*$ . This completes the proof.  $\square$

DEFINITION 3.12. Let  $F_1$  and  $F_2$  be filters of  $\mathfrak{X}$  and  $\mathfrak{X} = F_1 \bar{\oplus} F_2$ . If for any  $a \in F_1$  and  $b \in F_2$ , there exists  $x \in X$  such that  $x \equiv a \pmod{F_2}$  and  $y \equiv b \pmod{F_1}$ , then we say  $\mathfrak{X}$  is the *direct sum* of  $F_1$  and  $F_2$ , denoted by  $\mathfrak{X} = F_1 \oplus F_2$ .

We remark that  $\mathcal{DS}(\mathfrak{X})$  is generally not a filter of  $\mathfrak{X}$ . Let  $X := \{1, a, b\}$  be a set with the following Cayley table.

$$\begin{array}{c|ccc}
 \ominus & 1 & a & b \\
 \hline
 1 & 1 & a & b \\
 a & 1 & 1 & b \\
 b & b & b & 1
 \end{array}$$

Then  $\mathfrak{X} = (X; \ominus, 1)$  is a WFI-algebra. Then  $\mathcal{DS}(\mathfrak{X}) = \{1, b\}$  is not a filter of  $\mathfrak{X}$  since  $b \ominus a \in \mathcal{DS}(\mathfrak{X}), b \in \mathcal{DS}(\mathfrak{X})$  and  $a \notin \mathcal{DS}(\mathfrak{X})$ .

THEOREM 3.13. *If  $\mathcal{DS}(\mathfrak{X})$  is a filter of  $\mathfrak{X}$ , then  $X = \mathcal{S}(\mathfrak{X}) \oplus \mathcal{DS}(\mathfrak{X})$ .*

*Proof.* For any  $x \in X$ , let  $a \in B$  with  $x \preceq a$ . Then we have  $a \ominus x \in \mathcal{S}(\mathfrak{X})$ . Thus  $x \in \langle \mathcal{S}(\mathfrak{X}) \cup \mathcal{DS}(\mathfrak{X}) \rangle$ , and so  $x \in \langle \mathcal{S}(\mathfrak{X}) \cup \mathcal{DS}(\mathfrak{X}) \rangle$ . Therefore we have  $\mathfrak{X} = \mathcal{S}(\mathfrak{X}) \bar{\oplus} \mathcal{DS}(\mathfrak{X})$ . Also, for any  $b \in \mathcal{S}(\mathfrak{X})$  and  $p \in \mathcal{DS}(\mathfrak{X})$ , putting  $x := (p \ominus 1) \ominus b$ , we have

$$\begin{aligned}
 b \ominus x &= (p \ominus 1) \ominus (b \ominus b) = [p, 1]^2 = p \in \mathcal{DS}(\mathfrak{X}), \\
 x \ominus b &= p \ominus 1 \in \mathcal{DS}(\mathfrak{X}).
 \end{aligned}$$

Then  $x \equiv b \pmod{\mathcal{DS}(\mathfrak{X})}$ . On the other hand, we obtain

$$x \ominus 1 = [p, 1]^2 \ominus (b \ominus 1) = [p, 1]^3 = p \ominus 1.$$

Then we get  $p \ominus x \in \mathcal{S}(\mathfrak{X})$  because  $b = 1 \ominus b \preceq (p \ominus 1) \ominus (p \ominus b) = p \ominus x, b \in \mathcal{S}(\mathfrak{X})$  and  $\mathcal{S}(\mathfrak{X})$  is a filter of  $X$ . Moreover, we have

$$x \ominus p = x \ominus [p, 1]^2 = (p \ominus 1) \ominus (x \ominus 1) = 1 \in \mathcal{S}(\mathfrak{X}).$$

So,  $x \equiv p \pmod{\mathcal{S}(\mathfrak{X})}$ . Therefore we have  $X = \mathcal{S}(\mathfrak{X}) \oplus \mathcal{DS}(\mathfrak{X})$ . This completes the proof.  $\square$

#### 4. Conclusion

As we know, the primary aim of the theory of WFI-algebras is to determine the structure of all WFI-algebras. The main task of a structure theorem is to find a complete system of invariants describing the WFI-algebra up to isomorphism, or to establish some connection with other

mathematics branches. In addition, the filter theory plays an important role in studying WFI-algebras, and some interesting results have been obtained by several authors. In this paper we investigate the theory of decompositions in WFI-algebras, which is a useful tool for exploring the structure of WFI-algebras. Now we consider the subdirect sum and the direct sum of a filter family of a WFI-algebra (see Theorems 3.11 and 3.13). In the future we will discuss the direct product and the subdirect product of a WFI-algebraic family.

### References

- [1] Y. B. Jun, *Weak fuzzy implication algebras*, Adv. Stud. Contemp. Math. **7** (2003), no. 1, 41–52.
- [2] Y. B. Jun, *Perfect and concrete filters of WFI-algebras*, Bull. Korean Math. Soc. **44** (2007), no. 4, 691–699.
- [3] Y. B. Jun, *Osculatory WFI-algebras*, Commun. Korean Math. Soc. **23** (2008), no. 1, 41–47.
- [4] Y. B. Jun and C. H. Park, *Uncanny filters of WFI-algebras*, Honam Math. J. **30** (2008), no. 1, 119–126.
- [5] Y. B. Jun, C. H. Park, and E. H. Roh, *Characterizations of filters and ideals on WFI-algebras*, Honam Math. J. **28** (2006), no. 4, 471–484.
- [6] Y. B. Jun and S. Z. Song, *Simulative and mutant WFI-algebras*, Honam Math. J. **28** (2006), no. 4, 559–572.
- [7] Z. Li and C. Zheng, *Relations between fuzzy implication algebra and MV-algebra*, J. Fuzzy Math. **9** (2001), no. 1, 201–205.
- [8] W. M. Wu, *Fuzzy implication algebras*, Fuzzy Systems Math. **4** (1990), no. 1, 56–63.

\*

Department of Mathematics Education and Institute of Mathematics Education

Chinju National University of Education

Jinju 660-756, Republic of Korea

*E-mail*: ehroh@cue.ac.kr or ehroh0923@gmail.com