

## ON SOME $L_1$ -FINITE TYPE (HYPER)SURFACES IN $\mathbb{R}^{n+1}$

SEYED MOHAMMAD BAGHER KASHANI

ABSTRACT. We say that an isometric immersed hypersurface  $x : M^n \rightarrow \mathbb{R}^{n+1}$  is of  $L_k$ -finite type ( $L_k$ -f.t.) if  $x = \sum_{i=0}^p x_i$  for some positive integer  $p < \infty$ ,  $x_i : M \rightarrow \mathbb{R}^{n+1}$  is smooth and  $L_k x_i = \lambda_i x_i$ ,  $\lambda_i \in \mathbb{R}$ ,  $0 \leq i \leq p$ ,  $L_k f = \text{tr} P_k \circ \nabla^2 f$  for  $f \in C^\infty(M)$ , where  $P_k$  is the  $k$ th Newton transformation,  $\nabla^2 f$  is the Hessian of  $f$ ,  $L_k x = (L_k x^1, \dots, L_k x^{n+1})$ ,  $x = (x^1, \dots, x^{n+1})$ . In this article we study the following (hyper)surfaces in  $\mathbb{R}^{n+1}$  from the view point of  $L_1$ -finiteness type: totally umbilic ones, generalized cylinders  $S^m(r) \times \mathbb{R}^{n-m}$ , ruled surfaces in  $\mathbb{R}^{n+1}$  and some revolution surfaces in  $\mathbb{R}^3$ .

### 1. Introduction

Finite type submanifolds have been introduced in the late seventies by B. Y. Chen [2] to find the best possible estimate of the total mean curvature of a compact submanifold of a Euclidean space and to find a notion of degree for submanifolds of Euclidean space. Since then the subject has had a rapid development and so many mathematicians contribute to it, see the excellent survey of B. Y. Chen [3]. We recall that an isometrically immersed submanifold  $x : M^n \rightarrow \mathbb{R}^{n+k}$  is said to be of finite type if  $x$  has a finite decomposition as  $x = \sum_{i=0}^p x_i$ , for some positive integer  $p < +\infty$ , such that  $\Delta x_i = \lambda_i x_i$ ,  $\lambda_i \in \mathbb{R}$ ,  $0 \leq i \leq p$ , where  $\Delta$  is the Laplacian operator of  $M$ . If all  $\lambda_i$ 's are mutually different, then  $M^n$  is said to be of  $p$ -type. If in particular one of  $\lambda_i$ 's is zero, then  $M$  is said to be of null  $p$ -type.

As it is well known, the Laplacian operator of an isometrically immersed hypersurface  $M^n \subset \mathbb{R}^{n+1}$  is an (intrinsic) 2nd-order linear differential operator which arises naturally as the linearized operator of the first variation of the mean curvature for normal variation of the hypersurface. From this point of view, the Laplacian operator  $\Delta$  can be seen as the first one of  $n$  operators  $L_0 = \Delta, L_1, \dots, L_{n-1}$ , where  $L_k$  stands for the linearized operator of the first variation of the  $(k+1)$ th mean curvature (the  $(k+1)$ th elementary symmetric function in terms of the principal curvatures of  $M$ ) arising from normal variation of the hypersurface. These operators are given by  $L_k(f) = \text{tr}(P_k \circ \nabla^2 f)$  for

---

Received January 15, 2008.

2000 *Mathematics Subject Classification.* 53A05, 53B25, 53C40.

*Key words and phrases.* hypersurfaces,  $(L_1)$ -finite type.

any real smooth function  $f$  on  $M$ , where  $P_k$  denotes the  $k$ th Newton transformation associated to the 2nd fundamental form of the hypersurface and  $\nabla^2 f$  is the Hessian of  $f$ . Although in general the operators  $L_k$  are not elliptic, they still share nice properties with the Laplacian operator of  $M$ , see [1]. So, it seems natural to consider isometrically immersed hypersurfaces  $x : M^n \rightarrow \mathbb{R}^{n+1}$  in Euclidean space satisfying the condition  $x = \sum_{i=0}^p x_i$ , for some positive integer  $p$  such that  $L_k x_i = \lambda_i x_i$ ,  $\lambda_i \in \mathbb{R}$ ,  $0 \leq i \leq p$ . We call such hypersurfaces,  $L_k$ -finite type. This is an important and interesting generalization of finite type hypersurfaces, and with this paper, we begin to study such hypersurfaces. Here we consider totally umbilic hypersurfaces, generalized cylinders, ruled surfaces and some revolution hypersurfaces from the point of view of  $L_1$ -finiteness type.

We should mention that similar to Proposition 1 of [5] we have the following fact. If  $M$  is an  $L_1$ - $k$ -type surface, and if  $p(T) = \prod_{i=1}^k (T - \lambda_i)$ , then  $p(L_1)(x - x_0) = 0$ . The polynomial  $p$  is called the minimal polynomial of  $M$ .

## 2. Main result

**Proposition 2.1.** *If  $x : M^n \rightarrow \mathbb{R}^{n+1}$  is a totally umbilic isometric immersion, then either it is of  $L_k$ -null 1-type for every  $k$ ,  $0 \leq k \leq n - 1$ , or it is of  $L_k$ -1-type for  $1 \leq k \leq n - 1$ .*

*Proof.* If  $M$  is totally geodesic, then its shape operator  $S$  is  $S \equiv 0 \Rightarrow H_k = 0$  for  $0 \leq k \leq n \Rightarrow L_k x = c_k H_{k+1} N = 0$ ,  $H_k$  is the  $k$ th mean curvature of  $M$ , see [1], so  $M$  is of  $L_k$ -null 1-type for every  $k$ ,  $0 \leq k \leq n - 1$ .

If  $M$  is totally umbilic, but not totally geodesic, then its shape operator  $S$  is  $S = \lambda id$ , for some  $\lambda \in \mathbb{R}$ ,  $\lambda \neq 0$ ,  $\langle x, x \rangle = \lambda^2$ , so  $L_k x = c_k H_{k+1} N = c_k \binom{n}{k} \lambda^{k+1} N = c_k \binom{n}{k} \lambda^{k-1} x$ , see [6, Proposition 4.36]. So  $M$  is of  $L_k$ -1-type for every  $k$ ,  $1 \leq k \leq n - 1$ .  $\square$

**Proposition 2.2.** *If the isometrically immersed hypersurface  $M^n \subset \mathbb{R}^{n+1}$  is a generalized cylinder  $S^m(r) \times \mathbb{R}^{n-m}$ , then  $M$  is of  $L_k$ -null 1-type, if  $k + 1 > m$ , and it is of  $L_k$ -null 2-type, if  $k + 1 < m$ .*

*Proof.* Since  $L_k x = c_k H_{k+1} N$  and the shape operator of  $M$ ,  $S$  is

$$S = \text{diag}\left(\frac{1}{\sqrt{r}}, \dots, \frac{1}{\sqrt{r}}, 0, \dots, 0\right),$$

where the multiplicity of  $\frac{1}{\sqrt{r}}$  is  $m$ . If  $k + 1 > m \Rightarrow H_{k+1} = 0$ , so  $L_k x = 0$  hence  $M$  is of  $L_k$ -null 1-type. If  $k + 1 < m \Rightarrow H_{k+1} = \binom{m}{k+1} \frac{1}{r^{\frac{k+1}{2}}}$ ,  $N(x) = (x_1, \dots, x_{m+1}, 0, \dots, 0) \Rightarrow x = x_I + x_{II}$ ,  $x_I = (x_1, \dots, x_{m+1}) \times \{0\} \in \mathbb{R}^{n+1}$ ,  $x_{II} = \{0\} \times (x_{m+2}, \dots, x_{n+1}) \in \mathbb{R}^{n+1}$ ,  $L_k x_I = c_k \binom{m}{k+1} \frac{1}{r^{\frac{k+1}{2}}} x_I$ ,  $L_k x_{II} = 0$  hence  $M$  is of  $L_k$ -null 2-type.  $\square$

## 3. Ruled surfaces in $\mathbb{R}^{n+1}$

In this section we want to consider the ruled surfaces of  $\mathbb{R}^{n+1}$ . Here we follow [4] and give the result via Propositions 3.1 and 3.2.

**Proposition 3.1.** *If  $M$  is a cylindrical ruled surface in  $\mathbb{R}^{n+1}$ , then  $M$  is of  $L_1$ -finite type if and only if it is a cylinder over a curve of  $L_1$ -finite type.*

*Proof.* Let  $M$  be a cylinder over a curve  $\gamma$  in an affine hyperplane  $E^n$  which we can choose to have the equation  $x_{n+1} = 0$ . We can assume that  $\gamma$  is parameterized by its arc length, then a parametrization  $x$  of  $M$  is given by  $x(s, t) = \gamma(s) + t e_{n+1}$ . The induced metric on  $x$  is  $[\langle \gamma', \gamma' \rangle \ 0; \ 0 \ 1]$ ,  $L_1$  of  $M$  is of the form  $L_1 = \text{tr}(2H_1I - S) \circ H$ , where  $S$  is the shape operator of  $M$  which is of the form  $[\lambda \ 0; \ 0 \ 0]$ ,  $\lambda \in C^\infty(\mathbb{R})$ ,  $H = (H_{ij})$  is the Hessian operator. To find  $L_1$ , it is enough to know  $H_{11}, H_{22}$ . By the formula of the metric one gets that  $H_{11} = \frac{\partial^2}{\partial s^2}$ ,  $H_{22} = \frac{\partial^2}{\partial t^2}$ . By  $L'_1$  of  $\gamma$ , we mean  $L_1|_\gamma$  and  $L_1 x_{n+1} = L_1 t = 0$ . Thus as in [4], one obtains that  $M$  is of  $L_1$ -finite type if and only if each component of  $\gamma(s)$  is a finite sum of eigenfunctions of  $L_1$ , that is  $\gamma(s) = \Gamma_0 + \sum_{i=1}^p \Gamma_i(s, t)$  where  $L_1 \Gamma_i = \lambda_i \Gamma_i$ , it can be seen that  $\Gamma_i$  does not depend on  $t$  and  $(L_1|_\gamma) \Gamma_i = L_1 \Gamma_i = \lambda_i \Gamma_i$ ,  $1 \leq i \leq p$ , then  $M$  is of  $L_1$ -( $p+1$ )-type, unless one of the eigenfunctions which appear in the decomposition of  $\gamma$  has eigenvalue 0, in which case  $M$  is of  $L_1$ - $p$ -type.  $\square$

**Proposition 3.2.** *If  $M$  is a non cylindrical ruled surface in  $\mathbb{R}^{n+1}$ , then  $M$  is of  $L_1$ -null-1-type if and only if its Gaussian curvature is zero, i.e.,  $M$  is flat.*

*Proof.* If  $M$  is not cylindrical, we can decompose  $M$  into open pieces such that on each piece we can find a parametrization  $x$  of the form  $x(s, t) = \alpha(s) + t\beta(s)$ , where  $\alpha, \beta$  are curves in  $\mathbb{R}^{n+1}$  such that  $\langle \alpha', \beta \rangle = 0$ ,  $\langle \beta, \beta \rangle = 1$  and  $\langle \beta', \beta' \rangle = 1$ , i.e., the induced metric on  $x$  is  $[\ q \ 0; \ 0 \ 1]$  where  $q = t^2 + 2\langle \alpha', \beta' \rangle t + \langle \alpha', \alpha' \rangle = \|\alpha' + t\beta'\|^2$ . We have that

$$\begin{aligned} H_{ij} &= \frac{\partial^2}{\partial x_i \partial x_j} - \sum_k \Gamma_{ij}^k \frac{\partial}{\partial x_k}, \quad 1 \leq i, j, k \leq 2, \\ \frac{\partial}{\partial x_1} &= \frac{\partial}{\partial s}, \quad \frac{\partial}{\partial x_2} = \frac{\partial}{\partial t}, \\ \frac{\partial x}{\partial s} &= \alpha'(s) + t\beta'(s), \quad \frac{\partial x}{\partial t} = \beta(s), \\ \frac{\partial^2 x}{\partial s^2} &= \alpha''(s) + t\beta''(s), \quad \frac{\partial^2 x}{\partial t^2} = 0, \quad \frac{\partial^2 x}{\partial t \partial s} = \beta'(s). \end{aligned}$$

To calculate  $L_1 x$ , we need to know  $H_{11}, H_{22}$  and the shape operator of  $M$ . We obtain that

$$H_{11} = \frac{\partial^2}{\partial s^2} - \frac{1}{2} \frac{\partial q}{\partial s} \frac{\partial}{\partial s} - \frac{q}{2} \frac{\partial q}{\partial t} \frac{\partial}{\partial t}, \quad H_{22} = \frac{\partial^2}{\partial t^2}.$$

We have

$$S(x_s) = \frac{l}{m}x_s + \frac{m}{G}x_t, \quad S(x_t) = \frac{m}{E}x_s + \frac{n}{G}x_t,$$

$$l = \langle S(x_s), x_s \rangle = \langle N, x_{ss} \rangle, \quad N = \frac{x_s \times x_t}{\|x_s \times x_t\|}, \quad S(x_t) = \frac{m}{E}x_s,$$

$$x_s = \frac{\partial x}{\partial s}, \quad x_t = \frac{\partial x}{\partial t},$$

$l, m$  are polynomials in  $t$  of degree at most 2, with coefficients, functions in  $s$ ,  $E$  is a polynomial in  $t$  of degree 2, with coefficients functions in  $s$ ,  $G = \text{constant}$ , so the entries of  $S$  are rational functions in  $t$  with coefficients, functions in  $s$ . The nominators are of degree at most 2. Hence if  $P$  is a polynomial in  $t$  with coefficients, functions in  $s$  and  $\deg P = d$ , then

$$L_1(P(t)) = \frac{\widetilde{P}(t)}{q^3 Q}$$

where  $\widetilde{P}$  is a polynomial in  $t$  with functions in  $s$  as coefficients, and  $\deg \widetilde{P} \leq d + 6$ ,  $Q$  is a polynomial in  $t$  with functions as coefficients,  $\deg Q = 2$ .

If  $M$  is of  $L_1$ - $k$ -type,  $\exists c_1, \dots, c_k \in \mathbb{R}$ , s.t.,

$$(3.1) \quad L_1^{k+1}x + c_1 L_1^k x + \dots + c_k L_1 x = 0$$

We know that every component of  $x$  is a linear function in  $t$  with functions in  $s$  as coefficients. By the relation

$$L_1(P(t)) = \frac{\widetilde{P}(t)}{q^3 \cdot Q} \Rightarrow L_1^r x = \frac{P_r(t)}{q^{3r-1} \cdot Q^r},$$

where  $P_r$  is a vector field whose components are polynomial in  $t$  with functions in  $s$  as coefficients,  $\deg P_r \leq 1 + 6r$ . Hence if  $r$  goes up by one, the degree of the numerator of any component of  $L_1^r x$  goes up by at most 6, while the degree of the denominator goes up by 8. Hence the sum 3.1 can never be zero, unless of course  $L_1 x = 0$ , this means that  $H_2$ , the sectional (here the Gaussian) curvature of  $M$  is zero, (since  $\dim M = 2$ ,  $M \subseteq \mathbb{R}^{n+1}$ ). The converse is trivial.  $\square$

#### 4. Revolution hypersurfaces

In the final section we consider some of the revolution hypersurfaces. We follow [5] and study such hypersurfaces in two different cases, polynomial and rational kind revolution hypersurfaces. We begin with the first case.

**Definition 4.1.** Let  $M^2 \subseteq \mathbb{R}^3$  be a revolution hypersurface given by  $x(u, v) = ((g(u), h(u) \cos v), h(u) \sin v), (u, v) \in \mathbb{R}^2$ . Then  $M$  is called of polynomial (resp. rational) kind if  $g$  and  $h$  are polynomials in  $u$  (resp. if  $g$  is a rational function in  $h$ ).

#### 4.1. Polynomial kind revolution hypersurfaces

Let  $M^2 \subseteq \mathbb{R}^3$  be parameterized by  $x(u, v) = (g(u), h(u) \cos v, h(u) \sin v)$  where  $g, h$  are polynomials,  $(u, v) \in \mathbb{R}^2$ , we also assume that  $g'^2 + h'^2 = A^2$  where  $A$  is a polynomial in  $u$ . Then  $E = \langle x_u, x_u \rangle = g'^2 + h'^2$ ,  $G = \langle x_v, x_v \rangle = h^2$ ,  $F = \langle x_u, x_v \rangle = 0$ , so the matrix of the induced metric on  $M$  is  $\begin{bmatrix} g'^2 + h'^2 & 0 \\ 0 & h^2 \end{bmatrix}$ . If  $S$  is the shape operator of  $M$ , then  $S = \begin{bmatrix} \frac{1}{E} & 0 \\ 0 & \frac{1}{G} \end{bmatrix}$  where

$$l = \frac{-g'h'' + g''h'}{\sqrt{g'^2 + h'^2}}, \quad n = \frac{-g'h}{\sqrt{g'^2 + h'^2}}.$$

In order to find  $L_1 f = \text{tr}(2H_1 I - S) \circ H^f$ ,  $f \in C^\infty(M)$ , it is enough to calculate  $H_{11}^f$  and  $H_{22}^f$ , we obtain that

$$H_{11} = \frac{\partial^2}{\partial u^2} - \frac{g'g'' + h'h''}{g'^2 + h'^2} \frac{\partial}{\partial u}, \quad H_{22} = \frac{\partial^2}{\partial v^2}.$$

Following [5], we prove the next lemma.

**Lemma 4.2.** *Let  $F(u), G(u)$  be polynomial functions in  $u$  and  $M$  as introduced in the subsection (4.1). Then  $L_1(\frac{F}{G}) = \frac{F_1}{G_1}$  for some polynomial functions  $F_1, G_1$  with*

$$d(F_1) - d(G_1) \leq d(F) - d(G) - d(h) - 2$$

*Proof.* By applying  $L_1$  and straightforward computation, we obtain that  $L_1(\frac{F(u)}{G(u)}) = \frac{F_1(u)}{G_1(u)}$ , where

$$\begin{aligned} F_1 &= g'[(g'^2 + h'^2)(F'G - FG')'G - 2(g'^2 + h'^2)G'(F'G - FG') \\ &\quad - G(g'g'' + h'h'')(F'G - FG')], \\ G_1 &= G^3 h(g'^2 + h'^2)^{\frac{3}{2}} = G^3 h A^3. \end{aligned}$$

Then we get that

$$\begin{aligned} d(G_1) &= 3d(G) + d(h) + 3 \max\{d(g), d(h)\} - 3 \\ d(F_1) &\leq 2d(G) + d(F) + 3 \max\{d(g), d(h)\} - 5 \\ \Rightarrow d(F_1) - d(G_1) &\leq d(F) - d(G) - d(h) - 2. \end{aligned} \quad \square$$

**Theorem 4.3.** *Let  $M$  be a surface of revolution of the polynomial kind as introduced in the subsection (4.1). Then  $M$  is a surface of  $L_1$ -finite type if and only if  $M$  is either an open piece of a plane or an open piece of a circular cylinder, or it is flat.*

*Proof.* Let

$$(4.1) \quad L_1^i g = \frac{F_i}{G_i}, \quad i = 1, 2, 3, \dots,$$

then similar to (3.5) of [5], we have

$$(4.2) \quad d(F_{i+1}) - d(G_{i+1}) < d(F_i) - d(G_i) < \dots < d(F_1) - d(G_1) < d(g).$$

Assume that  $M$  of  $L_1$ -finite type, say of  $L_1$ - $k$ -type. Let

$$(4.3) \quad p(T) = T^k + c_1 T^{k-1} + \cdots + c_{k-1} T + c_k$$

be the minimal polynomial of  $M$ , then  $p$  has  $k$  distinct real roots, and from the relations 4.1 and 4.3 we have that

$$\frac{F_k}{G_k} + c_1 \frac{F_{k-1}}{G_{k-1}} + \cdots + c_{k-1} \frac{F_1}{G_1} + c_k(g - a) = 0$$

for some constant  $a$ . Let  $K = G_1 \cdots G_k$ , then

$$K \frac{F_k}{G_k} + c_1 K \frac{F_{k-1}}{G_{k-1}} + \cdots + c_{k-1} K \frac{F_1}{G_1} + c_k K(g - a) = 0.$$

If  $d(g) = 0$ ,  $M$  is an open portion of a plane. If  $d(g) > 0$ , then by (4.2) we get the inequalities

$$d(K(g - a)) > d(K \frac{F_1}{G_1}) > \cdots > d(K \frac{F_{k-1}}{G_{k-1}}) > d(K \frac{F_k}{G_k})$$

which is impossible unless

$$p(T) = T^2 + c_1 T, c_1 \neq 0, L_1 g = 0,$$

since the minimal polynomial  $p$  has  $k$  distinct real roots. That is

$$L_1^2 x + c_1 L_1 x = 0,$$

so we get that  $M$  is either of  $L_1$ -1-type or of  $L_1$ -null 2-type. If  $M$  is of  $L_1$ -1-type, then  $L_1 x = c_1 H_2 N = \lambda x$  for some  $\lambda \in \mathbb{R}$ , if  $\lambda = 0 \Rightarrow H_2 = 0$ , so  $M$  is flat. If  $\lambda \neq 0$ , then  $M$  has to be an open portion of an sphere, which is not possible, since  $M$  is a revolution hypersurface of polynomial kind in  $\mathbb{R}^3$ . If  $M$  is of  $L_1$ -null 2-type, we characterize it as follows. We have that  $x = c + x_p + x_q$  where  $c \in \mathbb{R}^3$ ,  $x_p, x_q : M \rightarrow \mathbb{R}^3$  are smooth,  $L_1 x_p = 0$ ,

$$\begin{aligned} L_1 x_q &= \lambda_q x_q, \lambda_q \in \mathbb{R}, \lambda_q \neq 0 \Rightarrow L_1 x_q = \lambda_q x_q = L_1 x = c_1 H_2 N \\ &\Rightarrow x_q = \frac{c_1 H_2 N}{\lambda_q} \\ (4.4) \quad &\Rightarrow L_1 \left( \frac{c_1 H_2 N}{\lambda_q} \right) = \frac{c_1}{\lambda_q} L_1(H_2 N) \lambda_q x_q = c_1 H_2 N. \end{aligned}$$

Now we calculate  $L_1(H_2 N)$  directly.

$$\begin{aligned} X \cdot \langle H_2 N, a \rangle &= \langle (X \cdot H_2) N, a \rangle - H_2 \langle S X, a \rangle \\ (4.5) \quad &\Rightarrow \nabla \langle H_2 N, a \rangle = \langle N, a \rangle \nabla H_2 - S(a^T) \\ &\Rightarrow \nabla_X (\nabla \langle H_2 N, a \rangle) \end{aligned}$$

$$\begin{aligned}
&= X \cdot \langle N, a \rangle \nabla H_2 + \langle N, a \rangle \nabla_X \nabla H_2 - \nabla_X (S(a^T)) \\
&= -\langle SX, a \rangle \nabla H_2 + \langle N, a \rangle \nabla_X \nabla H_2 - \nabla S(a^T, X) - S(\nabla_X a^T) \\
&= -\langle S(a^T), X \rangle \nabla H_2 + \langle N, a \rangle \nabla_X \nabla H_2 - (\nabla_{a^T} S)X - \langle N, a \rangle S^2 X \\
\Rightarrow L_1 \langle H_2 N, a \rangle &= \text{tr}(p_1 \circ \nabla_{a^T} S) - \langle N, a \rangle \text{tr} S^2 \circ P_1 - \sum_i \langle \langle S a^T, e_i \rangle \nabla H_2, P_1 e_i \rangle \\
&\quad + \langle N, a \rangle \sum_i \langle P_1(\nabla_{e_i} \nabla H_2), e_i \rangle \\
\Rightarrow L_1 \langle H_2 N, a \rangle &= L_1 \langle N, a \rangle + \langle N, a \rangle \sum_i H_{H_2}(e_i, P_1 e_i) - \langle S \circ P_1(\nabla H_2), a \rangle \\
\Rightarrow L_1(H_2 N) &= L_1 N + \left( \sum_i H_{H_2}(e_i, P_1 e_i) \right) N - S \circ P_1(\nabla H_2)
\end{aligned}$$

using the formula of  $L_1 N$  and comparing (4.5), (4.4) one gets that  $S \circ P_1(\nabla H_2) = -\nabla H_2$ . By using the relation  $P_1 = 2H_1 I - S$  and that  $S = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$  with respect to the ordered basis  $\{x_u, x_v\}$  one obtains that the relation  $S \circ P_1(\nabla H_2) = -\nabla H_2$  (by counting the degrees of the polynomials involved in this equation) holds only when  $h$  equals some constant  $\lambda$ ,  $\lambda \in \mathbb{R}, \lambda \neq 0$ , i.e.,  $x(u, v) = (g(u), \lambda \cos v, \lambda \sin v)$ ,  $\lambda \in \mathbb{R}, \lambda \neq 0$ , so  $M$  is a circular cylinder. We leave the proof of the converse to the reader as an easy exercise.  $\square$

#### 4.2. Revolution hypersurfaces of rational kind

Let  $M$  be parameterized by  $x(u, v) = (g(u), h(u) \cos v, h(u) \sin v)$ , we recall that  $M$  is said to be of rational kind if  $g$  is a rational function in  $h$ , i.e.,  $g$  is the quotient of two polynomial functions in  $h$ .

Following [5], we get the final result.

**Theorem 4.4.** *Let  $M$  be a surface of revolution of the rational kind such that if  $g' = \frac{Q}{R}$ ,  $(Q, R) = 1$ , then  $Q^2 + R^2 = A^2$ , where  $A$  is a polynomial. Then  $M$  is a surface of  $L_1$ -finite type if and only if  $M$  is flat.*

In fact Theorem 4.4 follows from the following more general result.

**Theorem 4.5.** *Let  $M$  be an  $L_1$ -finite type surface of revolution parameterized by  $x(t, v) = (g(t), t \cos v, t \sin v)$ , where  $g$  is a rational function in  $t$  and if  $g'(t) = \frac{Q}{R}$ ,  $(Q, R) = 1$ , then  $Q^2 + R^2 = A^2$ , where  $A$  is a polynomial in  $t$ . Then  $M$  is flat or  $\deg Q = \deg R = 2 + \deg(Q + R)$ .*

*Proof of Theorem 4.4.* Without loss of generality, we may assume that  $M$  is parameterized by  $x(t, v) = (g(t), t \cos v, t \sin v)$ , where  $g(t) = \frac{G(t)}{H(t)}$  for some polynomial functions  $G, H$ ,  $(G, H) = 1$ . Assume that  $M$  is of  $L_1$ -finite type, then by Theorem 4.5,  $M$  is flat. The converse is trivial.  $\square$

*Proof of Theorem 4.5.* Let  $M$  be a surface of revolution parameterized by  $x(t, v) = (g(t), t \cos v, t \sin v)$  as in Theorem 4.5. In order to find  $L_1$  we need

the metric, the shape operator  $S$  and  $H_{11}, H_{22}$ . The induced metric on  $x$  is  $[\begin{smallmatrix} g'^2+1 & 0 \\ 0 & t^2 \end{smallmatrix}]$ , the shape operator  $S$  is

$$\begin{bmatrix} \frac{g''}{(g'^2+1)^{\frac{3}{2}}} & 0 \\ 0 & \frac{g'}{t(g'^2+1)^{\frac{1}{2}}} \end{bmatrix}.$$

We obtain that

$$H_{11} = \frac{\partial^2}{\partial t^2} - \frac{g'g''}{g'^2+1} \frac{\partial}{\partial t}, \quad H_{22} = \frac{\partial}{\partial v^2}.$$

So,

$$\begin{aligned} L_1 &= \text{tr}(s_1 H_1 I - S) \circ H, \quad H = \begin{bmatrix} H_{11} & * \\ * & H_{22} \end{bmatrix} \\ \Rightarrow L_1 &= \text{tr} \begin{bmatrix} \frac{g'}{t(g'^2+1)^{\frac{1}{2}}} & 0 \\ 0 & \frac{g''}{(g'^2+1)^{\frac{3}{2}}} \end{bmatrix} \circ \begin{bmatrix} \frac{\partial^2}{\partial t^2} - \frac{g'g''}{g'^2+1} \frac{\partial}{\partial t} & * \\ * & \frac{\partial}{\partial v^2} \end{bmatrix} \\ \Rightarrow L_1 &= \frac{g'}{t(g'^2+1)^{\frac{1}{2}}} \left( \frac{\partial^2}{\partial t^2} - \frac{g'g''}{g'^2+1} \frac{\partial}{\partial t} \right) + \frac{g''}{(g'^2+1)^{\frac{3}{2}}} \frac{\partial^2}{\partial v^2} \end{aligned}$$

then if  $L_1(t \cos v) = \frac{Q_1}{R_1} \cos v \Rightarrow \frac{Q_1}{R_1} = \frac{-g'^2 g'' - t^2 g''}{t(g'^2+1)^{\frac{3}{2}}} = -\frac{g''(g'^2+t^2)}{t(g'^2+1)^{\frac{3}{2}}}$  if  $g' = \frac{Q}{R} \Rightarrow \frac{Q_1}{R_1} = \frac{(QR' - Q'R)(Q^2 + t^2 R^2)}{tR(Q^2 + R^2)^{\frac{3}{2}}}$ ,  $Q^2 + R^2 = A^2$ . If inductively  $L_1^{i+1}(t \cos v) = \frac{Q_{i+1}}{R_{i+1}} \cos v$ ,  $i = 1, 2, 3, \dots$  for some polynomial functions  $Q_i, R_i$ , put  $\bar{Q}_i = \bar{Q}_i R_i - Q_i R_i'$ ,  $\bar{Q}_i = (Q_i' R_i - Q_i R_i') R_i^2 - 2R_i R_i' (Q_i' R_i - Q_i R_i')$ , then have

$$\begin{aligned} \frac{Q_{i+1}}{R_{i+1}} &= \frac{g'}{t(g'^2+1)^{\frac{1}{2}}} \frac{\bar{Q}_i}{R_i^4} - \frac{g'^2 g''}{t(g'^2+1)^{\frac{3}{2}}} \frac{\bar{Q}_i}{R_i^2} - \frac{g''}{(g'^2+1)^{\frac{3}{2}}} \frac{Q_i}{R_i} \\ &= \frac{Q}{t(Q^2+R^2)^{\frac{1}{2}}} \times \frac{\bar{Q}_i}{R_i^4} - \frac{Q^2(Q'R - R'Q)}{tR(Q^2+R^2)^{\frac{3}{2}}} \times \frac{\bar{Q}_i}{R_i^2} - \frac{R(Q'R - R'Q)}{(Q^2+R^2)^{\frac{3}{2}}} \frac{Q_i}{R_i} \\ &= \frac{QR(Q^2+R^2)\bar{Q}_i - Q^2 R_i^2 (Q'R - R'Q)\bar{Q}_i - tR^2 R_i^3 (Q'R - R'Q)Q_i}{tRR_i^4(Q^2+R^2)^{\frac{3}{2}}} \end{aligned}$$

$$\begin{aligned} \Rightarrow Q_{i+1} &= QR(Q^2+R^2)\bar{Q}_i - Q_i^2 R_i^2 (Q'R - R'Q)\bar{Q}_i - tR^2 R_i^3 (Q'R - R'Q)Q_i, \\ R_{i+1} &= tRR_i^4(Q^2+R^2)^{\frac{3}{2}} = tRR_i^4 A^3. \end{aligned}$$

Assume that  $M$  is of  $L_1$ -finite type, say of  $k$ -type, let  $p(T) = T^k + c_1 T^{k-1} + \dots + c_{k-1} T + c_k$  be the minimal polynomial of  $M$ , then  $p$  has  $k$  distinct real roots. We have that

$$\frac{Q_k}{R_k} + c_1 \frac{Q_{k-1}}{R_{k-1}} + \dots + c_{k-1} \frac{Q_1}{R_1} + c_k t = 0.$$



Let  $K = R_1 \cdots R_k$ , then we get

$$K \frac{Q_k}{R_k} + c_1 K \frac{Q_{k-1}}{R_{k-1}} + \cdots + c_{k-1} K \frac{Q_1}{R_1} + c_k t K = 0.$$

Following ([5], cases i-iii of Theorem 3) by similar discussion about  $\deg(Q)$ ,  $\deg(R)$  according to the cases  $\deg(Q) > \deg(R)$  or  $\deg(Q) < \deg(R)$  or  $\deg(Q) = \deg(R)$ , we see that the only possibility is that either  $p(T) = T$  or  $\deg(Q) = \deg(R) = 2 + \deg(Q + R)$ . When  $p(T) = T \Rightarrow H_2 = 0$ , in this case,  $M$  is flat.  $\square$

**Acknowledgement.** The author would like to thank Tarbiat Modares university for its financial support and Warwick university for its hospitality while he was spending his sabbatical leave, during which this work has been done. He should thank gratefully L. J. Alias for introducing him the problem.

### References

- [1] L. J. Alias and N. Gürbüz, *An extension of Takahashi theorem for the linearized operators of the higher order mean curvatures*, *Geom. Dedicata* **121** (2006), 113–127.
- [2] B. Y. Chen, *Total Mean Curvature and Submanifolds of Finite Type*, Series in Pure Mathematics, 1. World Scientific Publishing Co., Singapore, 1984.
- [3] ———, *A report on submanifolds of finite type*, *Soochow J. Math.* **22** (1996), no. 2, 117–337.
- [4] B. Y. Chen, F. Dillen, L. Verstraelen, and L. Vrancken, *Ruled surfaces of finite type*, *Bull. Austral. Math. Soc.* **42** (1990), no. 3, 447–453.
- [5] B. Y. Chen and S. Ishikawa, *On classification of some surfaces of revolution of finite type*, *Tsukuba J. Math.* **17** (1993), no. 1, 287–298.
- [6] B. O'Neill, *Semi-Riemannian Geometry*, Pure and Applied Mathematics, 103. Academic Press, Inc., New York, 1983.

DEPARTMENT OF MATHEMATICS  
 FACULTY OF SCIENCES  
 TARBIAT MODARES UNIVERSITY  
 TEHRAN 14115-175, IRAN  
*E-mail address:* kashanim@modares.ac.ir