

Erratum

J. Microbiol. Biotechnol. (2008) 18(10), 1655–1658

Isolation and Structure Determination of a Proteasome Inhibitory Metabolite from a Culture of *Scytonema hofmanni*

Shim, Sang Hee¹, George Chlipala², and Jimmy Orjala^{2*}

¹*School of Biotechnology, Yeungnam University, Gyeongsan 712-749, Korea*

²*Department of Medicinal Chemistry and Pharmacognosy, University of Illinois at Chicago, Chicago, Illinois 60612, USA*

1. The second sentence of the second paragraph on page 1656 should appear as shown below.
As part of a project directed toward the discovery of anticancer agents from cyanobacteria, *Scytonema hofmanni* was obtained from the culture collection of Algae at the University of Texas at Austin (UTEX) as strain 1834.
2. The first sentence of the third paragraph on page 1656 should appear as shown below.
Scytonema hofmanni was cultured in three 1 l Elenmyer flasks, each containing 350 ml of media BG-11, under continuous illumination with fluorescent lamps at $29 \mu\text{mol m}^{-2}\text{s}^{-1}$.
3. Acknowledgement should appear as shown below.
This research was supported by the Yeungnam University research grants in 2007 and NIH grant R01GM075856. We thank Shunyan Mo, UIC, for her assistance.