

Co-expression of CdtA and CdtC subunits of cytolethal distending toxin from *Aggregatibacter actinomycetemcomitans*

Seung-Jae Lee¹, Kyung-Yeol Lee², Hyung-Seop Kim^{1*}

1. Department of Periodontology, College of Dentistry, Chonbuk National University

2. Department of Oral Microbiology and Institute of Oral Bioscience, Brain Korea 21 Project, Chonbuk National University

ABSTRACT

Purpose: Cytolethal distending toxin (CDT) is a family of heat-labile cytotoxins produced by several gram-negative mucosa-associated pathogens, including *Aggregatibacter actinomycetemcomitans*. CDT is well known to be capable of inducing growth arrest, morphological alterations, and eventually death in various cells. CDT belongs to a tripartite AB₂ toxin (CdtB: the enzymatic A subunit; CdtA and CdtC: the heterodimeric B subunit). Previous studies proposed that CdtA and CdtC together bind to a cell surface receptor and glycolipids act as a receptor for *A. actinomycetemcomitans* CDT (AaCDT). In this study, recombinant CdtA and CdtC proteins of AaCDT were co-expressed in a bacterial expression system and tested for their affinity for GM₁ ganglioside.

Methods: The genes for CdtA and CdtC from *A. actinomycetemcomitans* Y4 were utilized to construct the expression vectors, pRSET-cdtA and pET28a-cdtC. Both CdtA and CdtC proteins were expressed in *Escherichia coli* BL21(DE3) and then purified using hexahistidine (His6) tag. The identity of purified protein was confirmed by anti-His6 antibody and monoclonal anti-CdtA antibody. Furthermore, the affinity of recombinant protein to GM₁ ganglioside was checked through ELISA.

Results: Recombinant CdtA and CdtC proteins were expressed as soluble proteins and reacted to anti-His6 and monoclonal anti-CdtA antibodies. ELISA revealed that purified soluble CdtA-CdtC protein bound to GM₁ ganglioside, while CdtA alone did not.

Conclusions: Co-expression of CdtA and CdtC proteins enhanced the solubility of the proteins in *E. coli*, leading to convenient preparation of active CdtA-CdtC, a critical material for the study of AaCDT pathogenesis.

(*J Korean Acad Periodontol* 2009;39:231-237)

KEY WORDS: *Aggregatibacter actinomycetemcomitans*; toxin.

INTRODUCTION

Periodontitis is an inflammatory disease in the supporting tissues of the teeth caused by a composite of microorganisms. *Aggregatibacter actinomycetemcomitans* has been suspected to be one of the key pathogens in the etiology of human periodontitis.

A. actinomycetemcomitans is a gram-negative coccobacillus has been implicated not only in the patho-

genesis of localized aggressive periodontitis but also in several systemic diseases, such as endocarditis, meningitis, and osteomyelitis. This organism is able to produce a variety of virulence factors that are involved in the colonization of the oral cavity, the destruction and inhibition of regeneration of the periodontal tissues, and interference with host defense mechanisms. These virulence factors include collagenases, endotoxin, leukotoxin, and cytolethal distending toxin (CDT)¹.

CDTs are a family of heat-labile proteinous cytotoxins produced by several gram-negative pathogens associated with mucosal surfaces, including *Escherichia coli*, *Campylobacter jejuni*, *Shigella species*, *Haemophilus ducreyi*, and *A. actinomycetemcomitans*, with the capacity to induce growth arrest, morphological alter-

Correspondence: Dr. Hyung-Seop Kim

Department of Periodontology, College of Dentistry, Chonbuk National University, 634-18, Gumam-dong, Dukjin-gu, Jeonju, 512-712, Korea,

E-mail: cbuperio@chonbuk.ac.kr, Tel: 82-63-250-2116, Fax: 82-63-250-2259. Received: Jun 25, 2009; Accepted: Jul 28, 2009

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2008-331-E00377).

ations, and eventually death in the affected cells²⁾.

CDT is found and expressed in the majority of the *A. actinomycetemcomitans* isolates. Leung et al.³⁾ found that *A. actinomycetemcomitans* was isolated from 67% of aggressive periodontal diseased subjects and all but one contained all three cdt genes. Ahmed et al.⁴⁾ reported that 43 of 50 strains of *A. actinomycetemcomitans* isolated from periodontitis patients possessed all three cdt genes and expressed a cytotoxic activity. Also, Fabris et al.⁵⁾ found that 34 of 40 *A. actinomycetemcomitans* detected all three cdt genes and 39 samples caused distension of Chinese hamster ovary cells.

The periodontium is composed of epithelial cells, fibroblasts, cementoblasts, and osteoblasts. These various cell types respond differently to the CDT of *A. actinomycetemcomitans*. Belibasakis et al.⁶⁾ reported CDT of *A. actinomycetemcomitans* inhibits the proliferation of gingival fibroblasts and periodontal ligament cells without affecting their viability, and the cell cycle arrest associated with this inhibition is a combined G1 and G2/M phase arrest. In contrast, Kanno et al.⁷⁾ suggested that human periodontal ligament fibroblasts are resistant to the cytotoxic effects of the *A. actinomycetemcomitans* CDT.

Proliferation of human epithelial cell lines was rapidly inhibited by CDT^{7,8)}. By inducing growth arrest in oral epithelial cells, the toxin may interfere with the normal periodontal connective tissue remodeling equilibrium⁶⁾, or may facilitate the invasion and multiplication of *A. actinomycetemcomitans* in the deeper periodontal tissue⁸⁾, respectively.

A. actinomycetemcomitans up-regulates receptor activator of NF- κ B ligand (RANFL) expression in gingival fibroblasts and periodontal ligament cells, and that CDT is its component primarily responsible for this event⁹⁾. CDT can also stimulate RANKL expression in T-cells¹⁰⁾. RANKL expression activates osteoclasts involved in bone resorption. RANKL expression in the periodontium has been implicated in physiological tooth eruption and orthodontic tooth movement, as

well as in the pathological process of periodontitis¹¹⁾.

CDT toxin is encoded by three different genes: *cdtA*, *cdtB*, and *cdtC*, and expression of all three *cdt* genes is required in order to produce active CDT¹²⁾. CDT is suggested to be a unique tripartite AB₂ toxin in which CdtB is the active A subunit, and CdtA and CdtC constitute the heterodimeric B subunit^{2,13,14)}. Furthermore, it is generally agreed that the active subunit, CdtB, enters the cell while CdtA and CdtC remain associated with the cell surface¹⁴⁾. The CdtB protein has been shown to possess DNase activity¹³⁾ when introduced into or expressed within eukaryotic cells, and there seems to be a reasonable consensus that a functional CdtB molecule is essential for expression of toxicity by CDT²⁾. The cellular responses to DNA damage lead to a characteristic G₂/M cell cycle arrest, cellular distension, nuclear enlargement observed in intoxicated cells¹⁵⁾.

While progress in understanding the function of CdtB has been made, there is still little known about the precise roles of CdtA and CdtC in CDT function. Mao and DiRienzo⁸⁾ demonstrated by immunofluorescence assay that only CdtA gene derived from *A. actinomycetemcomitans* binds to the surfaces of Chinese hamster ovary cells independently. A study by Lee et al.¹⁶⁾ with *C. jejuni* showed both CdtA and CdtC were able to bind specifically and independently to the surface of HeLa cells. But, many studies find that CdtA and CdtC together bind to the cellular membrane^{14,17,18)}. In the crystal structures of the *A. actinomycetemcomitans* holotoxins, CdtA and CdtC form ricin-like lectin domains that may play a role in recognizing the cellular receptor that delivers the CdtB subunit into the target cells^{14,19)}. The cell surface receptor for the CDT has not yet been identified. However, there is evidence that *E. coli* CdtA-II and CdtC-II bind N-linked fucose-containing complex carbohydrates on the surfaces of HeLa cells¹⁷⁾. Other studies reported that glycosphingolipid GM₁ and GM₃ may play an important role as the receptor of *A. actinomycetemcomitans* CDT¹⁸⁾. In addition, cholesterol

ol-rich cytoplasmic membrane lipid rafts are also involved in the delivery of CdtB into target cell²⁰.

The goal of this study was to co-express CdtA and CdtC subunits of CDT from *A. actinomycetemcomitans* and to find whether purified CdtA-CdtC protein binds to GM₁ ganglioside.

MATERIALS AND METHODS

1. Bacterial strains and culture condition

A. actinomycetemcomitans Y4 (ATCC 43718) was cultured in Brain Heart Infusion broth medium for 24 h at 37°C in anaerobic chamber containing 85% N₂, 10% H₂, and 5% CO₂²¹. Growth was monitored by optical density (OD) at 600 nm, with all batch cultures having an initial OD₆₀₀ of approximately 0.01 after sub-culturing from overnight growth. For the co-expression of CdtA and CdtC subunit, *E. coli* BL21(DE3) transformed with pRSET-cdtA²² and pET28a-cdtC²³ was cultured in Luria-Bertari (LB) medium at 37°C with shaking at 200 rpm.

2. Co-expression of CdtA and CdtC subunits

E. coli BL21(DE3) transformed with the pRSET-cdtA and pET28a-cdtC plasmid was cultured into 5 ml of LB broth containing 50 ug/ml for each of ampicillin and kanamycine. At the following day, the bacteria were freshly inoculated into the 500 ml LB broth containing 50 ug/ml ampicillin/kanamycine and further grown at 37°C with vigorously shaking until OD₆₀₀ reached to 0.8 and then induced by 1 mM of isopropyl-β-D-thiogalactopyranoside (IPTG) for another 3 h at 30°C

3. Purification of CdtA-CdtC protein

Induced bacteria were harvested by centrifugation,

and the resulting pellet was resuspended in binding buffer (20 mM Tris-HCl pH 8.0, 0.5 M NaCl, and 5 mM imidazole). The resuspended bacteria were disrupted with an ultrasonic disruptor²⁴. The sonicated samples were centrifuged at 13000 rpm for 10 min and then the supernatant was saved as the soluble fractions. The soluble fractions were then loaded into the nickel-chelated agarose (Ni-NTA, Qiagen). The column was washed with binding buffer and eluted with binding buffer containing 200 mM imidazole.

4. SDS-PAGE and western blot analysis of purified CdtA-CdtC protein

The fractions containing recombinant protein (His6-tagged protein) were dialyzed and then resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Recombinant CdtA-CdtC protein were mixed with 100 μl of gel loading buffer and heated in a boiling water bath for 5 min. The samples were applied to 10-20% polyacrylamide gel. After electrophoresis, protein bands were transferred to a nitrocellulose membrane. The membrane was blocked with 3% BSA in TBS containing 0.1% Tween-20 (TBST) buffer for 1h at room temperature with shaking. The blocked membrane was incubated with anti-His6 antibody or monoclonal anti-CdtA antibody tagged with AP (alkaline phosphatase) and then washed in TBST. The membrane was treated with BCIP/NBT to induce developing reaction and incubated in the dark. The membrane was rinsed with H₂O to stop the developing reaction.

5. GM₁ ELISA of purified CdtA-CdtC protein

GM₁ ganglioside was used to coat Enzyme-Linked ImmunoSorbent Assay (ELISA) plate at 37°C for 2 h and the coated plates was blocked by 1% skim milk. After washed using PBS (pH 7.5), the plates were incubated with purified CdtA-CdtC protein or CdtA in PBS at 37°C for 2 h. Then the plates were washed us-

ing PBS and incubated with anti-His6 antibody or monoclonal anti-CdtA antibody in PBS at 37°C for 2 h. The plates were then washed with PBS, and further incubated with anti-mouse IgG conjugated with alkaline phosphatase in PBS (1:5000 dilution) at 37°C for 2 h and then applied with the substrate which is converted by the enzyme to elicit a chromogenic signal. The result was quantified using a spectrophotometer for OD at 405 nm.

RESULTS

1. Purification of CdtA-CdtC protein

CdtA-CdtC protein was expressed in *E. coli* BL21(DE3) transformd with pRSET-cdtA and pET28a-cdtC, and

then purified by Ni-NTA affinity column. Recombinant CdtA and CdtC proteins were clearly shown in the expected locations of SDS-PAGE gel (Fig. 1).

Both CdtA and CdtC proteins were expressed as His6-tagged fusion proteins which were confirmed by anti-His6 antibody by Western blot. CdtA was shown just below 25 kDa and CdtC around 20 kDa (Fig. 2).

Monoclonal anti-CdtA antibody also only detected CdtA at 25kDa (Fig. 3). A protein at a higher band around 35 kDa was also purified by Ni-NTA column and responded to anti-His6 and anti-CdtA antibodies.

2. GM₁ ELISA of purified CdtA-CdtC protein

By using ELISA, purified CdtA-CdtC protein was evaluated whether it bound to GM₁ ganglioside. The




Figure 1. SDS-PAGE analysis of purified CdtA and CdtC protein. After expression of recombinant proteins from pRSET-cdtA and pET28a-cdtC, the transformed bacteria were sonicated and centrifuged. The supernatant was loaded onto Ni-NTA agarose column and Flow-through (FT) was saved. The sample-loaded column was washed three times (W1~W3) and eluted (E1~E3) as described in Materials and Methods. rCdtA indicated purified recombinant CdtA protein from pET-cdtA and served as control.




Figure 2. Western blot analysis of purified CdtA and CdtC protein using anti-His6 antibody. rCdtA-CdtC protein was purified from *E. coli* transformed with both pRSET-cdtA and pET28a-cdtC, while rCdtA from *E. coli* transformed with pET-cdtA. Protein samples were electrophoresed in SDS-PAGE gel, transferred to nitrocellulose membrane, and detected by anti-His6 antibody.




Figure 3. Western blot analysis of purified CdtA and CdtC protein using monoclonal anti-CdtA antibody. Protein samples were purified by Ni-NTA column from bacterial lysates from *E. coli* transformed with both pRSET-cdtA and pET28a-cdtC (rCdtA-CdtC), and *E. coli* transformed with pET-cdtA (rCdtA). Samples were separated in SDS-PAGE gel and then transferred to nitrocellulose membrane, and detected by anti-CdtA antibody. Bands around 25 kDa was clearly shown but band at 20 kDa not shown.




Figure 4. GM₁ ELISA of purified CdtA–CdtC protein using anti-His6 antibody. ELISA plate was coated with GM₁ ganglioside and further incubated with diverse concentration of CdtA and CdtA–CdtC protein. After washing, the remaining amount of CdtA and CdtA–CdtC protein was quantified using anti-His6 antibody at OD₄₀₅.




Figure 5. GM₁ ELISA of purified CdtA–CdtC protein using monoclonal anti-CdtA antibody. ELISA plate was coated with GM₁ ganglioside and further incubated with diverse concentration of CdtA and CdtA–CdtC protein. After washing, the remaining amount of CdtA and CdtA–CdtC protein was quantified using monoclonal anti-CdtA antibody at OD₄₀₅.

peak OD₄₀₅ value of purified CdtA–CdtC protein was measured around 0.5. CdtA showed to have lower OD₄₀₅ value than CdtA–CdtC protein at any dilution (Fig. 4).

Monoclonal anti-CdtA antibody was also applied to check GM₁ affinity of purified CdtA–CdtC protein and CdtA. Recombinant CdtA–CdtC protein showed a higher OD₄₀₅ value than recombinant CdtA protein (Fig. 5).

DISCUSSION

The crystal structures of the CDT cytotoxin from *Aggregatibacter actinomycetemcomitans* include a pronounced groove between the CdtA and CdtC subunits in the heterotoxin complex¹⁹. A CDT groove mutant exhibited diminished binding to cells²⁵. This data support the hypothesis that the groove formed between CdtA and CdtC are involved in the binding of the toxin to target cells. Also, many studies reported that CdtA and CdtC together bind to the cellular membrane^{14,17,18}.

In this study, CdtA and CdtC subunits were ex-

pressed as soluble proteins from *E. coli* transformed with both cdtA- and cdtC-expressing vectors, while each of CDT subunit expressed as an insoluble protein. These expressed CDT subunits were purified with Ni-NTA column and also responded to anti-His6 and anti-CdtA antibodies (Fig. 1–3). CdtA protein was located at two different bands in SDS-PAGE and Western blot (Fig. 1–3) and some studies also report similar results^{26,27}. They reported that it was caused by a covalent bond between cysteine residue of CdtA N-terminus and lipid²⁷. Therefore, both of doublet bands in Fig. 1–3 are supposed to contain CdtA subunit.

The cell surface receptor for CDT has not been clearly identified. However, it has been reported that both CdtA and CdtC recognize N-linked fucose-containing glycoproteins¹⁷ or gangliosides, such as GM₁ and GM₃¹⁸, in vitro. Therefore purified CdtA–CdtC protein was evaluated whether it bound to GM₁ ganglioside through ELISA. Recombinant CdtA–CdtC protein had higher affinity to GM₁ ganglioside than CdtA (Fig. 4 and 5). Either anti-His6 antibody or monoclonal anti-CdtA antibody showed similar results.

The exact role of CdtA/CdtC heterodimer in CDT toxicity is not yet clearly understood, and further work is required to fully identify the role of CdtA/CdtC heterodimer, especially in its mechanism of host cell binding. Therefore, the soluble CdtA-CdtC protein will serve to investigate the pathological process of *A. actinomycetemcomitans* and also to be a potential vaccine material. More interestingly, there is a possibility of using CDT as a mucosal adjuvant through its binding activity to cell surface receptor^{27,28}. Up to now, cholera toxin or enterotoxigenic *E. coli* toxin has been mainly utilized as a mucosal adjuvant but these toxins belong to AB₅ family. But CDT belongs to AB₂ and its simple structure and/or lesser toxicity fit to develop a new mucosal adjuvant.

In conclusion, CdtA and CdtC proteins were co-expressed as soluble protein and CdtA-CdtC protein bound to GM₁ ganglioside. Further studies will be necessary to characterize precisely CdtA-CdtC protein.

REFERENCES

1. Henderson B, Nair SP, Ward JM, Wilson M. Molecular pathogenicity of the oral opportunistic pathogen *Actinobacillus actinomycetemcomitans*. *Annu Rev Microbiol* 2003;57:29-55.
2. Lara-Tejero M, Galan JE. Cytolethal distending toxin: limited damage as a strategy to modulate cellular functions. *Trends Microbiol* 2002;10:147-152.
3. Leung WK, Ngai VKS, Yau JYY et al. Characterization of *Actinobacillus actinomycetemcomitans* isolated from young Chinese aggressive periodontitis patients. *J Periodontol Res* 2005;40:258-268.
4. Ahmed HJ, Svensson LA, Cope LD et al. Prevalence of cdtABC genes encoding cytolethal distending toxin among *Haemophilus ducreyi* and *Actinobacillus actinomycetemcomitans* strains. *J Med Microbiol* 2001;50: 860-864.
5. Fabris AS, DiRienzo JM, Wikstrom M, Mayer MP. Detection of cytolethal distending toxin activity and cdt genes in *Actinobacillus actinomycetemcomitans* isolates from geographically diverse populations. *Oral Microbiol Immunol* 2002;17:231-238.
6. Belibasakis G, Johansson A, Wang Y et al. Inhibited proliferation of human periodontal ligament cells and gingival fibroblasts by *Actinobacillus actinomycetemcomitans*: involvement of the cytolethal distending toxin. *Eur J Oral Sci* 2002;110:366-373.
7. Kanno F, Korostoff J, Volgina A, DiRienzo JM. Resistance of human periodontal ligament fibroblasts to the cytolethal distending toxin of *Actinobacillus actinomycetemcomitans*. *J Periodontol* 2005;76:1189-1201.
8. DiRienzo JM, Song M, Wan LSY, Ellen RP. Kinetics of KB and HEp-2 cell responses to an invasive, cytolethal distending toxin producing strain of *Actinobacillus actinomycetemcomitans*. *Oral Microbiol Immunol* 2002;17: 245-251.
9. Belibasakis GN, Johansson A, Wang Y et al. The cytolethal distending toxin induces receptor activator of NF-kappaB ligand expression in human gingival fibroblasts and periodontal ligament cells. *Infect Immun* 2005;73:342-351.
10. Belibasakis GN, Brage M, Lagergård T, Johansson A. Cytolethal distending toxin upregulates RANKL expression in Jurkat T-cells. *APMIS* 2008;116:499-506.
11. Mogi M, Ootogoto J, Ota N, Togari A. Differential expression of RANKL and osteoprotegerin in gingival crevicular fluid of patients with periodontitis. *J Dent Res* 2004;83:166-169.
12. Pickett CL, Cottle DL, Pesci EC, Bikah G. Cloning, sequencing, and expression of the *Escherichia coli* cytolethal distending toxin genes. *Infect Immun* 1994;62:1046-1051.
13. Elwell CA, Dreyfus LA. DNase I homologous residues in CdtB are critical for cytolethal distending toxin-mediated cell cycle arrest. *Mol Microbiol* 2000;37:952-963.
14. Nestic D, Hsu Y, Stebbins CE. Assembly and function of a bacterial genotoxin. *Nature* 2004;429:429-433.
15. Peres SY, Marches O, Daigle F et al. A new cytolethal distending toxin (CDT) from *Escherichia coli* producing CNF2 blocks HeLa cell division in G2/M phase. *Mol Microbiol* 1997;24:1095-1107.
16. Lee RB, Hassane DC, Cottle DL, Pickett CL. Interactions of *Campylobacter jejuni* cytolethal distending toxin subunits CdtA and CdtC with HeLa cells. *Infect Immun* 2003;71:4883-4890.
17. McSweeney LA, Dreyfus LA. Carbohydrate-binding specif-

- icity of the *Escherichia coli* cytolethal distending toxin CdtA-II and CdtC-II subunits. *Infect Immun* 2005;73:2051-2060.
18. Mise K, Akifusa S, Watarai S et al. Involvement of ganglioside GM3 in G2/M cell cycle arrest of human monocyctic cells induced by *Actinobacillus actinomycetemcomitans* cytolethal distending toxin. *Infect Immun* 2005;73:4846-4852.
 19. Yamada T, Komoto J, Saiki K, Konishi K, Takusagawa F. Variation of loop sequence alters stability of cytolethal distending toxin (CDT): crystal structure of CDT from *Actinobacillus actinomycetemcomitans*. *Protein Sci* 2006;15:362-372.
 20. Boesze-Battaglia K, Besack D, McKay T et al. Cholesterol-rich membrane microdomains mediate cell cycle arrest induced by *Actinobacillus actinomycetemcomitans* cytolethal-distending toxin. *Cell Microbiol* 2006;8:823-836.
 21. Saiki K, Konishi K, Gomi T, Nishihara T, Yoshikawa M. Reconstitution and purification of cytolethal distending toxin of *Actinobacillus actinomycetemcomitans*. *Microbiol Immunol* 2001;45:497-506.
 22. Ko SY, Jeong DK, Ryu SH, Kim HS. Cloning and protein expression of *Actinobacillus actinomycetemcomitans* cytolethal distending toxin subunit CdtA. *J Korean Acad Periodontol* 2007;37:339-351.
 23. Lee ES, Park SY, Lee EU, Kim HS. Cloning and protein expression of *Aggregatibacter actinomycetemcomitans* cytolethal distending toxin C. *J Korean Acad Periodontol* 2008;38:317-324.
 24. Mao X, DiRienzo JM. Functional studies of the recombinant subunits of a cytolethal distending holotoxin. *Cell Microbiol* 2002;4:245-255.
 25. Nestic D, Stebbins CE. Mechanisms of assembly and cellular interactions for the bacterial genotoxin CDT. *PLoS Pathog* 2005;1:214-224.
 26. Ueno Y, Ohara M, Kawamoto T et al. Biogenesis of the *Actinobacillus actinomycetemcomitans* cytolethal distending toxin holotoxin. *Infect Immun* 2006;74:3480-3487.
 27. Lagergard T, Lundqvist A, Wising C, Gabrielsson V, Ahlman K. Formaldehyde treatment increases the immunogenicity and decreases the toxicity of *Haemophilus ducreyi* cytolethal distending toxin. *Vaccine* 2007;25:3606-3614.
 28. Mbwana J, Ahmed HJ, Ahlman K et al. Specificity of antibodies directed against the cytolethal distending toxin of *Haemophilus ducreyi* in patients with chancroid. *Microb Pathog* 2003;35:133-137.