

High Prevalence of *Haplorchis taichui*, *Phaneropsolus molenkampii*, and Other Helminth Infections among People in Khammouane Province, Lao PDR

Jong-Yil Chai^{1,*}, Eun-Taek Han², Eun-Hee Shin^{1,3}, Woon-Mok Sohn⁴, Tai-Soon Yong⁵, Keeseon S. Eom⁶, Duk-Young Min⁷, Jin-Young Um⁸, Min-Sung Park⁹, Eui-Hyug Hoang⁹, Bounlay Phommasack⁹, Bounaloth Insiengmay⁹, Soon-Hyung Lee^{1,8} and Han-Jong Rim¹⁰

¹Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799, Korea; ²Department of Parasitology, College of Medicine, Kangwon National University, Chuncheon 200-701, Korea; ³Seoul National University Bundang Hospital, Gyeonggi-do 463-707, Korea; ⁴Department of Parasitology and Institute of Health Sciences, Gyeongsang National University School of Medicine, Jinju 660-751, Korea; ⁵Department of Parasitology, College of Medicine, Yonsei University, Seoul 120-752, Korea; ⁶Department of Parasitology, College of Medicine, Chungbuk National University, Chongju 361-763, Korea; ⁷Department of Microbiology and Immunology, School of Medicine, Eulji University, Daejeon 301-746, Korea; ⁸Korea Association of Health Promotion, Seoul 157-704, Korea; ⁹Department of Hygiene and Prevention, Ministry of Public Health, Vientiane, Lao PDR; ¹⁰Department of Parasitology, College of Medicine, Korea University, Seoul 136-705, Korea

Abstract: The prevalence of liver and intestinal helminth infections, including *Opisthorchis*, *Haplorchis*, *Phaneropsolus*, hookworms, *Enterobius*, and *Taenia*, was surveyed in Khammouane province, Lao PDR. Fecal specimens were collected from 1,242 people (590 men and 652 women) in 3 Mekong riverside villages and were examined by the Kato-Katz thick smear technique. The overall helminth egg positive rate was 81.1%. The positive rate for small trematode eggs, including *Opisthorchis viverrini*, heterophyids, and leicithodendriids, was 81.1% and the positive rate for hookworms was 6.7%. To obtain adult worms, 35 people who were positive for small trematode eggs were treated with 20-30 mg/kg praziquantel and 10-15 mg/kg pyrantel pamoate, and then purged. Diarrheic stools were collected from 33 of these people and searched for helminth parasites using a stereomicroscope. Mixed infections with various helminths (*Haplorchis taichui*, *Haplorchis yokogawai*, *Prosthodendrium molenkampii*, *Phaneropsolus bonnei*, echinostomes, hookworms, *Trichostrongylus* spp., *Trichuris trichiura*, *Enterobius vermicularis*, and/or *Taenia saginata*) were found. The total number of helminth specimens collected was 20,907 (approximately 634 per person). The most common species was *H. taichui*, followed by *P. molenkampii*, *O. viverrini*, *P. bonnei*, *E. vermicularis*, hookworms, and *Trichostrongylus* spp. These results show that diverse species of intestinal nematodes, trematodes, and cestodes are infecting humans in Khammouane province, Lao PDR.

Key words: *Opisthorchis viverrini*, *Haplorchis taichui*, *Haplorchis yokogawai*, *Prosthodendrium molenkampii*, *Phaneropsolus bonnei*, hookworm, *Trichostrongylus*, *Taenia saginata*, prevalence, Khammouane province, Laos

INTRODUCTION

Soil-transmitted and fecal-borne nematodes, and food-borne liver and intestinal trematodes and cestodes, are important parasites from public health points of view [1-3]. These parasites include *Ascaris*, hookworms, *Enterobius*, *Opisthorchis*, *Clonorchis*, echinostomes, heterophyids (*Heterophyes*, *Metagonimus*, *Haplorchis*, and *Centrocestus*), leicithodendriids (*Prosthodendrium* and *Phaneropsolus*), *Diphyllobothrium*, and *Taenia* [3-5]. To prevent and control these parasites, epidemiological studies are urgent and pre-requisite.

In Lao PDR, which is located in the middle of the Indochina Peninsula, the Mekong River runs through the whole length of the country from north to south. Until the 1990s, the liver fluke, *Opisthorchis viverrini*, and soil-transmitted nematodes, including *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworms (*Ancylostoma duodenale* and *Necator americanus*), were the major helminth species known to be infecting Laotians [6-9]. However, in 1991, *Haplorchis taichui* worms were recovered from 5 Laotian students studying in Czechoslovakia [10]. Since then, mixed infections with *O. viverrini*, *H. taichui*, *Haplorchis pumilio*, *Haplorchis yokogawai*, *Centrocestus formosanus* (as *C. caninus*), *Prosthodendrium molenkampii*, *Phaneropsolus bonnei*, *Echinochasmus japonicus*, and other echinostomes have been recovered from residents of Vientiane Municipality, Saravane province, and Savan-

• Received 2 June 2009, revised 30 July 2009, accepted 3 August 2009.

* Corresponding author (cyj@snu.ac.kr)

nakheth province [11-13]. In addition to these helminths, Sayasone et al. [13] also recovered *Taenia saginata* tapeworms. Interestingly, the intensity of infection of each parasite species varied with the locality surveyed. For example, in Vientiane Municipality, *O. viverrini* infection was dominant, whereas in Saravane province *H. taichui* infection was dominant [11].

Intestinal nematode recovery data after anthelmintic treatment and purgation has never been reported in Lao PDR. Therefore, the present study aimed to determine the infection status of food-borne intestinal trematodes and cestodes, including *O. viverrini*, *H. taichui*, and *Taenia* species, and soil-transmitted nematodes, including hookworms and *Trichostrongylus*, among the riverside residents of Khammouane province.

MATERIALS AND METHODS

Khammouane province, Lao PDR, is located about 300 km distance from Vientiane Municipality and about 120 km north of Savannakhet province (Fig. 1). Three small riverside villages (Bane Nong Bone, Bane Mahaxay, and Bane Phova) located near tributaries of the Mekong River were selected for this study. Most residents are agricultural workers. Some catch freshwater fish and aquatic insect larvae from small streams and ponds, and




Fig. 1. Map showing the surveyed area of Lao PDR (arrow: Khammouane).

consume them improperly cooked.

A total of 1,242 fecal samples, one sample from each person, were collected from residents (590 men and 652 women; aged 6-80 years) of the 3 villages, in March 2003. Samples were transported to the Malaria Station in Khammouane within 2-3 days of collection and were stored at 4°C until examined. The Kato-Katz thick smear technique was used to detect helminth eggs. As it was not possible to differentiate eggs of *O. viverrini* from those of heterophyids and lecithodendriids, these eggs were collectively recorded as small trematode eggs. Fecal examinations of these residents and anthelmintic treatments were officially approved by the Ministry of Public Health, Laos, under the terms of the Korea-Laos Agreement on Parasite Control in Laos (1999-2004).

A total of 35 people who tested positive for small trematode eggs were selected for adult worm recovery at the Malaria Station. After obtaining informed consent, they were treated with a mixed single dose of 20-30 mg/kg praziquantel (Distocide®, Shinpoong Pharm. Co., Seoul, Republic of Korea) and 10-15 mg/kg of pyrantel pamoate (Combantrin®, Pfizer, New York, USA), and then purged with magnesium salt. Whole diarrheic stools passed successively 4-5 times were collected from 33 people. Worms were collected using a glass pipette and were washed several times in water. Worms were counted and some were fixed with 10% formalin under cover slip pressure, acetocarmine-stained, and morphologically identified using a light microscope.

Fecal examination results were analyzed with respect to age and sex of the subjects using the Student's *t*-test and the chi-

Table 1. Helminth eggs found in the feces of residents in Khammouane province, Laos (March 2003)

| Helminth species | No. of residents (%) | | |
|---|----------------------|------------|--------------|
| | Men | Women | Total |
| No. examined ^a | 590 | 652 | 1,242 |
| No. helminth egg positive cases (%) | 490 (83.1) | 517 (79.3) | 1,007 (81.1) |
| <i>Ascaris lumbricoides</i> | 5 (0.8) | 5 (0.8) | 10 (0.8) |
| Hookworms | 49 (8.3) | 34 (5.2) | 83 (6.7) |
| <i>Trichuris trichiura</i> | 39 (6.6) | 36 (5.5) | 75 (6.0) |
| <i>Trichostrongylus</i> spp. | 1 (0.2) | 0 (0.0) | 1 (0.08) |
| <i>Opisthorchis viverrini</i> and small trematode eggs ^b | 499 (84.6) | 508 (77.9) | 1,007 (81.1) |
| Echinostomes | 3 (0.5) | 4 (0.6) | 7 (0.6) |
| <i>Fasciolopsis buski</i> | 0 (0.0) | 2 (0.3) | 2 (0.2) |
| <i>Taenia</i> spp. | 27 (4.6) | 31 (4.8) | 58 (4.7) |

^aFecal examination was performed by the Kato-Katz smear technique. One smear was examined for each person.

^bIncluding the eggs of Heterophyidae and Lecithodendriidae.

square test. *P*-values of < 0.05 were considered statistically significant.

RESULTS

Fecal examination

The overall helminth egg positive rate was 81.1% (Table 1), and there were mixed infections involving different kinds of

trematodes and nematodes. The most frequently encountered were small trematode eggs, including those of *O. viverrini*, heterophyids, and lecitodendriids. The positive rate of small trematode eggs was 81.1% (Table 1). Other parasite eggs detected were hookworms (6.7%), *Trichuris trichiura* (6.0%), *Taenia* spp. (4.7%), *Ascaris lumbricoides* (0.8%), echinostomes (0.6%), *Fasciolopsis buski* (0.2%), and *Trichostrongylus* spp. (0.08%). The egg positive rates were not significantly (*P* > 0.01) different between men

Table 2. Adult flukes collected from residents in Khammouane province, Laos, after treatment with praziquantel and pyrantel pamoate (March 2003)

| Age and sex of resident | No. of helminth specimens collected* | | | | | | | | | | | Total |
|-------------------------|--------------------------------------|-------------------|---------------------|----------------------|------------------|----------------------------|-------------------------|---------------------|------------------------------|------------------------|----------------------------|--------|
| | <i>O. viverrini</i> | <i>H. taichui</i> | <i>H. yokogawai</i> | <i>P. molenkampi</i> | <i>P. bonnei</i> | Echino-stomes ^b | Hook-worms ^b | <i>T. trichiura</i> | <i>Trichostrongylus</i> spp. | <i>E. vermicularis</i> | <i>Taenia</i> ^c | |
| 80M | 2 | 469 | 8 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 482 |
| 65M | 1 | 96 | 13 | 7 | 2 | 0 | 0 | 0 | 1 | 8 | 1 | 129 |
| 61M | 21 | 23 | 0 | 0 | 53 | 0 | 0 | 0 | 1 | 13 | 0 | 111 |
| 60F | 19 | 115 | 0 | 5 | 1 | 0 | 0 | 0 | 0 | 6 | 0 | 146 |
| 59M | 0 | 222 | 0 | 10 | 0 | 0 | 0 | 0 | 0 | 27 | 0 | 259 |
| 58M | 27 | 208 | 0 | 69 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 308 |
| 56M | 41 | 241 | 0 | 2 | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 287 |
| 55M | 0 | 204 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 2 | 1 | 211 |
| 50M | 4 | 399 | 0 | 76 | 145 | 0 | 0 | 0 | 1 | 11 | 0 | 636 |
| 50F | 0 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 7 |
| 48M | 205 | 110 | 0 | 256 | 29 | 3 | 3 | 1 | 0 | 1 | 0 | 608 |
| 43M | 1 | 36 | 0 | 6 | 47 | 1 | 8 | 0 | 0 | 6 | 1 | 106 |
| 42M | 7 | 730 | 0 | 96 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 837 |
| 42F | 26 | 197 | 0 | 190 | 48 | 25 | 1 | 0 | 0 | 1 | 1 | 489 |
| 41F | 22 | 479 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 503 |
| 40M | 11 | 102 | 0 | 3,967 | 129 | 0 | 8 | 0 | 0 | 1 | 0 | 4,218 |
| 40F | 51 | 111 | 0 | 304 | 5 | 1 | 0 | 0 | 7 | 7 | 1 | 487 |
| 37M | 23 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 23 |
| 35F | 12 | 13 | 17 | 9 | 5 | 0 | 1 | 0 | 2 | 3 | 0 | 62 |
| 34M | 0 | 137 | 0 | 40 | 125 | 1 | 0 | 0 | 6 | 0 | 1 | 310 |
| 32M | 41 | 32 | 0 | 42 | 9 | 2 | 2 | 0 | 1 | 12 | 0 | 141 |
| 32F | 369 | 2,246 | 0 | 5 | 3 | 0 | 12 | 0 | 0 | 20 | 0 | 2,655 |
| 30F | 68 | 28 | 0 | 6 | 197 | 14 | 0 | 0 | 0 | 2 | 2 | 317 |
| 28M | 94 | 11 | 0 | 4 | 1 | 0 | 34 | 0 | 0 | 0 | 2 | 146 |
| 25M | 4 | 3,045 | 0 | 2 | 31 | 1 | 2 | 0 | 0 | 5 | 0 | 3,090 |
| 23M | 60 | 144 | 0 | 2,225 | 58 | 0 | 2 | 0 | 35 | 14 | 2 | 2,540 |
| 18M | 46 | 136 | 0 | 162 | 16 | 0 | 4 | 0 | 0 | 0 | 1 | 365 |
| 17M | 49 | 170 | 0 | 6 | 0 | 0 | 1 | 0 | 0 | 24 | 0 | 250 |
| 16M | 0 | 770 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 771 |
| 16F | 132 | 116 | 0 | 4 | 2 | 0 | 1 | 0 | 0 | 5 | 0 | 260 |
| 13M | 0 | 32 | 0 | 6 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 40 |
| 9M | 21 | 62 | 0 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 93 |
| 6M | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 |
| Total | 1,377 | 10,691 | 38 | 7,510 | 910 | 52 | 83 | 1 | 56 | 174 | 15 | 20,907 |
| (Mean No./ person) | (50) | (324) | (1) | (228) | (28) | (2) | (3) | (0.03) | (2) | (5) | (1) | (634) |

**O. viverrini*, *Opisthorchis viverrini*; *H. taichui*, *Haplorchis taichui*; *H. yokogawai*, *Haplorchis yokogawai*; *P. molenkampi*, *Prosthodendrium molenkampi*; *P. bonnei*, *Phaneropsolus bonnei*; *T. trichiura*, *Trichuris trichiura*; *E. vermicularis*, *Enterobius vermicularis*; ^bSpecies not determined; ^cThe whole or a part of strobilae with or without a scolex were recovered. All were morphologically identified as *Taenia saginata*.

and women (Table 1), or between different age groups (data not shown).

Worm collection

The adult worm collection was completed in 33 people (24 men and 9 women; age range 6-80 years) (Table 2) and a total of 20,907 helminth specimens were recovered. Between 7 and 4,218 were collected per person, with an average of 634. Intestinal flukes were recovered from 31 cases (93.9%), and *O. viverrini* was from 27 cases (81.8%). The numbers of trematode specimens were 10,691 *H. taichui*, 7,510 *P. molenkampi*, 1,377 *O. viverrini*, 910 *P. bonnei*, 52 echinostomes (species to be determined), and 38 *H. yokogawai* (Table 2). Intestinal trematodes, including *Haplorchis*, lecithodendriids, and echinostomes, comprised 91.8% of all helminth specimens recovered, whereas *O. viverrini* and intestinal nematodes comprised only 6.6% and 1.5%, respectively (Table 2).

The highest worm load of *H. taichui* collected from a single person was 3,045, and the average number of *H. taichui* recovered per treated person was 324. Specimens of *H. yokogawai* were collected only from 3 persons. As for lecithodendriids, the number of *P. molenkampi* worms ranged from 0 to 3,967 (av. 228 worms per person), and the number of *P. bonnei* ranged from 0 to 197 (av. 28 worms per person). Data on echinostomes will be published separately.

The number of nematode and cestode specimens was 174 *E. vermicularis*, 83 hookworms (*A. duodenale* and *N. americanus*), 56 *Trichostrongylus* spp., 1 *T. trichiura*, and 15 strobilae of *T. saginata* (Table 2). With regard to *E. vermicularis*, the worms were collected from 22 of 33 treated persons and the worm load averaged 5 worms (mostly females) per treated person. The worm load was higher among 17-32 and 59-65 year age groups. Hookworms were collected from 16 of 33 treated persons, averaged 3 worms per treated person, and consisted of *A. duodenale* and *N. americanus* in a 1 : 1 ratio. *Trichostrongylus* spp. specimens consisted of male and female worms (about 0.8 : 1), and were recovered from a total of 9 persons (Table 2).

DISCUSSION

Our results suggested that most of the people who were positive for small trematode eggs had mixed-infections with several species of intestinal flukes, including heterophyids and lecithodendriids, and the liver fluke, *O. viverrini*. Fecal examinations to detect helminth eggs have considerable limitations for under-

standing the prevalence and intensity of liver and intestinal flukes in Lao PDR, which was already known in the Republic of Korea [3] and Thailand [14,15]. In Lao PDR, this kind of limitation was reported in Vientiane Municipality and Saravane province [11] and also in Savannakhet province [12].

The relative prevalence of each fluke species varied remarkably by locality surveyed. For example, in Savannakhet and Vientiane, *O. viverrini* comprised up to 43.5% and 62.9% of all fluke specimens recovered, whereas in Saravane, intestinal flukes accounted for over 99.0% of all recovered flukes [11,12]. By comparison, in a recent study performed in Vientiane, Savannakhet, and Saravane, the worm loads of *O. viverrini* (av. 186 worms per worm-positive patient) and *H. taichui* (av. 207 worms) were quite similar in an analysis of 97 worm-recovered cases [13]. In our study, intestinal fluke infections predominated in Khammouane, although the degree of predominance (91.8%) was a little less than in Saravane [11].

There is a possibility that not all of the flukes (in particular, *O. viverrini*) were collected from the praziquantel treated and purged patients, since the diarrheic stools were collected only 4-5 times (for 5-6 hr) [11,12] or 6-8 times (for 24 hr) [13] after praziquantel administration. It may take longer for all *O. viverrini* to be expelled from the bile duct and gall bladder into the stools. However, data can be compared between surveyed areas, if the same protocol was applied. Thus, the individual worm loads of *O. viverrini* could be compared between different provinces. In Savannakhet province, the number of expelled worms averaged 115 worms per treated person [12], followed by an average of 58 worms in Vientiane Municipality and an average of 22 worms in Saravane province [11]. In our study, an average of 50 worms was recovered in Khammouane, which is quite similar to Vientiane [11].

The average worm load of *H. taichui* per treated person was remarkably high (8,514) in Saravane [11] compared to Khammouane (324; this study), Savannakhet (103) [12], and Vientiane (27) [11]. The highest worm load of lecithodendriid flukes was found in Khammouane (an average of 228 for *P. molenkampi* and an average of 28 for *P. bonnei*; this study), followed by Savannakhet (an average of 61 and 1, respectively) [12], Saravane (an average of 25 and 20, respectively), and Vientiane (an average of 3 and 0, respectively) [11]. These results suggest that the relative predominance of trematode species varies greatly by locality in Lao PDR.

It is of note that a total of 4 nematode species and a cestode species were recovered from the residents of Khammouane. The

higher worm loads of *E. vermicularis* among the persons at the ages of 17-32 and 59-65 year could be explained as that these people may have more contact with children, since they may have young children or grand children. Despite the relatively low prevalence of hookworm eggs (6.7%), hookworm specimens were collected from 16 of 33 treated persons. This may indicate that the actual prevalence of hookworms may be higher than the egg positive rate. Only 1 specimen of *T. trichiura* was recovered, which seems to have been due to the negligible efficacy of pyrantel pamoate and praziquantel against this nematode. A least expected result, because of a low egg positive rate (0.08%) in the fecal examination, was that a total of 56 specimens of *Trichostrongylus* spp. were recovered from 9 persons. A specimen collected from a Laotian was identified morphologically as *T. colubriformis* and its ITS-1 sequence was most similar to that of *T. colubriformis* [16]. However, the presence of *T. orientalis*, a common species in Asian countries [17], should not be ruled out in Lao PDR. The egg positive rate of *Taenia* spp. was 4.7%, but 12 of 33 treated persons expelled *Taenia* tapeworms after praziquantel treatment, which suggests a considerably higher prevalence of this tapeworm infection in this area.

ACKNOWLEDGEMENTS

We thank Dr. Sithat Insiengmay and the staff of the Center for Laboratory and Epidemiology, Department of Hygiene and Prevention, Ministry of Public Health, Vientiane, and the staff of the Khammouane Provincial Health Department, Khammouane, Lao PDR, for their help during the collection of fecal samples and for the preparation of Kato-Katz smears. We also thank the staff of the Korea Association of Health Promotion, Seoul, Republic of Korea, who participated in the Korea-Laos Cooperation Project on Parasite Control in Lao PDR (1999-2004). This study was supported by BK21 Human Life Sciences, Ministry of Education, Republic of Korea.

REFERENCES

- World Health Organization. Controlling Disease Due to Helminth Infections (eds. Crompton DWT et al.). Geneva, Switzerland. World Health Organization. 2003, p 1-248.
- Chai JY, Murrell KD, Lymbry A. Fishborne parasitic zoonoses: status and issues. *Int J Parasitol* 2005; 35: 1233-1254.
- Chai JY, Lee SH. Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* 2002; 51: 129-154.
- Shin EH, Guk SM, Lee SH, Chai JY. Trends in parasitic diseases in the Republic of Korea. *Trends Parasitol* 2008; 24: 143-150.
- Jeon HK, Kim KH, Chai JY, Yang HJ, Rim HJ, Eom KS. Sympatric distribution of three human *Taenia* tapeworms collected between 1935 and 2005 in Korea. *Korean J Parasitol* 2008; 46: 235-241.
- Sornmani S, Pathammavong O, Bunnag T, Impand P, Intarakhao C, Thirachantra S. An epidemiological survey of human intestinal parasites in Vientiane, Laos. *Southeast Asian J Trop Med Public Health* 1974; 5: 541-546.
- Kobayashi J, Vannachone B, Xeutvongsa A, Manivong K, Ogawa S, Sato Y, Pholsena K. Prevalence of intestinal parasitic infection among children in two villages in Lao PDR. *Southeast Asian J Trop Med Public Health* 1996; 27: 562-565.
- Chai JY, Hongvanthong B. A small-scale survey of intestinal helminth infections among the residents near Pakse, Laos. *Korean J Parasitol* 1998; 36: 55-58.
- Rim HJ, Chai JY, Min DY, Cho SY, Eom KS, Hong SJ, Sohn WM, Yong TS, Deodato G, Standgaard H, Phommasack B, Yun CY, Hoang EH. Prevalence of intestinal parasite infections on a national scale among primary schoolchildren in Laos. *Parasitol Res* 2003; 91: 267-272.
- Giboda M, Ditrich O, Scholz T, Viengsay T, Bouaphanh S. Human *Opisthorchis* and *Haplorchis* infections in Laos. *Trans R Soc Trop Med Hyg* 1991; 85: 538-540.
- Chai JY, Park JH, Han ET, Guk SM, Shin EH, Lin A, Kim JL, Sohn WM, Yong TS, Eom KS, Min DY, Hwang EH, Phommasack B, Insiengmai B, Rim HJ. Mixed infections with *Opisthorchis viverrini* and intestinal flukes in residents of Vientiane Municipality and Saravane Province in Laos. *J Helminthol* 2005; 79: 283-289.
- Chai JY, Han ET, Guk SM, Shin EH, Sohn WM, Yong TS, Eom KS, Lee KH, Jeong HG, Ryang YS, Hoang EH, Phommasack B, Insiengmai B, Lee SH, Rim HJ. High prevalence of liver and intestinal fluke infections among residents of Savannakhet Province in Laos. *Korean J Parasitol* 2007; 45: 213-218.
- Sayasone S, Vonghajak Y, Vanmany M, Rasphone O, Tesana S, Utzinger J, Akkhavong K, Odermatt P. Diversity of human intestinal helminthiasis in Lao PDR. *Trans R Soc Trop Med Hyg* 2009; 103: 247-254.
- Manning GS, Lertprasert P, Watanasirmit K, Chetty C. A description of newly discovered intestinal parasites endemic to northeastern Thailand. *J Med Ass Thailand* 1971; 54: 466-474.
- Radomyos P, Radomyos B, Tungtrongchitr A. Multi-infection with helminths in adults from northeast Thailand as determined by post-treatment fecal examination of adult worms. *Trop Med Parasitol* 1994; 45: 133-135.
- Yong TS, Lee JH, Sim SB, Lee JW, Min DY, Chai JY, Eom KS, Sohn WM, Lee SH, Rim HJ. Differential diagnosis of *Trichostrongylus* and hookworms eggs via PCR using ITS-1 sequence. *Korean J Parasitol* 2007; 45: 69-74.
- John DT, Petri WA Jr. Markell and Voge's Medical Parasitology. 9th ed. St. Louis, Missouri, USA. Saunders Elsevier. 2006, p 266.