

Disease-specific pluripotent stem cells

Hoon-Chul Kang, M.D.

Department of Pediatrics, Pediatric Epilepsy Clinics, Severance Children's Hospital, Epilepsy Research Institute, Yonsei University College of Medicine, Seoul, Korea

Received: 4 July 2010, Revised: 10 July 2010
Accepted: 15 July 2010

Corresponding author: Hoon-Chul Kang, M.D.
Department of Pediatrics, Yonsei University College of Medicine, 134 Shinchon-dong, Seodaemun-gu, Seoul 120-752, Korea
Tel: +82.2-2228-2050, Fax: +82.393-9118
E-mail: hipo0207@yuhs.ac

Copyright © 2010 by The Korean Pediatric Society

Induced pluripotent stem (iPS) cells are generated by epigenetic reprogramming of somatic cells through the exogenous expression of transcription factors. Recently, the generation of iPS cells from patients with a variety of genetic diseases was found to likely have a major impact on regenerative medicine, because these cells self-renew indefinitely in culture while retaining the capacity to differentiate into any cell type in the body, thereby enabling disease investigation and drug development. This review focuses on the current state of iPS cell technology and discusses the potential applications of these cells for disease modeling; drug discovery; and eventually, cell replacement therapy.

Key words: Induced pluripotent stem cells, Transcription factors

Introduction

Since their first isolation, embryonic stem (ES) cells have been regarded as the gold standard for potential use in cell-based regenerative medicine. The pluripotent nature of these cells provides them with, the ability to self-renew indefinitely in culture while retaining the capacity to differentiate into any cell type in the body¹. However, the fact that ES cells can only be derived from early-stage embryos precludes the establishment of autologous cell lines for patients and immune rejection hinders the use of nonautologous ES cell lines².

The advent of mammalian somatic cell nuclear transfer (SCNT) first appeared to offer solutions for overcoming these roadblocks. SCNT is a technique whereby the DNA of an unfertilized egg is replaced by the DNA of a somatic cell. Although significant advancements toward this goal have been made, successful SCNT into human cells remains an elusive goal and is fraught with social and logistical concerns³. Alternative methods for deriving pluripotent cells, such as cell fusion and culture-induced reprogramming, have been developed, but these approaches still

suffer from severe practical and technical limitations⁴.

In 2006, Takahashi⁵ and Yamanaka reported the astonishing discovery that somatic cells could be reprogrammed to adopt the pluripotent stem cell state. The generation of pluripotent cells by exogenous expression of transcription factors circumvents many previous limitations because this approach is not technically demanding and does not require embryonic material or oocytes. We therefore believe that this discovery which is now known as "induced pluripotent stem cell technology" (iPS) will have a significant impact on regenerative medicine. In this article we review the current methodologies used for generating iPS cells and then discuss their potential clinical applications.

Reprogramming somatic cells towards pluripotency by defined transcription factors⁵

To establish the transcription factors necessary for reprogramming of differentiated cells, mouse embryonic fibroblasts and tail-tip fibroblasts of mice were initially used. Reprogrammed somatic cells were identified by selecting for the reactivation of

the Fbx-15 gene locus with a neomycin-reporter cassette whose expression is restricted to the early stage of embryonal development. Cells which are only resistant to G418 selection allowed growth of ES-cell-like colonies and cells derived by sub-culturing these ES-like colonies were designated as “induced pluripotent stem cells.” It was found that of the 24 candidate genes that are implicated in the establishment and maintenance of the pluripotent state, Oct4 (Pou5f1), Sox2, c-Myc, and Klf4, were sufficient to mediate reprogramming. These iPS cells had fully demethylated Oct4 and Nanog promoters and could contribute to the formation of germline-competent chimeras upon injection of blastocysts, a characteristic which demonstrates true pluripotency.

Preferred methods for preparation of iPS cells

First-generation iPS cells were generated by retroviral transduction⁵. Since then, the technique has been optimized and reproduced in a number of different processes⁶. Several important variables include the choice of cell type for reprogramming, the choice of the combination of genes and small molecules used for reprogramming, as well as the gene transfer method (Fig. 1).

For the first reprogramming attempts of both mouse and human cells, fibroblasts were used as the starting cell population⁵. Since the original success of fibroblast reprogramming, multiple cell types have been used for reprogramming, including keratinocytes, neural progenitor cells, hepatocytes, stomach epithelial cells, pancreatic beta cells and intestinal epithelial cells⁷. Several factors must therefore be considered in determining the optimal cell type for a given application: (1) the ease with which reprogramming factors can be introduced, which varies according to cell type and delivery approach; (2) the availability and ease of derivation of the given cell type; and (3) the age and source of the cells used⁶. An inverse correlation has been effectively demonstrated for the differentiation

stage of target cells. For example, with mouse hematopoietic stem and progenitor cells being more efficiently reprogrammed than terminally differentiated B and T lymphocytes^{8,9}. While fibroblasts are likely to remain the choice cell type in basic research efforts, iPS cells derived for therapeutic purposes will require the donor cells to be easily attainable, less likely to contain genetic aberrations, and easy to reprogram using transient approaches⁶.

Since the first report of reprogramming, it has become evident that there is some flexibility in the factors needed to reprogram somatic cells. Of the original four transcription factor-encoding genes, c-Myc is not strictly required for the reprogramming of somatic cells and Klf4 and Sox2 have been shown to be dispensable in reprogramming strategies utilizing cell types that endogenously express them. Oct4 is the only factor that cannot be replaced by other family members and the only one that has been required in every reprogramming strategy in either mouse or human cells⁸.

To aid in the production of epigenetic modifications during the reprogramming process, several studies have examined the use of small-molecule compounds. Importantly, small molecules, including DNA methyltransferase inhibitors¹⁰, the histone deacetylase inhibitor valproic acid¹¹, and the histone methyltransferase inhibitor BIX-0 1294¹², have been shown to substantially improve the efficiency of reprogramming, even in cases without inclusion of exogenous Klf4 and c-Myc (for valproate) or exogenous Sox2 and c-Myc (for BIX-01294). In addition, inhibition of DNA methyltransferase activity by 5-aza-cytidine also was found to improve reprogramming efficiency¹³. Recently, another small molecule, transforming growth factor RI kinase inhibitor II, has also been reported to replace Sox2 by the induction of Nanog¹⁴. Of note, the effects of these small-molecule agents are not directed towards specific reprogramming events, and it remains to be determined whether full reprogramming through the exclusive use of chemicals is possible.

iPS cells have been generated using a number of different gene transfer methods, including retroviral, lentiviral, and adenoviral vectors and nonviral plasmids, and recently by direct recombinant protein delivery⁸. Initial generations of mouse and human iPS cells employed retroviral vectors⁵ and constitutive lentiviruses¹⁵. These viral systems, however, have been criticized because they are permanently integrated into the genome and endeavors to make iPS cells more therapeutically applicable have led to the pursuit of non-integrative approaches. A refined system which employs the Crelox system or transposases in which reprogramming factors could be delivered into the host genome and removed without any residual elements has been recently developed¹⁶. More importantly, iPS cell generation has now been achieved without genomic




Fig. 1. Overview of the processes for inducing the derivation of pluripotent stem (iPS) cells and clinical applications.

integration. This has been accomplished using adenoviruses¹⁷⁾, repeated plasmid transfection¹⁸⁾, and recombinant proteins¹⁹⁾ and via the use of episomal vectors²⁰⁾ in human cells. However, the reprogramming efficiency obtained using these delivery systems tends to be dramatically lower for most cell types relative to the standard retroviral/lentiviral mediated reprogramming⁷⁾. Ultimately, the logistical, financial, and practical aspects of each technique must be taken into account. It is likely that the optimal cell reprogramming method will be determined by the intended application of the reprogrammed cells. Currently, for example, the fastest and most efficient method of creating human iPS cell lines through lentiviral and retroviral transduction might represent the preferred approaches for generating iPS cells for use in large-scale drug-screening programs and disease modeling⁸⁾.

Clinical relevance

The recent introduction of iPS cell technology has opened a new avenue not only for cell replacement therapy but also for disease modeling and drug development²¹⁾. The generation of human iPS cells from patients with a variety of genetic diseases with either Mendelian or complex inheritance including X-linked adrenoleukodystrophy (X-ALD), amyotrophic lateral sclerosis, Duchenne and Becker muscular dystrophy, Parkinson's disease, Huntington's disease, Gaucher disease type III, and Down syndrome is also possible. Such disease-specific iPS cells offer an unprecedented opportunity to recapitulate both normal and pathologic human tissue formation *in vitro*, thereby enabling disease investigation, drug development and cell therapy²¹⁾.

Conclusion

As a result of its tremendous potential in a wide variety of clinical and research applications, there is great interest in cellular reprogramming. The recent successes in iPS cell derivation without the use of viral vectors and genomic integration from human cells has brought the realization of the therapeutic potential of iPS cell technology closer than ever. Importantly, however, there is still much to learn about how this reprogramming process works. The key steps involved in this process consist of the choice of reprogramming factors and molecules, their delivery method, and the choice of target cell type. Given the high level of interest in iPS cells and the large number of laboratories around the world that are now studying these cells, we are hopeful that iPS cell technology will have a positive impact on therapeutic interventions.

References

- 1) Sell S. Stem cells: What are they? Where do they come from? Why are they here? When do they go wrong? Where are they going? In: Sell S, editor, Stem cells handbook. 1st ed. Totowa: Humana Press Co, 2004:1-18.
- 2) Kirschstein R. The stem cell. In: Kirschstein R, editor. Stem cells: scientific progress and future research directions. 1st ed. Maryland: National Institutes of Health, 2001:1-4.
- 3) Wakayama T, Hayashi Y, Ogura A. Participation of the female pronucleus derived from the second polar body in full embryonic development of mice. *J Reprod Fertil* 1997;110:263-6.
- 4) Hochedlinger K, Jaenisch R. Nuclear reprogramming and pluripotency. *Nature* 2006; 441:1061-7.
- 5) Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006;126:663-76.
- 6) Maherali N, Hochedlinger K. Guidelines and techniques for the generation of induced pluripotent stem cells. *Cell Stem Cell* 2008;3:595-605.
- 7) Cox JL, Rizzino A. Induced pluripotent stem cells: what lies beyond the paradigm shift. *Experimental Biology and Medicine* 2010;235:148-58.
- 8) Kiskinis E, Eggan K. Progress toward the clinical application of patient-specific pluripotent stem cells. *J Clin Invest* 2010;120:51-9.
- 9) Eminli S, Foudi A, Stadtfeld M, Maherali N, Ahfeldt T, Mostoslavsky G, et al. Differentiation stage determines potential of hematopoietic cells for reprogramming into induced pluripotent stem cells. *Nat Genet* 2009;41:968-76.
- 10) Mikkelsen TS, Hanna J, Zhang X, Ku M, Wernig M, Schorderet P, et al. Dissecting direct reprogramming through integrative genomic analysis. *Nature* 2008;454:49-55.
- 11) Huangfu D, Maehr R, Guo W, Eijkelenboom A, Snitow M, Chen AE, et al. Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nat Biotechnol* 2008;26:795-7.
- 12) Shi Y, Despons C, Do JT, Hahm HS, Schöler HR, Ding S. Induction of pluripotent stem cells from mouse embryonic fibroblasts by Oct4 and Klf4 with small-molecule compounds. *Cell Stem Cell* 2008;3:568-74.
- 13) Mikkelsen TS, Hanna J, Zhang X, Ku M, Wernig M, Schorderet P, et al. Dissecting direct reprogramming through integrative genomic analysis. *Nature* 2008;454:49-55.
- 14) Wernig M, Meissner A, Cassady JP, Jaenisch R. c-Myc is dispensable for direct reprogramming of mouse fibroblasts. *Cell Stem Cell* 2008;2:10-2.
- 15) Belloch R, Venere M, Yen J, Ramalho-Santos M. Generation of induced pluripotent stem cells in the absence of drug selection. *Cell Stem Cell* 2007;1: 245-7.
- 16) Soldner F, Hockemeyer D, Beard C, Gao Q, Bell GW, Cook EG, et al. Parkinson's disease patient-derived induced pluripotent stem cells free of viral reprogramming factors. *Cell* 2009;136:964-77.
- 17) Stadtfeld M, Nagaya M, Utikal J, Weir G, Hochedlinger K. Induced pluripotent stem cells generated without viral integration. *Science* 2008;322:945-9.
- 18) Gonzalez F, Barragan Monasterio M, Tiscornia G, Montserrat Pulido N, Vassena R, et al. Generation of mouse-induced pluripotent stem cells by transient expression of a single nonviral polycistronic vector. *Proc Natl Acad Sci* 2009;106:8918-22.

- 19) Kim D, Kim CH, Moon JI, Chung YG, Chang MY, Han BS, et al. Generation of human induced pluripotent stem cells by direct delivery of reprogramming proteins. *Cell Stem Cell* 2009;4:472-6.
- 20) Yu J, Hu K, Smuga-Otto K, Tian S, Stewart R, Slukvin II, Thomson JA. Human induced pluripotent stem cells free of vector and transgene sequences. *Science* 2009;324:797-801.
- 21) Park IH, Arora N, Huo H, Maherali N, Ahfeldt T, Shimamura A, et al. Disease-specific induced pluripotent stem cells. *Cell* 2008;134:877-86.