

Cytotoxic Sesquilignans from the Roots of *Saururus chinensis*

Young-Won Chin, Xing-Fu Cai, Kyung-Seop Ahn, Hyeong-Kyu Lee, and Sei-Ryang Oh*

Bio-Therapeutics Research Institute, Korea Research Institute of Bioscience and Biotechnology, Ochanggeup, Cheongwongun, ChungBuk 363-883, Korea. *E-mail: seiryang@kribb.re.kr
Received April 26, 2010, Accepted May 25, 2010

Key Words: *Saururus chinensis*, Sesquilignans, Cytotoxicity, Saururaceae

Saururus chinensis Hort. ex Loudon (Saururaceae) is a perennial herb distributed in China and Korea, and has been used as a folk medicine for the treatment of edema, gonorrhea, jaundice, pneumonia, and several inflammatory diseases in Korea.¹ Previous studies of *S. chinensis* reported the occurrence of lignans,²⁻⁹ aristolactams,¹⁰ flavonoids,¹¹ and furanoditerpenes,¹² and a wide range of biological activities including antioxidant activity,⁵ hepatoprotective activity,⁶ cytotoxic activity,¹³⁻¹⁸ anti-inflammatory activity,¹⁹⁻²² anti-atherogenic activity,^{21,23} and immunosuppressive activity.²⁴ This paper reports the structures elucidation of the two new lignans and six known compounds, along with their cytotoxicity.

Compound **1** was obtained as a colorless powder and its molecular formula was determined to be C₃₁H₃₈O₈, based on the [M-H]⁻ peak at *m/z* 537.2471 (calcd 537.2488) in the HRESIMS. The ¹H-NMR spectroscopic data of compound **1** indicated the presence of a tetrahydrofuran-type lignan unit, as judged from the signals for two methines at δ_H 2.34 (H-8 and H-8'), for oxymethines at δ_H 5.65 (H-7 and H-7'), and for methyl groups at δ_H 0.78 (H-9) and 0.74 (H-9'), as well as the signals at δ_H 7.20 (d, *J* = 1.7 Hz, H-2), 7.30 (*J* = 8.2 Hz, H-5), 7.36 (br d, *J* = 8.2 Hz, H-6), 7.19 (d, *J* = 1.7 Hz, H-2'), 7.34 (d, *J* = 8.2 Hz, H-5'), and 7.09 (dd, *J* = 8.2, 1.7 Hz, H-6') corresponding to two 1,3,4-trisubstituted benzene rings.^{4,25} An additional phenylpropanoid unit was observed in compound **1** from the proton signals for two oxymethines at δ_H 4.95 (H-8''), and 5.42 (H-7''),

for a methyl at δ_H 1.60 (H-9'') and for a 1,3,4-trisubstituted benzene ring at δ_H 7.58 (d, *J* = 1.6 Hz, H-2''), 7.01 (d, *J* = 8.2 Hz, H-5''), and 7.07 (dd, *J* = 8.2, 1.6 Hz, H-6'').⁴ The location of the phenylpropanoid group was confirmed by the HMBC correlation between H-8'' (δ_H 4.95) and C-4' (δ_C 147.5), suggesting this phenylpropanoid is attached to C-4' on the tetrahydrofuran-type lignan moiety through an ether linkage. The relative stereochemistry of the tetrahydrofuran ring in compound **1** was established by the observed NOE correlations of H-9 with H-8', H-2, and H-6 as well as H-9' with H-8, H-2', and H-6', indicating the 7,8-*cis*-8,8'-*trans*-7',8'-*cis* configuration.^{4,26} In addition, the chemical shifts of C-9'' (δ_C 15.3), C-7'' (δ_C 75.7), and C-8'' (δ_C 81.1), and the coupling constant (*J* = 3.4 Hz) of H-7'' was supportive of the relative configuration of C-7'' and C-8'' as being *erythro*.^{4,26} Furthermore, the positive Cotton effect at 231 nm enabled to assign the configuration of C-7'' and 8'' as *R* and *S*, respectively.²⁷ Thus, the structure of compound **1** was determined as 7''*R*,8''*S*-saucerneol, a diastereomer of (-)-saucerneol (**3**).

Saucerneol J (**2**) had a molecular formula (C₃₀H₃₆O₈) and exhibited a close resemblance to compound **1** in their ¹H and ¹³C NMR spectroscopic data except for the presence of three methoxy groups. There were differences in the chemical shifts and coupling constants of H-7' in compounds **2** (δ_H 4.54, *J* = 9.3 Hz) and **1** (δ_H 5.65, *J* = 6.4 Hz), indicating the opposite configuration at C-7' position in compound **2** compared to compound **1**.




Figure 1. Structures of compounds **1-2**.




Figure 2. Key ¹H-¹H COSY (—) and HMBC (H→C) and NOESY (---) correlations of compound **1**.

Table 1. ^1H - and ^{13}C -NMR data (δ) of **1-2** (Pyridine- d_5)^a

position	1		2	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1	-	136.5 (s)	-	135.6 (s)
2	7.20 d, 1.7	111.6 (d)	7.25 d, 1.7	112.1 (d)
3	-	151.5 (s)	-	151.5 (s)
4	-	147.6 (s)	-	147.8 (s)
5	7.30 d, 8.2	117.5 (d)	7.28 d, 8.1	116.5 (d)
6	7.36 br d, 8.2	120.1 (d)	7.31 br d, 8.1	120.6 (d)
7	5.65 d, 6.4	84.6 (d)	5.30 d, 8.6	82.7 (d)
8	2.34 ddq, 12.8, 6.4, 6.4	44.5 (d)	2.24 ddq, 8.6, 7.0, 6.5	45.7 (d)
9	0.78 d, 6.4	15.3 (q)	0.73 d, 7.0	14.9 (q)
1'	-	133.8 (s)	-	133.2 (s)
2'	7.19 d, 1.7	112.2 (d)	7.42 d, 1.7	111.2 (d)
3'	-	148.9 (s)	-	149.2 (s)
4'	-	147.5 (s)	-	148.5 (s)
5'	7.34 d, 8.2	116.6 (d)	7.30 d, 8.1	116.1 (d)
6'	7.09 dd, 8.2, 1.7	119.8 (d)	7.32 dd, 8.1, 1.7	120.5 (d)
7'	5.65 d, 6.4	84.3 (d)	4.54 d, 9.3	87.4 (d)
8'	2.34 ddq, 12.8, 6.4, 6.4	44.5 (d)	1.93 ddq, 9.3, 6.5, 6.5	47.9 (d)
9'	0.74 d, 6.4	15.3 (q)	1.05 d, 6.5	14.5 (q)
1''	-	136.8 (s)	-	135.1 (s)
2''	7.58 d, 1.6	112.2 (d)	7.59 d, 1.6	111.3 (d)
3''	-	150.2 (s)	-	148.9 (s)
4''	-	149.5 (s)	-	147.7 (s)
5''	7.01 d, 8.2	112.6 (d)	7.31 d, 8.1	115.8 (d)
6''	7.07 dd, 8.2, 1.6	120.3 (d)	7.29 dd, 8.1, 1.6	120.7 (d)
7''	5.42 d, 3.4	75.7 (d)	5.42 d, 3.9	74.8 (d)
8''	4.95 dq, 6.2, 3.4	81.1 (d)	4.90 dq, 6.2, 3.9	80.3 (d)
9''	1.60 d, 6.2	15.3 (q)	1.60 d, 6.2	14.3 (q)
3-OCH ₃	3.84 s	56.5 (q)	3.82 s	56.4 (q)
3'-OCH ₃	3.85 s	56.5 (q)	3.86 s	56.4 (q)
3''-OCH ₃	3.82 s	56.3 (q)	3.80 s	56.3 (q)
4''-OCH ₃	3.77 s	56.4 (q)	-	-

^aAssignments were based on ^1H - ^1H COSY, DEPT, HMQC and HMBC spectra.

The CD spectroscopic data exhibited the positive Cotton effect at 232 nm in the same manner. Therefore, the structure of compound **2** was confirmed to be 7'-*epi*-7''*R*,8''*S*-4''-demethyl-saucerneol.

The known compounds were in good agreement with previously reported NMR data and were consequently identified as (-)-saucerneol (**3**) with a $[\alpha]_{\text{D}}$ value of -71.1 (*c* 0.1 MeOH) [Lit. $[\alpha]_{\text{D}}$ -91.8],²⁵ *threo,erythro*-manassantin A (**4**),¹⁶ 4-*O*-demethylmanassantin B (**5**),¹⁸ *erythro,erythro*-manassantin A (**6**),¹⁶ manassantin B (**7**)²⁸ and manassantin A (**8**).¹⁷

All the compounds isolated were evaluated against HL-60 (human promyelocytic leukemia) cells. Compounds **1-8** exhibited cytotoxicity (IC_{50} , 0.5, 7.1, 3.3, 5.2, 3.6, 2.3, 8.5 and 0.8 μM , respectively) against HL-60 cell lines (camptothecin, IC_{50} 0.8 μM).

Experimental Section

General experimental procedures. Melting points were determined on a Kofler micro-hotstage (uncorrected). Optical rota-

tions were measured on a JASCO P-1020 polarimeter. UV spectra were measured on a Shimadzu UV-1601 UV-visible. CD spectroscopic data were obtained from JASCO-720 CD spectrometer. The NMR spectra were recorded on a Varian Unity 400 FT-NMR spectrometers with the tetramethylsilane as an internal standard. Chemical shifts are presented in ppm. HRESIMS were measured on a Waters Q-ToF Premier mass spectrometer. Column chromatography (CC) was performed on silica gel (70 - 230 and 230 - 400 mesh, Merck), reverse-phase C18 gel (40 μm , Nacalai Inc., Japan). Thin layer chromatography (TLC) was performed on Kieselgel 60 F₂₅₄ (Merck) or RP-18 F₂₅₄ (Merck) plates. Spots were visualized by spraying 10% aqueous H₂SO₄ solution on the plates and heating them for 5 min.

Plant material. The roots of *Saururus chinensis* was collected at Jeju (Korea) in July 2008 and dried at room temperature. A voucher specimen (00250) is deposited at the Plant Extract Bank, Korea Research Institute of Bioscience and Biotechnology, Daejeon, Korea.

Extraction and isolation. The dried roots of *S. chinensis*

(6.5 kg) was extracted with MeOH at room temperature (3 × 20 L) to obtain 0.65 kg of the solid extract. The MeOH extract was suspended in H₂O and extracted with EtOAc (3 × 3 L) to give the EtOAc-soluble fractions (85 g). The EtOAc-soluble fraction (84 g) was chromatographed on a silica gel column eluted with a stepwise gradient of hexane and EtOAc to yield 14 fractions (fr. SC1-SC14). Fr. SC12 (1.2 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 7 : 3) to yield 19 sub-fractions (fr. SC12-1-SC12-19). Fr. SC12-15 (0.14 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 2 : 1) to give compounds **1** (13.2 mg) and **3** (98.1 mg). Fr. SC12-19 (0.103 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 2 : 1) to give compound **4** (17.8 mg). Fr. SC14 (0.88 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 8 : 2) to yield seven sub-fractions (Fr. SC14-1-SC14-7). Fr. SC14-3 (0.12 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 3 : 2) to give compound **2** (5.3 mg). Fr. SC14-6 (0.36 g) was chromatographed on a RP C-18 column (MeOH/H₂O, 3 : 2) to give compounds **5** (76.7 mg), **6** (5.3 mg), **7** (18.3 mg), and **8** (43.3 mg).

erythro-Saucerneol (1): Colorless powder, mp 85 - 86 °C. $[\alpha]_D^{25}$ -18 (c 0.1, MeOH). UV λ_{max} (MeOH) nm (log ϵ): 206 (3.74), 284 (2.94). ¹H- and ¹³C-NMR data see Tables 1. HRESIMS *m/z* 537.2471 [M-H]⁻ (Calcd for C₃₁H₃₇O₈: 537.2488). CD (c 0.0004 MeOH): $[\theta]_{231}^{25} +12,075$.

Saucerneol J (2): Colorless powder, mp 75 - 76 °C. $[\alpha]_D^{25}$ -10 (c 0.1, MeOH). UV λ_{max} (MeOH) nm (log ϵ): 206 (3.81), 282 (2.67). ¹H- and ¹³C-NMR data see Tables 1 and 2. HRESIMS *m/z* 523.2310 [M-H]⁻ (Calcd for C₃₀H₃₅O₈: 523.2332). CD (c 0.0006 MeOH): $[\theta]_{232}^{25} +18,121$.

Cytotoxicity evaluation. All the isolates were assessed with the HL-60 (human promyelocytic leukemia) cells according to the established protocol.²⁹

Acknowledgments. This research was supported by a grant of KRIBB Research Initiative Program (KGS2241012).

References and Notes

- Chung, B. S.; Shin, M. G. *Dictionary of Korean Folk Medicine*; Young Lim Publishing: Seoul, 1990; pp 813-814.
- Ahn, B. T.; Lee, S.; Lee, S. B.; Lee, E. S.; Kim, J. G.; Bok, S. H.; Jeong, T. S. *J. Nat. Prod.* **2001**, *64*, 1562.
- Hwang, B. Y.; Lee, J. H.; Jung, H. S.; Kim, K. S.; Nam, J. B.; Hong, Y. S.; Paik, S. G.; Lee, J. J. *Planta Med.* **2003**, *69*, 1096.
- Hwang, B. Y.; Lee, J. H.; Nam, J. B.; Hong, Y. S.; Lee, J. J. *Phytochemistry* **2003**, *64*, 765.
- Lee, W. S.; Baek, Y. I.; Kim, J. R.; Cho, K. H.; Sok, D. E.; Jeong, T. S. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5623.
- Sung, S. H.; Kim, Y. C. *J. Nat. Prod.* **2000**, *63*, 1019.
- Sung, S. H.; Lee, E. J.; Cho, J. H.; Kim, H. S.; Kim, Y. C. *Biol. Pharm. Bull.* **2000**, *23*, 666.
- Seo, C. S.; Zheng, M. S.; Woo, M. H.; Lee, C. S.; Lee, S. H.; Jeong, B. S.; Chang, H. W.; Jahng, Y.; Lee, E. S.; Son, J. K. *J. Nat. Prod.* **2008**, *71*, 1771.
- Sung, S. H. *Fitoterapia* **2006**, *77*, 487.
- Kim, S. R.; Sung, S. H.; Kang, S. Y.; Koo, K. A.; Kim, S. H.; Ma, C. J.; Lee, H. S.; Park, M. J.; Kim, Y. C. *Planta Med.* **2004**, *70*, 391.
- Kang, T. H.; Cho, H.; Oh, H.; Sohn, D. H.; Kim, Y. C. *Fitoterapia* **2005**, *76*, 115.
- Hwang, B. Y.; Lee, J. H.; Nam, J. B.; Kim, H. S.; Hong, Y. S.; Lee, J. J. *J. Nat. Prod.* **2002**, *65*, 616.
- Seo, B. R.; Lee, K. W.; Ha, J.; Park, H. J.; Choi, J. W.; Lee, K. T. *Carcinogenesis* **2004**, *25*, 1387.
- Seo, B. R.; Yoo, C. B.; Park, H. J.; Choi, J. W.; Seo, K.; Choi, S. K.; Lee, K. T. *Biol. Pharm. Bull.* **2004**, *27*, 1594.
- Song, S. Y.; Lee, I.; Park, C.; Lee, H.; Hahm, J. C.; Kang, W. K. *Int. J. Mol. Med.* **2005**, *16*, 517.
- Hahm, J. C.; Lee, I. K.; Kang, W. K.; Kim, S. U.; Ahn, Y. J. *Planta Med.* **2005**, *71*, 464.
- Hossain, C. F.; Kim, Y. P.; Baerson, S. R.; Zhang, L.; Bruick, R. K.; Mohammed, K. A.; Agarwal, A. K.; Nagle, D. G.; Zhou, Y. D. *Biochem. Biophys. Res. Commun.* **2005**, *333*, 1026.
- Hodges, T. W.; Hossain, C. F.; Kim, Y. P.; Zhou, Y. D.; Nagle, D. G. *J. Nat. Prod.* **2004**, *67*, 767.
- Hwang, B. Y.; Lee, J. H.; Koo, T. H.; Kim, H. S.; Hong, Y. S.; Ro, J. S.; Lee, K. S.; Lee, J. J. *Planta Med.* **2002**, *68*, 101.
- Lee, A. K.; Sung, S. H.; Kim, Y. C.; Kim, S. G. *Br. J. Pharmacol.* **2003**, *139*, 11.
- Rho, M. C.; Kwon, O. E.; Kim, K.; Lee, S. W.; Chung, M. Y.; Kim, Y. H.; Hayashi, M.; Lee, H. S.; Kim, Y. K. *Planta Med.* **2003**, *69*, 1147.
- Moon, T. C.; Kim, J. C.; Song, S. E.; Suh, S. J.; Seo, C. S.; Kim, Y. K.; Jin, M.; Yang, J. H.; Son, J. K.; Jahng, Y.; Kim, C. H.; Chang, H. W. *Int. Immunopharmacol.* **2008**, *8*, 1395.
- Lee, W. S.; Lee, D. W.; Baek, Y. I.; An, S. J.; Cho, K. H.; Choi, Y. K.; Kim, H. C.; Park, H. Y.; Bae, K. H.; Jeong, T. S. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3109.
- Park, S. Y.; Lee, S. H.; Choi, W. H.; Koh, E. M.; Seo, J. H.; Ryu, S. Y.; Kim, Y. S.; Kwon, D. Y.; Koh, W. S. *Planta Med.* **2007**, *73*, 674.
- Hanessian, S.; Reddy, G. J.; Chahal, N. *Org. Lett.* **2006**, *8*, 5477.
- Tofern, B.; Jenett-Siems, K.; Siems, K.; Jakupovic, J.; Eich, E.; *Phytochemistry* **2000**, *53*, 119.
- Huo, C.; Liang, H.; Zhao, Y.; Wang, B.; Zhang, Q. *Phytochemistry* **2008**, *69*, 788.
- Park, H. J.; Kim, R. G.; Seo, B. R.; Ha, J.; Ahn, B. T.; Bok, S. H.; Lee, Y. S.; Kim, H. J.; Lee, K. T. *Planta Med.* **2003**, *69*, 947.
- Park, B. Y.; Oh, S. R.; Ahn, K. S.; Kwon, O. K.; Lee, H. K. *Int. Immunopharmacol.* **2008**, *8*, 967.