

## Protective Effect of Enzymatic Extracts from *Sargassum coreanum* on H<sub>2</sub>O<sub>2</sub>-induced Cell Damage

Seok-Chun Ko<sup>1</sup>, Sung-Myung Kang<sup>1</sup>, Seung-Hong Lee<sup>1</sup>, Ginnae Ahn<sup>1</sup>,  
Kil-Nam Kim<sup>2</sup>, Yong-Tae Kim<sup>3</sup>, Jin-Soo Kim<sup>4</sup>,  
Min Soo Heu<sup>5</sup> and You-Jin Jeon<sup>1,6\*</sup>

<sup>1</sup>Department of Marine Life Science, Jeju National University, Jeju, 690-756, Korea

<sup>2</sup>Jeju Biodiversity Research Institute (JBRI) and Jeju Hi-Tech Industry  
Development Institute (HiDI), Jeju 697-943, Korea

<sup>3</sup>Department of Food Science and Biotechnology, Kunsan National University,  
Kunsan 573-701, Korea

<sup>4</sup>Department of Seafood Science and Technology/Institute of Marine Industry,  
Gyeongsang National University, Tongyeong 650-160, Korea

<sup>5</sup>Department of Food Science and Nutrition/Institute of Marine Industry,  
Gyeongsang National University, Jinju 660-701, Korea

<sup>6</sup>Marine and Environmental Research Institute, Jeju National University,  
Jeju 695-814, Korea

In our previous study, we preliminarily demonstrated that Celluclast and Neutrased extracts exhibited the strongest H<sub>2</sub>O<sub>2</sub>-scavenging activities among five carbohydrases (Viscozyme, Celluclast, Termamyl, Ultraflo and AMG) and five proteases (Kojizyme, Alcalase, Flavourzyme, Protamex and Neutrased) extracts. Thus, Celluclast and Neutrased extracts were selected for use in further experiments and were separated into four different molecular weight fractions (<5, 5-10, 10-30 and >30 kDa). Among them, the 5-10 kDa fraction showed the highest H<sub>2</sub>O<sub>2</sub>-scavenging activity. The 5-10 kDa fraction also strongly enhanced cell viability against H<sub>2</sub>O<sub>2</sub>-induced oxidative damage. Furthermore, the fraction reduced the proportion of apoptotic cells induced by H<sub>2</sub>O<sub>2</sub>, as demonstrated by decreased sub-G<sub>1</sub> hypodiploid cells and decreased apoptotic body formation by flow cytometry. These results indicated that the 5-10 kDa fraction of the Celluclast and Neutrased extracts from *S. coreanum* exhibited strong antioxidant activity over H<sub>2</sub>O<sub>2</sub>-mediated cell damage *in vitro*.

Key words: *Sargassum coreanum*, Enzymatic extract, H<sub>2</sub>O<sub>2</sub>, Cell damage

### Introduction

Reactive oxygen species (ROS) include free radicals, such as the peroxy (ROO·), nitric oxide (NO·), hydroxyl (·OH), and superoxide radicals (O<sub>2</sub><sup>·-</sup>), and non free-radicals, such as singlet oxygen (<sup>1</sup>O<sub>2</sub>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), that are formed during normal metabolic processes (Heo et al., 2008). ROS can readily oxidize cellular components such as lipids, proteins, and lipoproteins (Lee et al., 2009). High levels of ROS are considered to be responsible for various pathological conditions including cancer, cardiovascular disease, and aging (Cox and Choen,

1996; Ames, 1998). Among the ROS, H<sub>2</sub>O<sub>2</sub> plays a pivotal role because it is generated in nearly all types of oxidative stress and it can diffuse freely in and out of cells and tissues (Halliwell and Aruoma, 1991).

Synthetic antioxidants, such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), and tert-butylhydroquinone (TBHQ), have been used to reduce oxidative damage in the human body (Kim et al., 2008; Sherwin, 1990). However, the use of these antioxidants in human is also associated with some side effects, such as liver damage and carcinogenesis (Lindenschmidt et al., 1986). Thus, much attention has been focused on natural antioxidants in functional foods instead of synthetic antioxidants.

Seaweeds are known to provide an abundance of

\*Corresponding author: youjinj@jejunu.ac.kr

bioactive materials with valuable pharmaceutical and biomedical potential. According to the results of previous studies, the bioactive substances contained in seaweeds differ from those in land plants, and various of these constituents have been demonstrated to possess antioxidant, antitumor, anticoagulant, and antibacterial effects (Kashiwagi et al., 1980; Gonzalez et al., 1982; Kosovel et al., 1988). In particular, brown seaweeds possess a variety of biological compounds, including fucoxanthin, xanthophyll, and polyphenol-containing phlorotannins. Several studies on these kinds of compounds have pointed out a variety of biological benefits, including antioxidant, anticoagulant, anti-hypertensive, antibacterial, and antitumor activities (Athukorala and Jeon, 2005; Heo et al., 2005; Kotake-Nara et al., 2005; Mayer and Hmann, 2004; Nagayama et al., 2002).

In a previous study (Heo et al., 2005), a novel extraction technique using food-grade digestive enzymes, including proteases and carbohydrases, was used to extract products with a variety of bioactivities including antioxidant, anti-inflammatory, and anti-hypertensive effects (Kim et al., 2008; Athukorala and Jeon, 2005; Ahn et al., 2008). These seaweed-derived enzymatic extracts have been shown to have some important benefits, including higher yields, water-solubility, and safety, because this technique does not require the use of organic solvents or other toxic chemicals. In the present study, *S. coreanum*, a brown seaweed, was extracted enzymatically with different proteases and carbohydrases used in the food industry to prepare water-soluble extracts, and then the antioxidant effects of the resulting extracts were assessed with regard to their ability to scavenge H<sub>2</sub>O<sub>2</sub> and their protective effects against H<sub>2</sub>O<sub>2</sub>-induced cell damage.

## Materials and Methods

### Materials

*S. coreanum* was collected along the coast of Jeju Island in South Korea. The sample was washed three times with tap water to remove salt, epiphytes, and sand attached to the surface of the samples. Finally, the sample was carefully rinsed using fresh water and stored at -20°C. The frozen samples were then lyophilized and homogenized in a grinder prior to extraction. For enzymatic digestion, Celluclast 1.5 LFG, which catalyzes the breakdown of cellulose into glucose, cellobiose, and higher glucose polymers, and Neutrase 0.8 L, an endoprotease, were used. Propidium iodide (PI), 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl tetrazolium bromide (MTT), and Hoechst

33342 were all purchased from Sigma (St. Louis, MO, USA). All other chemicals and reagents used were of analytical grade.

### Preparation of enzymatic extracts from *S. coreanum*

Enzymatic extraction of *S. coreanum* was conducted in accordance with the method described by Heo et al. (2005). *S. coreanum* was pulverized into powder with a grinder. Buffer solution (100 mL) was added to 1 g of the dried sample, and then 100 µL (or mg) of each enzyme was added. The enzymatic hydrolysis reaction was performed for 12 h to achieve optimal hydrolysis (Heo et al., 2003). After the reaction, the enzyme was inactivated by heating at 100°C for 10 min. The extracts were clarified by centrifugation (5,000×g, 20 min) to remove non-extracted residue. The resultant mixtures were filtered with Whatman No. 1 filter paper, and the pH of the filtrates was adjusted to pH 7 with 1 M HCl/NaOH. Finally, the enzymatic extracts of *S. coreanum* were lyophilized and maintained at -20°C until use.

### Molecular weight fractionation of enzymatic extracts

Celluclast and Neutrase extracts were fractionated using Millipore's Labscale TFF system (Millipore, Billerica, MA, USA) with ultrafiltration membranes (5, 10 and 30 kDa). The resulting fractions were then collected according to molecular weight (<5, 5-10, 10-30 and >30 kDa) and evaluated with regard to H<sub>2</sub>O<sub>2</sub>-scavenging activity and effects on cell proliferation.

### Total polyphenolic compounds

Total polyphenolic compounds were quantified according to a protocol similar to that described by Chandler and Dodds (1983). *S. coreanum* hydrolysate (1 mL) was mixed in a test tube containing 1 mL of 95% ethanol, 5 mL of distilled water, and 0.5 mL of 50% Folin-Ciocalteu reagent. The mixture was then allowed to react for 5 min, after which 1 mL of 5% Na<sub>2</sub>CO<sub>3</sub> was added. The mixture was then placed in a dark room for 1 h, and the absorbance was measured at a wavelength of 725 nm in a UV-vis spectrophotometer (Opron 3000 Hanson Tech. Co., Ltd., Korea). A gallic acid standard curve was prepared to calculate the polyphenolic content.

### ABTS radical scavenging assay

Hydrogen peroxide scavenging activity was determined by a colorimetric assay according to the method of Müller (1985). Phosphate buffer (100 µL,

0.1 M, pH 5.0) and sample solution were mixed in a 96-microwell plate. Hydrogen peroxide (20  $\mu$ L, 10 mM) was added to the mixture, which was then incubated for 5 min at 37°C. After incubation, 30  $\mu$ L of 1.25 mM ABTS and 30  $\mu$ L of peroxidase (1 unit/mL) were added to the mixture, and it was then incubated for 10 min at 37°C. The absorbance was read with an ELISA plate reader at 405 nm.

### Cell culture

Cells of a Chinese hamster lung fibroblast line (V79-4) were maintained at 37°C in an incubator in a humidified atmosphere at 5% CO<sub>2</sub>. The cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% heat-inactivated fetal calf serum, streptomycin (100 mg/mL), and penicillin (100 unit/mL).

### Assessment of cell viability

Cell viability was then estimated using the MTT assay, a test of metabolic competence predicated on the assessment of mitochondrial performance. It is a colorimetric assay, dependent on the conversion of yellow tetrazolium bromide to its purple formazan derivative by mitochondrial succinate dehydrogenase in viable cells (Mosmann, 1983). The cells were seeded in 96-well plates at a concentration of  $1.0 \times 10^5$  cells/mL. After 16 h, the cells were treated with enzymatic extracts at different concentrations (25, 50 and 100  $\mu$ g/mL). Then, 10  $\mu$ L of H<sub>2</sub>O<sub>2</sub> (1 mM) was added to the cell culture medium and incubated for 24 h at 37°C. MTT stock solution (50  $\mu$ L; 2 mg/mL) was then added to each well to a total reaction volume of 200  $\mu$ L. After 4 h of incubation, the plates were centrifuged (800 $\times$ g, 5 min), and the supernatants were aspirated. The formazan crystals in each well were dissolved in 150  $\mu$ L of dimethylsulfoxide (DMSO), and the absorbance was measured with an ELISA plate reader at 540 nm. Relative cell viability was evaluated in accordance with the quantity of MTT converted to the insoluble formazan salt. The optical density of the formazan generated in the control cells was deemed to represent 100% viability. The data are expressed as mean percentages of viable cells versus the respective control.

### Nuclear staining with Hoechst 33342

Nuclear morphology of cells was evaluated using the cell-permeable DNA dye, Hoechst 33342. Cells with homogeneously stained nuclei were considered viable, whereas the presence of chromatin condensation and/or fragmentation was considered indicative of apoptosis (Gschwind and Huber, 1995; Lizard et

al., 1995). The V79-4 cells were placed in 24-well plates at a concentration of  $1.0 \times 10^5$  cells/mL. At 16 h after plating, the cells were treated with various concentrations of the sample fractions and further incubated for 1 h prior to exposure to H<sub>2</sub>O<sub>2</sub> (1 mM). After 24 h, 1.5  $\mu$ L of Hoechst 33342 (stock, 10 mg/mL), a DNA-specific fluorescent dye, was added to each well, followed by a 10 min incubation at 37°C. The stained cells were then observed under a fluorescence microscope equipped with a CoolSNAP-Pro color digital camera to examine the degree of nuclear condensation.

### Comet assay analysis

The cell suspension was mixed with 75  $\mu$ L of 0.5% low-melting-point agarose (LMPA) and added to slides precoated with 1% normal-melting-point agarose (NMPA). The slides were kept at 4°C for 10 min, then covered with another 75  $\mu$ L of 0.5% LMPA and kept at 4°C for 40 min to solidify the agarose. The slides were then immersed in lysis solution (2.5 M NaCl, 100 mM EDTA, 10 mM Tris, 1% sodium lauryl sarcosine, 1% Triton X-100) at 4°C for 1 h. The slides were next placed in an electrophoresis tank containing 300 mM NaOH and 10 mM Na<sub>2</sub>EDTA (pH 13.0) for 40 min for DNA unwinding. For electrophoresis of the DNA, an electric current of 25 V/300 mA was applied at 4°C for 20 min. The slides were washed three times with a neutralizing buffer (0.4 M Tris, pH 7.5) at 4°C for 5 min and then treated with ethanol for another 5 min before staining with 50  $\mu$ L of ethidium bromide (20  $\mu$ g/mL). Measurements were made with an image analysis system (Kinetic Imaging, Komet 5.0, UK) and a fluorescence microscope (LEICA DMLB, Germany) to determine the percentage of fluorescence in the tail (tail intensity, TI; 50 cells from each of two replicate slides).

### Flow cytometry analysis

Flow cytometry analyses were conducted to determine the proportion of apoptotic sub-G<sub>1</sub> hypodiploid cells (Nicoletti et al., 1991). The V79-4 cells were placed in 6-well plates at a concentration of  $1.0 \times 10^5$  cells/mL, and 16 h after plating, the cells were treated with various concentrations of the sample fraction (100  $\mu$ g/mL). After a further 1 h incubation, H<sub>2</sub>O<sub>2</sub> (1 mM) was added to the culture. After 24 h, the cells were harvested at the indicated times and fixed for 30 min in 1 mL of 70% ethanol at 4°C. The cells were then washed twice with PBS and incubated for 30 min in the dark in 1 mL of PBS containing 100  $\mu$ g PI and 100  $\mu$ g RNase A at 37°C.

Flow cytometric analysis was conducted using a FACSCalibur flow cytometer (Becton Dickinson, San Jose, CA, USA). Effects on the cell cycle were determined by measuring changes in the percentage of cell distribution at each phase of the cell cycle and were assessed by histograms generated by the Cell Quest and Mod-Fit computer programs (Wang et al., 1999).

## Results and Discussion

Recently, due to the relatively limited resources of the land, many researchers have focused on natural marine resources with various biological activities. Reactive oxygen species (ROS) form as natural by-products from normal oxygen metabolism and perform important functions in cell signaling. Cells are normally able to defend themselves against ROS damage by the use of antioxidant enzymes, including superoxide dismutases and catalases (Heo and Jeon, 2008). However, an imbalance between ROS and antioxidant defense mechanisms can result in oxidative modification of the cellular membrane or intracellular molecules (Halliwell et al., 1992; El-Habit et al., 2000). Recently, interest has been increasing in the therapeutic potential of natural plants as sources of antioxidants for the reduction of such free radical-induced tissue injuries. It has been suggested that many plants may possess therapeutically useful antioxidant activities (Kim et al., 2006). In this study, we focused on natural water-soluble antioxidants isolated from *S. coreanum*, which were prepared via enzymatic hydrolysis using different carbohydrate-degrading enzymes and proteases. We then investigated the antioxidant effects of these *S. coreanum* enzymatic extracts after the administration of H<sub>2</sub>O<sub>2</sub> treatment.

In a previous study, we preliminarily demonstrated that Celluclast and Neutrased extracts exhibited the strongest H<sub>2</sub>O<sub>2</sub>-scavenging activities among five carbohydrases (Viscozyme, Celluclast, Termamyl, Ultraflo and AMG) and five proteases (Kojizyme, Alcase, Flavourzyme Protamex and Neutrased) extracts (Athukorala et al., 2003). Thus, the Celluclast and Neutrased extracts were selected for further study. To characterize the molecular weight of the active compounds, the extracts were passed through ultrafiltration membranes. Then, the fractions (unfractionated sample, <5, 5-10, 10-30 and >30 kDa) at different concentrations (0.1, 0.25 and 0.5 mg/mL) were investigated separately with regard to their H<sub>2</sub>O<sub>2</sub> scavenging activities. The 5-10 kDa fraction of the Celluclast and Neutrased extracts showed the most

potent H<sub>2</sub>O<sub>2</sub>-scavenging activity among the fractions, with IC<sub>50</sub> values of 0.177 and 0.182 mg/mL, respectively (Table 1). Previous reports have examined high H<sub>2</sub>O<sub>2</sub> scavenging ability in seaweeds (Athukorala et al., 2003; Siriwardhana et al., 2004). In particular, brown seaweeds has been shown to exhibit profound H<sub>2</sub>O<sub>2</sub>-scavenging and DPPH radical-scavenging activities (Heo et al., 2005; Heo and Jeon, 2008; Kim et al., 2006).

The fraction that had the lowest IC<sub>50</sub> of the tested extracts was selected for use in the next set of experiments. Meanwhile, all of the molecular weight fractions were subjected to total polyphenolic assays to determine the polyphenolic contents of each extract. The total polyphenolic amount in each extract is shown in Table 1. The highest polyphenolic content was found in the 5-10 kDa fraction of Celluclast (13.98 mg/100 mL) and Neutrased (15.58 mg/100 mL), whereas the lowest content was seen in the <5 kDa fraction of Celluclast (4.2 mg/100 mL) and Neutrased (3.38 mg/100 mL). As reported previously (Siriwardhana et al., 2004), a correlation exists between total polyphenolic content and radical-scavenging activity of seaweed extracts; this was confirmed in the present study. Many researchers have reported a positive correlation between ROS-scavenging activity and total phenolic compound content (Kim et al., 2006; Ahn et al., 2007, Oki et al., 2002). As the 5-10 kDa fraction of the Celluclast and Neutrased extracts generated in this study showed such high H<sub>2</sub>O<sub>2</sub>-scavenging activity, this fraction was further evaluated with regard to its protective effects against the cellular damage induced by H<sub>2</sub>O<sub>2</sub>. The protective effect of the 5-10 kDa fraction of Celluclast and Neutrased extracts on H<sub>2</sub>O<sub>2</sub>-induced cellular damage to the V79-4 cell line is shown in Fig. 1. In this study, both of the tested extracts controlled H<sub>2</sub>O<sub>2</sub>-induced

Table 1. IC<sub>50</sub> values of H<sub>2</sub>O<sub>2</sub> scavenging activity and total phenolic contents of the different molecular weight fractions of *S. coreanum* Celluclast and Neutrased extract

Treated enzymes	Molecular weight fractions	IC <sub>50</sub> values (mg/mL)	Total phenolic contents (mg/100 mL)
Celluclast	Original extract	0.545	8.69
	>30 kDa	0.379	11.80
	10-30 kDa	0.367	9.82
	5-10 kDa	0.182	13.98
	<5 kDa	0.932	3.38
Neutrased	Original extract	0.536	9.45
	>30 kDa	0.275	13.62
	10-30 kDa	0.251	15.38
	5-10 kDa	0.177	15.58
	<5 kDa	0.876	4.20




Fig. 1. Protective effects of 5-10 kDa fractions of the Celluclast and Neutrased extracts from *S. coreanum* against H<sub>2</sub>O<sub>2</sub>-induced oxidative damage in V79-4 cells. Experiments were conducted in triplicate, and data are expressed as the average percentage change from control  $\pm$  S.D. (♦) 5-10 kDa fractions of Celluclast extract and (■) 5-10 kDa fractions of Neutrased extract.

cellular damage in a dose-dependent manner. The addition of H<sub>2</sub>O<sub>2</sub> to the cell culture medium without enzymatic extracts resulted in a 46% cell survival rate, but the addition of extracts coupled with exposure to H<sub>2</sub>O<sub>2</sub> resulted in a dose-dependent increase in cell survival rates. In particular, the highest cell viability, 72% and 76%, was recorded with the 5-10 kDa fraction of Celluclast and Neutrased extract, respectively, at concentrations of 100 µg/mL.

Due to the profound antioxidant activity shown by the 5-10 kDa fraction, the fraction was evaluated further with regard to its ability to protect against DNA damage. DNA damage is one of the most sensitive biological markers for the evaluation of oxidative stress and illustrates the imbalance between free radical generation and the efficiency of the antioxidant system (Gutteridge, 1995; Kassie et al., 2000). Hoechst 33342 dye specifically stains DNA




Fig. 2. Protective effect of 5-10 kDa fractions of the Celluclast and Neutrased extracts from *S. coreanum* against H<sub>2</sub>O<sub>2</sub>-induced apoptosis in V79-4 cells. The cells were treated with 5-10 kDa fractions of the Celluclast and Neutrased extracts at concentrations of 50 and 100 µg/mL. After 1 h, 1 mM of H<sub>2</sub>O<sub>2</sub> was added to the plate. After incubation of 24 h, the apoptotic body formation was observed under a fluorescence microscope after Hoechst 33342 staining. (A) Negative control; (B) positive control (cells treated with 1 mM H<sub>2</sub>O<sub>2</sub>); (C) cells treated with 50 µg/mL 5-10 kDa fraction of Celluclast extract + 1 mM H<sub>2</sub>O<sub>2</sub>; (D) cells treated with 100 µg/mL 5-10 kDa fraction of Celluclast extract + 1 mM H<sub>2</sub>O<sub>2</sub>; (E) cells treated with 50 µg/mL 5-10 kDa fraction of Neutrased extract + 1 mM H<sub>2</sub>O<sub>2</sub>; (F) cells treated with 100 µg/mL 5-10 kDa fraction of Neutrased extract + 1 mM H<sub>2</sub>O<sub>2</sub>.

and has been used extensively for the detection of nuclear shrinkage, including chromatin condensation, nuclear fragmentation, and the appearance of apoptotic bodies, all of which are indicative of apoptosis (Heo and Jeon 2009). As shown in Fig. 2, the negative control, which contained no extracts or H<sub>2</sub>O<sub>2</sub>, harbored intact nuclei (Fig. 2A), and the positive control (cells treated with 1 mM H<sub>2</sub>O<sub>2</sub>) showed a significant level of nuclear fragmentation, indicative of apoptosis (Fig. 2B). However, addition of the 5-10 kDa fractions with H<sub>2</sub>O<sub>2</sub> reduced the number of apoptotic bodies (Fig. 2C-F), suggesting the ability of the 5-10 kDa fractions to protect against nuclear fragmentation in H<sub>2</sub>O<sub>2</sub> challenge and indicating that the cells were protected against oxidative stress-associated cellular injury. The protective effects of 5-10 kDa fractions on cell damage were also confirmed via a comet assay, which is a rapid and sensitive fluorescent microscopic method for the detection of primary DNA damage at the individual cellular level and is used to evaluate the genotoxicity of test substances (Bagchi et al., 2000). The percentage of the fluorescence tail DNA intensity of V79-4 cells subjected to 30 min of treatment with the negative control differed significantly from that of the positive control (H<sub>2</sub>O<sub>2</sub> preincubated in PBS). This increase in H<sub>2</sub>O<sub>2</sub>-induced cell damage was inhibited significantly, in a dose-dependent fashion, by preincubating the H<sub>2</sub>O<sub>2</sub> and 5-10 kDa fractions together with the negative cells (Fig. 3). The inhibitory activities of the 5-10 kDa fraction of the Celluclast and Neutrased extracts on DNA damage were 67 and 68%, respectively, at concentrations of 100 µg/mL. We also examined photomicrographs of different DNA migration profiles after treatment with different concentrations of the 5-10 kDa fractions of Celluclast and Neutrased extracts (Fig. 4). In the positive control (cells treated only with H<sub>2</sub>O<sub>2</sub>), the DNA was severely and systemically damaged, and the quantities of tail DNA were significantly increased, compared with those of the negative control. However, when we added the 5-10 kDa fraction of Celluclast and Neutrased extracts to the cells, we determined that the amounts of tail DNA were increasingly reduced with increasing concentrations of the 5-10 kDa fractions of the Celluclast and Neutrased extracts. Reactive oxygen species such as H<sub>2</sub>O<sub>2</sub> are highly reactive and are capable of inflicting damage to a variety of biological macromolecules, including DNA, RNA, proteins, and lipids (Yan et al., 1999). In addition to our morphological evaluations, the protective effects of the 5-10 kDa fraction were confirmed by flow cytometry. DNA content analyses conducted following the H<sub>2</sub>O<sub>2</sub>




Fig. 3. Inhibitory effect of 5-10 kDa fractions of the Celluclast and Neutrased extracts from *S. coreanum* on H<sub>2</sub>O<sub>2</sub>-induced DNA damage. Cells were treated with 5-10 kDa fractions at concentrations of 25, 50, and 100 µg/mL. After 30 min, 1 mM H<sub>2</sub>O<sub>2</sub> was added and incubated for 15 min. The percentage of cellular DNA damage was detected by comet assay. (A) Inhibitory effect of 5-10 kDa fraction of the Celluclast extract on H<sub>2</sub>O<sub>2</sub>-induced DNA damage. (B) Inhibitory effect of 5-10 kDa fraction of the Neutrased extract on H<sub>2</sub>O<sub>2</sub>-induced DNA damage. (□) % Fluorescence in tail; (●) inhibitory effect of cell damage.

treatment of V79-4 cells revealed an increase in the proportion of cells with sub-G<sub>1</sub> DNA content to 18.5% (Fig. 5). This result indicated that apoptosis was induced by the added H<sub>2</sub>O<sub>2</sub>. However, the cells that were pretreated with the 5-10 kDa fraction of the Celluclast and Neutrased extracts showed significantly reduced sub-G<sub>1</sub> DNA contents (2.68 and 1.8%, respectively). Cells in the human body are basically under continuous attack by physical agents such as solar radiation, a variety of chemical compounds, and reactive oxygen species, which are natural by-products of metabolic processes (Kim et al., 2006). These substances can all induce DNA damage. If this DNA damage remains unrepaired, a cascade of biological consequences can be initiated within the cell population. Unresolved DNA damage can also




Fig. 4. Photomicrographs of DNA damage and migration observed under 5-10 kDa fractions of the Celluclast and Neutrased extracts from *S. coreanum* where the tail moments were decreased. (A) Negative control; (B) positive control (cells treated with 1 mM  $H_2O_2$ ); (C) cells treated with 25  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Celluclast extract + 1 mM  $H_2O_2$ ; (D) cells treated with 50  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Celluclast extract + 1 mM  $H_2O_2$ ; (E) cells treated with 100  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Celluclast extract + 1 mM  $H_2O_2$ ; (F) cells treated with 25  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Neutrased extract + 1 mM  $H_2O_2$ ; (G) cells treated with 50  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Neutrased extract + 1 mM  $H_2O_2$ ; (H) cells treated with 100  $\mu\text{g}/\text{mL}$  5-10 kDa fraction of Neutrased extract + 1 mM  $H_2O_2$ .




Fig. 5. Effects of the 5-10 kDa fraction on cell cycle pattern and apoptotic cell proportion in V79-4 cells with H<sub>2</sub>O<sub>2</sub> (1 mM) via flow cytometric analysis. (A) Control; (B) H<sub>2</sub>O<sub>2</sub>-treated; (C) cells treated with 100 µg/mL 5-10 kDa fraction of Celluclast extract + 1 mM H<sub>2</sub>O<sub>2</sub>; (D) cells treated with 100 µg/mL 5-10 kDa fraction of Neutrased extract + 1 mM H<sub>2</sub>O<sub>2</sub>.

promote cancer development via several mechanisms (Nardella et al., 1996). The antioxidative activities of *E. cava*, as with other brown seaweeds, may be primarily attributable to carotenoid pigments (fucoxanthin), polyphenols (phlorotannins), vitamins (vitamins C, E), sulfated polysaccharides (laminarin, fucoidans), or their breakdown products, as well as proteins or their breakdown products (Yan et al., 1999; Nardella et al., 1996; Peña-Ramos and Xiong, 2002). In this enzymatic hydrolysis, polyphenolics, vitamin C, sulfated polysaccharides, and proteins, among the many materials with antioxidative potential, may be considered among the most effective due to their native water-soluble nature.

### Acknowledgements

This research was supported by the grant from the Marine Bioprocess Research Center of the Marine Bio 21 Center, funded by Ministry of Marine Affairs and Fisheries, Republic of Korea.

### References

- Ahn G, Kim KN, Cha SH, Song CB, Lee JH, Heo MS, Yeo IK, Lee NH, Jee YH, Kim JS, Heu MS and Jeon YJ. 2007. Antioxidant activities of phlorotannins purified from *Ecklonia cava* on free radical scavenging using ESR and H<sub>2</sub>O<sub>2</sub>-mediated DNA damage. *Eur Food Res Technol* 226, 71-79.
- Ahn G, Park E, Kim DS, Jeon YJ, Shin T, Park JW, Woo HC, Lee KW and Jee Y. 2008. Anti inflammatory effects of enzymatic extract from *Ecklonia cava* on TPA-induced ear skin edema. *Food Sci Biotechnol* 17, 745-750.
- Ames B. 1998. Micronutrients prevent cancer and delay aging. *Toxicol Lett* 102, 5-18.
- Athukorala Y, Lee KW, Park EJ, Heo MS, Yeo IK, Lee YD and Jeon YJ. 2003. Reduction of lipid peroxidation and H<sub>2</sub>O<sub>2</sub>-mediated DNA damage by a red alga (*Grateloupia filicina*) methanolic extract. *J Sci Food Agri* 85, 251-65.
- Athukorala Y and Jeon YJ. 2005. Screening for angiotensin I-converting enzyme inhibitory activity of *Ecklonia cava*. *J Food Sci Nutr* 10, 134-139.
- Bagchi D, Bagchi M, Stohs SJ, Das DK, Ray SD, Kuszynski CA, Joshi SS and Pruess HG. 2000. Free radicals and grape seed proanthocyanidin extract: importance in human health and disease prevention. *Toxicology* 148, 187-197.
- Chandler SF and Dodds JH. 1983. The effect of phosphate, nitrogen and sucrose on the production of phenolics and solasidine in callus cultures of *Solanum laciniatum*. *Plant Cell Rep* 2, 105-108.



- Cox DA and Choen ML. 1996. Effects of oxidized low density lipoproteins on vascular contraction and relaxation. *Pharmacol Rev* 48, 3-9.
- El-Habit OHM, Saada HN and Azab KS. 2000. The modifying effect of b-carotene on gamma radiation-induced elevation of oxidative reactions and genotoxicity in male rats. *Mutat Res* 466, 179-186.
- Gonzalez AG, Darias V and Estevez E. 1982. Chemotherapeutic activity of polyhalogenated terpenes from Spanish algae. *Planta Med* 44, 44-46.
- Gschwind M and Huber G. 1995. Apoptotic cell death induced by b-amyloid 1-42 peptide is cell type dependent. *J Neurochem* 65, 292-300.
- Gutteridge JMC. 1995. Lipid peroxidation and antioxidants as biomarker of tissue damage. *Clin Chem* 41, 1819-1828.
- Halliwell B and Aruoma OI. 1991. DNA damage by oxygen-derived species. *FEBS Lett* 281, 9-19.
- Halliwell B, Gutteridge JMC and Cross CE. 1992. Free radicals, antioxidants, and human disease: where are we now. *J Lab Clin Med* 119, 598-620.
- Heo SJ, Ko SC, Kang SM, Kang HS, Kim JP, Kim SH, Lee KW, Cho MG and Jeon YJ. 2008. Cytoprotective effect of fucoxanthin isolated from brown algae *Sargassum siliquastrum* against H<sub>2</sub>O<sub>2</sub>-induced cell damage. *Eur Food Res Technol* 228, 145-151.
- Heo SJ and Jeon YJ. 2009. Evaluation of diploretol isolated from *Ishige okamurae* for radical scavenging activity and its protective effect against H<sub>2</sub>O<sub>2</sub>-induced cell damage. *Process Biochem* 44, 412-418.
- Heo SJ and Jeon YJ. 2008. Radical scavenging capacity and cytoprotective effect of enzymatic digests of *Ishige okamurae*. *J Appl Phycol* 20, 1087-1095.
- Heo SJ, Jeon YJ, Lee JH, Kim HT and Lee KW. 2003. Antioxidant effect of enzymatic hydrolyzate from a Kelp, *Ecklonia cava*. *Algae* 18, 341-347.
- Heo SJ, Park EJ, Lee KW and Jeon YJ. 2005. Antioxidant activities of enzymatic extracts from brown seaweeds. *Bioresour Technol* 96, 1613-1623.
- Kashiwagi M, Mynderse JS, Moore RE and Norton TR. 1980. Antineoplastic evaluation of pacific basin marine algae. *J Pharm Sci* 69, 735-738.
- Kassie F, Parzefall W and Knasmuller S. 2000. Single cell gel electrophoresis assay: a new technique for human biomonitoring studies. *Mutat Res* 463, 13-31
- Kim EK, Lee SJ, Lim BO, Jeon YJ, Song MD, Park TK, Lee KH, Kim B, Lee SR, Moon SH, Jeon BT and Park PJ. 2008. Antioxidative and neuroprotective effects of enzymatic extracts from leaves of *Perilla frutescens* var. japonica. *Food Sci Biotechnol* 17, 279-286.
- Kim KN, Heo SJ, Song CB, Lee J, Heo MS, Yeo IK, Kang KA, Hyun JW and Jeon YJ. 2006. Protective effect of *Ecklonia cava* enzymatic extracts on hydrogen peroxide-induced cell damage. *Process Biochem* 41, 2393-2401.
- Ko SC, Kang SM, Ahn G, Yang HP, Kim KN and Jeon YJ. 2009. Antioxidant activity of enzymatic extracts from *Sargassum coreanum*. *J Kor Soc Food Sci Nutr* Submitted.
- Kosovel V, Avanzini A, Scarcia V and Furlani A. 1988. Algae as possible sources of antitumoral agents. Preliminary evaluation of the *in vitro* cytostatic activity of crude extracts. *Pharmacol Res Commun* 20, 27-31.
- Kotake-Nara E, Asai A and Nagao A. 2005. Neoxanthin and fucoxanthin induce apoptosis in PC-3 human prostate cancer cells. *Cancer Lett* 220, 75-84.
- Lee SH, Karawita R, Affan A, Lee JB, Lee KW, Lee BJ, Kim DW and Jeon YJ. 2009. Potential of benthic diatoms *Achnanthes longipes*, *Amphora coffeaeformis* and *Navicula* sp. (Bacillariophyceae) as antioxidant sources. *Algae* 24, 47-55.
- Lindenschmidt RC, Trika AF, Guard ME and Witschi HP. 1986. The effect of dietary butylated hydroxyl toluene on liver and colon tumor development in mice. *Toxicol* 38, 151-160.
- Lizard G, Fournel S, Genestier L, Dhedin N, Chaput C, and Flacher M. 1995. Kinetics of plasma membrane and mitochondrial alterations in the cells undergoing apoptosis. *Cytometry* 21, 275-283.
- Mayer AMS and Hmann MT. 2004. Marine pharmacology in 2000: marine compounds with antibacterial, anti-coagulant, antifungal, anti-inflammatory, antimalarial, antiplatelet, antiprotozoal, anti-tuberculosis, and antiviral activities; affecting the cardiovascular, immune and nervous systems and other miscellaneous mechanisms of action. *Mar Biotechnol* 6, 37-52.
- Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Meth* 65, 55-63.
- Müller HE. 1985. Detection of hydrogen peroxide produced by microorganism on ABTS-peroxide medium. *Zentralbl Bakteriol Mikrobiol Hyg* 259, 151-158.
- Nagayama K, Iwamura Y, Shibata T, Hirayama I and Nakamura T. 2002. Bactericidal activity of phlorotannins from the brown alga *Ecklonia kurome*. *J Antimicrob Chemother* 50, 889-893.
- Nardella A, Chaubet F, Boisson-vidal C, Blondin C, Durand P and Jozefonvicz J. 1996. Anticoagulant low molecular weight fucans produced by radical process and ion exchange chromatography of high molecular weight fucans extracted from the brown seaweed *Ascophyllum nodosum*. *Carbohydr Res* 289, 201-208.

- Nicoletti I, Migliorati G, Pagliacci MC, Grignani F and Riccardi C. 1991. A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J Immunol Meth* 139, 271-279.
- Oki T, Masuda M, Furuta S, Nishibia Y, Terahara N and Suda I. 2002. Involvement of anthocyanins and other phenolic compounds in radical-scavenging activity of purple-fleshed sweet potato cultivars. *J Food Sci* 67, 1752-1756.
- Peña-Ramos EA and Xiong YL. 2002. Antioxidant activity of soy protein hydrolysates in a liposomal system. *J Food Sci* 67, 2952-2956.
- Sherwin ER. 1990. Antioxidants. In *Food Additives*. Branen AI, Dvidson PM, Salminen S, eds. Marcel Dekker, New York.
- Siriwardhana N, Lee KW, Kim SH, Ha JH, Jeon YJ. 2003. Antioxidant activity of *Hizikia fusiformis* on reactive oxygen species scavenging and lipid peroxidation inhibition. *Food Sci Technol Int* 9, 339-346.
- Siriwardhana N, Lee KW, Kim SH, Ha JW, Park GT, Jeon YJ. 2004. Lipid peroxidation inhibitory effects of *Hizikia fusiformis* methanolic extract on fish oil and linoleic acid. *Food Sci Technol Int* 10, 65-72.
- Wang XW, Zhan Q, Coursen JD, Khan MA, Kontny HU and Yu L. 1999. GADD45 induction of a G2/M cell cycle checkpoint. *Proc Natl Acad Sci USA* 96, 3706-3711.
- Yan X, Chuda Y, Suzuki M and Nagata T. 1999. Fucoxanthin as the major antioxidant in *Hizikia fusiformis*, a common edible seaweed. *Biosci Biotechnol Biochem* 63, 605-607.

(Received 19 January 2010; Revised 22 February 2010;  
Accepted 15 March 2010)